PROFESSOR CHENGDAO LI (Orcid ID : 0000-0002-9653-2700)

Article type : Original Article

A global barley panel revealing genomic signatures of breeding in modern Australian cultivars

Camilla Beate Hill¹, Tefera Tolera Angessa¹, Xiao-Qi Zhang¹, Kefei Chen^{2,3}, Gaofeng Zhou^{1,2}, Cong Tan¹, Penghao Wang¹, Sharon Westcott², Chengdao Li^{*1,2}

¹Western Crop Genetics Alliance, Agricultural Sciences, College of Science, Health, Engineering and Education, Murdoch University, Murdoch, WA

²Agriculture and Food, Department of Primary Industries and Regional Development, South Perth, WA, Australia

³Statistics for the Australian Grains Industry (SAGI) West, Faculty of Science and Engineering, Curtin University, Bentley, WA, Australia

* Corresponding author

Email: C.Li@murdoch.edu.au

Running title

Genomic signatures of barley breeding

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/TPJ.15173](https://doi.org/10.1111/TPJ.15173)

This article is protected by copyright. All rights reserved

Keywords

barley, breeding, population analysis, selective sweeps, genetic diversity, phenology, next-generation sequencing, GWAS, gene ontology, genotyping-by-sequencing

Summary

The future of plant cultivar improvement lies in the evaluation of genetic resources from currently available germplasm. Today's gene pool of crop genetic diversity has been shaped during domestication and more recently by breeding. Recent efforts in plant breeding have been aimed at developing new and improved varieties from poorly adapted crops to suit local environments. However, the impact of these breeding efforts is poorly understood. Here, we assess the contributions of both historical and recent breeding efforts to local adaptation and crop improvement in a global barley panel by analysing the distribution of genetic variants with respect to geographic region or historical breeding category. By tracing the impact breeding had on the genetic diversity of barley released in Australia, where the history of barley production is relatively young, we identify 69 candidate regions within 922 genes that were under selection pressure. We also show that modern Australian barley varieties exhibit 12% higher genetic diversity than historical cultivars. Finally, field-trialling and phenotyping for agriculturally relevant traits across a diverse range of Australian environments suggests that genomic regions under strong breeding selection and their candidate genes are closely associated with key agronomic traits. In conclusion, our combined dataset and germplasm collection provide a rich source of genetic diversity that can be applied to understanding and improving environmental adaptation and enhanced yields.

Introduction

The diversity of the existing genetic pool for commercially important plant species has been shaped during plant domestication, human migration, varietal selection processes and, more recently, breeding. However, there is concern that breeding efforts have eroded genetic variation, thereby resulting in a narrow range of genotypes in the current gene pools of domesticated crops (van de Wouw *et al.*, 2010a, b). Although regionally adapted landraces and wild relatives represent the most diverse germplasm reservoirs, the introgression of desirable alleles into elite germplasm used by breeders—whilst minimising the introduction of other genes from the wild germplasm that might reduce the agronomic fitness of the elite cultivar—has been challenging and time consuming (Dempewolf *et al.*, 2017). As a result of these challenges and the often limited availability of high-density markers and detailed information for key adaptive traits, the high degree of genetic diversity in wild crop relatives has been poorly exploited.

Changes in global climate and short-term variations in growing environments pose unprecedented challenges to maintain and further enhance crop yields. Among other effects, climate change substantially alters phenological cycles, thereby posing a significant challenge to growers, who must modify crop management practices such as sowing dates in order to achieve optimal flowering times (Hill *et al.*, 2016). Conservation and maintenance of current crop genetic diversity for future breeding of varieties is particularly important to help mitigate future adverse impacts of climate change on crop production.

While climate change threatens the supply of agricultural products, global demand is increasing for resource-intensive foods including meat and dairy, and alcoholic beverages including beer (Gandi *et al.*, 2014; Colen *et al.*, 2016). Barley (*Hordeum vulgare* L.) is a globally important and versatile crop used for both livestock feed and brewing malts. Despite of its large and repeat-rich genome (~5.1 Gb) distributed over seven chromosomes, it is a widely utilized diploid cereal model for genetic studies in the *Triticeae*, a botanical tribe which includes polyploid bread wheat and rye. Since the barley reference genome became publicly available (Mascher *et al.*, 2017), several genetic studies have explored the origin, domestication, and geographic spread of modern barley (Bustos-Korts *et al.*, 2019; Dreissig *et al.*, 2019), and showed that the Fertile Crescent is the main centre of domestication and genetic diversity. Population genetic studies have examined a variety of aspects of barley genetic variation on a global scale, and have identified a striking degree of variability in traits related to flowering time, grain yield, and tolerance to abiotic and biotic factors (Merchuk-Ovnat *et al.*, 2018; Phan *et al.*, 2018; He *et al.*, 2019; Hill *et al.*, 2019a, b). This suggests that current barley germplasm resources might be harnessed to meet future challenges imposed by climate change.

Although it was only in the late 18th century that barley was first introduced to Australia (Friedt *et al.*, 2011), it is currently one of the world's largest barley producers (<http://faostat.fao.org>). The first introduced cultivars were poorly adapted, late-maturing, European barleys, and were susceptible to the hot, dry conditions typical of Australia. It was not until the 1960s that first Australian breeding programmes were established that specifically targeted different barley-producing regions in Western Australia, South Australia, Victoria, New South Wales, Queensland and Tasmania. Only then breeding material from North Africa and North America was introduced to improve disease resistance against powdery mildew and cereal cyst nematodes as well as phenological adaptation of Australian barley varieties, which was further aided by the

development of molecular marker technologies in the 1980s. However, the genetic impact of these breeding efforts and the extent of genetic diversity within current cultivated barley germplasm reservoirs is poorly understood.

In our previous studies (Hill *et al.*, 2019a,b), we applied a target-capture method to detect polymorphisms in 174 phenology-related genes of 895 barley accessions and identified target genes linked with phenology and grain yield traits. Analysis of allele diversity within the targeted 174 genes showed that modification of phenology genes was a key element to breeding new varieties for the challenging growing environments characteristic for the Australian Wheatbelt. In this study, we assessed the genome-wide contributions of both historical and recent breeding efforts towards local adaptation and crop improvement, using Australian barley as a model for the profound impact that breeding efforts can have on developing a previously poorly adapted crop suitable to local environments. In order to accomplish this, we analysed the distribution of a genome-wide set of over 33,000 genetic variants in terms of geographic region and historical breeding category and determined the genomic regions under selection and underlying candidate genes that may have been shaped by selective breeding. Associations identified between both known and novel genes and agronomic traits demonstrate the value of our global barley panel for both fundamental and applied studies. Lastly, both our selective sweep analyses and our genome-wide association studies (GWASs) highlight targets for future gene and functional allele discovery.

Results

Barley genomic diversity

To examine the origins and patterns of genetic diversity within the currently available barley cultivar gene pool, we assembled a global panel of 632 genotypes to represent barley genotypes from major global barley breeding programmes, including both historical cultivars and modern cultivars from 43 countries (S1 File). As the barley genome is quite large (~5.1 Gb), and the genome consists of >80% mobile and repeated elements, whole-genome re-sequencing is a cost-intensive approach to comprehensively catalogue genetic diversity. To circumvent this limitation, we used a combination of three sequencing methods to capture variation in and around the gene-

containing regions of the 632 barley genotypes. The panel of 632 genotypes geographically diverse barley cultivars was genotyped using target capture (Hill *et al.*, 2019a, b), low-coverage whole-genome sequencing (WGS), and genotyping-by-sequencing (GBS) by Diversity Arrays Technology (DArTseq). In total, 15,328 single-nucleotide polymorphisms (SNPs) and insertions and deletions (InDels) were detected via low-coverage WGS, 4,260 SNPs via target-enrichment sequencing, and 18,551 SNPs via DArTseq were distributed across 5,171 barley genes. The mapping of genetic markers from all 632 genotypes onto the current barley reference genome sequence (Mascher *et al.*, 2017) (IBSC v2) revealed 38,139 high-confidence genetic variants. A total of 33,486 filtered genetic markers (32,645 SNPs and 841 InDels) with a minor allele frequency (MAF) > 0.01 were used in the present study (S1 Table). As expected for target capture and DArTseq analyses (which focused on actively transcribed genes), the distribution of genetic variants across the seven chromosomes exhibited a visible gradient across chromosome compartments, from distal regions with relatively high gene density to pericentromeric regions with fewer genes (S1 Fig).

For a more detailed analysis of the genomic changes that have occurred during the history of Australian barley breeding, Australian genotypes were separated into four historical subgroups based on release date: Category A (historically relevant cultivars used in 20th century breeding programmes in Australia, released between 1903 and 1998), Category B (modern cultivars with specific regional adaptations that were released between 1999 and 2005 as a result of focused barley breeding programmes across different Australian states), Category C (most recently released elite cultivars, released between 2006 and 2019); and Category D (unreleased breeding and research lines). A detailed description of the Australian varieties, including the year of release and breeder, is provided in S1 File.

The polymorphism information content (PIC) was estimated to evaluate the frequency of nucleotide variants across the entire barley population, as well as in subpopulations based on geographical regions or historical subgroups of Australian barley. From the genetic variant data for all 632 accessions in the barley panel, the PIC was estimated to be 0.17 (Table 1), although we observed marked differences among different geographical regions (Australia, Africa, Asia, Europe, North America, and South America) and among historical subgroups of Australian barley. Within historical subgroups of Australian cultivars, the observed mean PIC values were slightly higher for varieties released between 2006 and 2019 (CatC) and unreleased research and breeding

lines (CatD) (both 0.16) than for varieties released between 1903 and 1998 (CatA, 0.14) or between 1999 and 2005 (CatB, 0.15).

Population structure within the global collection of domesticated barley varieties

Underlying population structure is known to be a confounding factor in GWASs, particularly for adaptive traits such as flowering time (Liu and Yan, 2019). Known sources of population structure in domesticated barley varieties include the separation of two-row and six-row barleys, which occurred early in domestication (~8,000 years ago), as well as the separation of spring and winter barleys, which accelerated the migration of barley through the modification of the vernalisation requirement and photoperiod response (Malysheva-Otto *et al.*, 2006; Komatsuda *et al.*, 2007). Next, we therefore investigated the population structure of the global barley germplasm collection used in this study using ADMIXTURE (Alexander *et al.*, 2011) to select the optimal number of subpopulations (K), which we predicted was approximately K = 12 (Figs 1a–e).

Phylogenetic trees were constructed based on the genetic distances of the entire population (Figs 1f and g), as well as on the genetic distances of 47 Australian cultivars selected to represent the diversity of germplasm used in Australian barley breeding (Fig 1h) using the Neighbour-joining (NJ) clustering method. Distinct clusters were detected based on geographic location (Fig 1f), row type, and growth habit (Fig 1g). As expected, no clear clustering pattern was observed based on historical subgroups (Fig 1h), as within our Australian barley panel, many of the cultivars that were first to be released are ancestors of modern cultivars. Principal component analysis (PCA) was performed with separation based on row type, growth habit, or geographic region (Fig 2), corroborating the results of phylogenetic analyses. Taken together, our data suggest that three major factors account for the partitioning of diversity within the global barley panel: geographic origin (Asia, Middle East, North America, and South America) (Fig 2a), growth habit (winter vs. spring) (Fig 2b), and row type (six-row vs. two-row) (Fig 2c). However, no clear clustering pattern was observed among European, African, and Australian genotypes, which is likely due to the extensive movement of germplasm from Europe to Australia and extensive use of diverse African lines in Australian and European breeding programmes (Friedt *et al.*, 2011).

To understand the patterns of linkage disequilibrium (LD) between different chromosomes, we calculated r^2 values between pairs of genetic variants for all 632 genotypes, as well as in the four Australian subpopulations. LD was estimated for each subpopulation as a function of physical

distance (Fig 3 and S2 Fig). Genetic marker pairs were sorted into 100-kb bins based on the distance between pairs, and mean r^2 values were estimated for each bin. Long-range LD was more extensive in historical barley cultivars (CatA and CatB) than in more recently released barley cultivars (CatC) owing to the greater extent of allelic association in the early period of barley breeding, thereby confirming the narrow initial gene pool of early breeding programmes (Friedt *et al.*, 2011).

Selection footprints of barley breeding

To explore selection footprints resulting from breeding within the global barley panel, next we investigated genetic diversity parameters in for the whole population, between barley genotypes sourced from different geographic regions, and within Australian subgroups based on release date. We first examined the degree of polymorphism along each chromosome within different geographic region and among historical Australian barley groups by calculating the nucleotide diversity statistic π (Nei *et al.*, 1979) (S3 Fig and S4 Fig). The distribution of nucleotide diversity indicates limited allelic diversity in domesticated barley genotypes and that modern breeding processes had measurably altered overall genetic diversity, which increased during a relatively short breeding period in Australian barley varieties (Fig 4a, S2 Table).

The nucleotide diversity index π varied between barley varieties from the six geographic regions, with the highest and lowest genetic diversities observed in Asian and African barleys, respectively. A gradual increase in nucleotide diversity was detected when comparing historical cultivars (CatA) to the later (CatB) and the most recently released Australian cultivars (CatC). Our results are consistent with a continuous increase in diversity through breeding, as observed between CatA and CatB (representing a ~8.9% higher nucleotide diversity in cultivars released between 1998 and 2005 than in cultivars released between 1903 and 1998). The continuous increase in diversity between groups CatA and CatC (representing a ~12.5% higher nucleotide diversity in cultivars released between 2005 and 2019 than in cultivars released between 1903 and 1998) reflect the increased use of exotic germplasm bred into modern Australian barleys (Friedt *et al.*, 2011) and breeding improvement in early Australian barley breeding programmes. These findings show that despite six decades of intense breeding of Australian barley cultivars, higher genetic variation exists within the current breeding gene pool compared to historical varieties.

We also calculated subpopulation-specific estimates of Tajima's D to compare the average number of pairwise differences and the number of segregating sites between samples within each of our geographic and historical subpopulations in Australia. The sign of Tajima's D provides an interpretation of natural selection, where balancing selection results in a positive, and positive selection results in a negative Tajima's D . Subpopulation-specific estimates of Tajima's D differed extensively, and included both negative and positive values, but with a strong and consistent skew towards positive mean values for all subpopulations (Fig 4b, S5 Fig and S6 Fig). In addition, we observed an excess of rare alleles relative to expectation (corresponding to negative Tajima's D values in the top 1% tail of the empirical distribution) for five phenology-related genes in historical Australian barley groups and seven phenology-related genes in geographic subpopulations, including known phenology genes *FLOWERING LOCUS T2* (*HvFT2*) and *AGAMOUS 1* (*HvAG1*) in North American and in CatD Australian subpopulations (Figs 4c and d). We also observed positive Tajima's D values, indicating a lack of rare alleles relative to expectation, which corresponds to a sudden population contraction, likely associated with the introduction of barley varieties to Australia and other countries (Figs 4c and d). Positive Tajima's D values were detected for phenology-related genes in historical Australian barley groups, including *FLOWERING LOCUS T1* (*HvFT1*) for the earliest- (CatA) and latest-released (CatC) Australian barley cultivars (Fig 4c), as well as for the North American and Asian varieties (Fig 4d).

To unravel genomic regions targeted by breeders in efforts to improve barley production in Australia in the last 120 years, we next explored loci in the barley genome that harbour selective sweeps related to breeding. To accomplish this, we examined population differentiation using the fixation index (F_{ST}) (S7 Fig), reduction of diversity (ROD), and cross-population composite likelihood ratio (Chen *et al.*, 2010) (XP-CLR) test scores, and compared these results within the Australian panel between the groups CatB and CatC groups (which have been subjected to recent breeding efforts) and those for the oldest group, CatA (Fig 5, S7 Fig, S8 Fig, S9 Fig).

We identified substantial population differences (high F_{ST} , Figs 5a and b) and genomic regions with substantially lower levels of diversity in more recently released cultivars than in historical cultivars (CatB and CatC groups, high ROD, Figs 5c and d) as possible candidate regions that were under selection during breeding in the recent past. To further assess the extent of genetic differentiation between early and recent Australian barley cultivars, we also used likelihood ratio

(XP-CLR) tests (Chen *et al.*, 2010) to identify genomic regions that had been differentially selected between the groups (Fig 5b). Regions above the 99th percentile of XP-CLR selection signals were considered candidates that had undergone selection during breeding, revealing 8 regions from the individual comparisons (CatA–CatB and CatA–CatC), of which 6 were adjacent to high- F_{ST} loci (Fig 5b). Based on the XP-CLR analysis, regions that had undergone selection contained 459 variants from 4 genes, with different regions between the two historical subpopulations.

In total, we identified 69 candidate regions with 922 potential genes that were potentially under selection during crop breeding, post-domestication and diversification in Australian barley (S2 File, S3 File). Among those genes, we identified 17 unique phenology-related genes, including gibberellin metabolism-related genes (the gibberellin oxidases *HvGA2ox8*, *HvGA20ox2*, *HvGA20ox4*, and *HvGA2ox4* and the gibberellin receptor genes *HvGID1L2* and *HvGID1L3*), *HvFT3* (also known as *PHOTOPERIOD 2*, *HvPPD-H2*), *EARLY FLOWERING 4-like4* (*HvELF4-like4*), and a homologue of *FLOWERING LOCUS T1* (*HvFT1-1*). More than 53% of the detected SNPs within these genes exhibited large differences in allele frequency among the different historical categories ($\geq 20\%$; S4 Table). To investigate possible functions of all candidate genes, we performed gene ontology and pathway enrichment analyses, revealing that genes under selection during barley breeding were related to responses to oxidoreductase activity, peroxidase activity, and antioxidant activity (S4 File).

We next predicted the variant effects of the 3,105 genetic variants located within the 69 candidate regions under selection in the 47 Australian barley cultivars within CatA, CatB, and CatC. Most genetic variants were detected in downstream (28%) or upstream gene regions (22%), while a large proportion of variants that fell within coding regions were missense variants (44%). Using a sorting intolerant from tolerant (SIFT) analysis, we identified 119 missense tolerated and deleterious mutations in 42 genes (S5 File). Of these variants, 29 were detected and annotated within 12 genes that exhibited large differences in allele frequency among the three historical categories ($\geq 20\%$; S5 Table). These 12 genes—which based on the categories we conclude have been under selection during the past 120 years—were observed only on chromosome 7H. Hence, these genes are attractive candidates for further investigation, as they may hold the potential to enhance agronomic traits in Australian barleys through breeding.

GWAS of agronomic traits

We field trialled and scored all 632 barley genotypes in the global barley panel for three key agronomic traits—flowering time (using days to Zadoks stage 49 [ZS49] as an equivalent for flowering time (Alqudah and Schnurbusch, 2017)), grain yield, and plant height—in sixteen independent field experiments at field sites located across the Western Australian wheatbelt region (Geraldton, Merredin, Katanning, Perth, and Esperance) conducted between 2015 and 2017. To evaluate the trait stability of the global barley panel across all locations and years, we calculated the coefficient of variation (CV) and heritability of each trait across all field trials (S6 File, S7 File). Z scores calculated per genotype for each field trial and trait revealed specific genotypes in the global barley panel with stable, consistent, and robust trait characteristics across the different field trials (S7 File). For example, the 2-row hulless and very early maturing Canadian variety CDC Speedy was one of the earliest-flowering varieties, irrespective of location or year, whereas Spanish landrace 355 was consistently late-flowering and tall-growing across all environments and years.

We then performed a multi-environment and multi-year GWAS to test if genetic variation identified from the global barley panel is associated with the key agronomic traits flowering time, grain yield, and plant height. We used 33,486 filtered genetic markers with a MAF > 0.01 for GWASs based on two statistical models, generalized linear models (GLMs) and mixed linear models (MLMs). Manhattan plots and quantile-quantile (QQ) plots of the three traits are provided in S10 Fig, S11 Fig, and S12 Fig. A graphical genotype map of selected significant genetic variants detected for all three traits is shown in Fig 6.

First, we consider the genetic variation associated with flowering time. Across all field trials, we identified 1,132 significant unique marker-trait associations (MTAs) (false discovery rate [FDR] of $P < 0.05$) for flowering time located within 327 unique genes with functional annotations, each explaining up to 18.7% of the phenotypic variation (S8 File, S10 Fig). These regions include known phenology-related genes, such as *HvPPD-H1*, *PHYTOCHROME C* (*HvPhyC*), *PROTEIN KINASE 2A* (*HvCK2A*), *HvADA2*, *PHYTOCHROME-ASSOCIATED PROTEIN 2* (*HvPAP2*), and *VERNALIZATION H1* (*HvVRN-H1*) (Hill *et al.*, 2016; Hill *et al.*, 2019a, b; He *et al.*, 2019). Furthermore, a total of 246 MTAs were considered to be ‘stable’, 76 MTAs were considered to be ‘consistent’, and 73 MTAs were considered to be ‘robust’ (S8 File). More than 30% of the significant MTAs were also detected in our previous study (Hill *et al.*, 2019a), in which a GWAS

was performed using 4,600 SNPs (in comparison to the 33,486 SNPs used in the present study) obtained from target enrichment sequencing data for phenology genes combined with field trial data from 2015 and 2016. Here, highly significant MTAs with no prior known associations were located within the genes HORVU5Hr1G096560 (disease resistance protein), HORVU5Hr1G095040 (beta glucosidase C), and HORVU5Hr1G104240 (zinc finger A20 and AN1 domain-containing stress-associated protein 6) on chromosome 5H (S8 File).

Notably, we also detected associations with candidate phenology-related genes that were not included in the previous target-enrichment sequencing study (Hill *et al.*, 2019a) but that have annotations linking them to roles in flowering time, including CCT and PRR motifs characteristic of key phenology genes such as *HvCO1*, *HvVRN-H2*, and *HvPPD-H1* (Hill *et al.*, 2016). These genes included HORVU1Hr1G011030 and HORVU5Hr1G125620 (both annotated as COP1-interacting protein-related), HORVU2Hr1G055130 and HORVU7Hr1G044380 (both annotated as CONSTANS, CO-like, and TOC1 [CCT] motif family protein, and HORVU3Hr1G092330 and HORVU6Hr1G008870 (both annotated as pentatricopeptide repeat-containing protein).

Next, we consider the genetic variation associated with grain yield. We identified 118 significant unique MTAs for grain yield, 30 of which were ‘robust’, with each MTA explaining up to 7.2% of the phenotypic variation (none were consistent or stable). We identified genetic variants within 30 functionally annotated genes (S9 File, S11 Fig). These regions included known phenology-related genes, such as *HvFT2* and *HvGA20ox2*. Like the results for flowering time, more than 30% of the significant MTAs for grain yield were also detected in our previous study (Hill *et al.*, 2019a).

Here, highly significant MTAs with no prior known associations were located within the genes HORVU2Hr1G125100 (peroxidase superfamily protein), HORVU7Hr1G002260 (disease resistance protein CC-NBS-LRR class family), and HORVU7Hr1G045290 (aluminum-activated malate transporter 9).

Finally, for plant height, we identified 1,279 significant unique MTAs within 395 functionally annotated genes, each explaining up to 8.9% of the phenotypic variation, including several gibberellin oxidase genes (*HvGA20ox4*, *HvGA20ox4*, and *HvGA20ox1*) and in particular the *sdw1/denso* gene (*HvGA20ox2*), a major determinant of plant height (Jia *et al.*, 2015) (S10 File, S12 Fig). Furthermore, a total of 190 MTAs were considered to be ‘stable’, one MTA was considered to be ‘consistent’, and 61 MTAs were considered to be ‘robust’ (S9 File). Of these

significant MTAs, approximately 8% were detected in our previous study (Hill *et al.*, 2019a). Highly significant MTAs with no prior known associations were located within the genes HORVU3Hr1G021140 (gigantea protein GI), HORVU3Hr1G022170 (homeobox-leucine zipper protein ROC4), and HORVU3Hr1G089160 (AP2-like ethylene-responsive transcription factor). Our data also revealed relevant candidate genes for functional annotation that are known to have pleiotropic effects on several agronomic traits, as exemplified by the major flowering time and plant height associations detected on chromosome 5H, where *HvPhyC* was a major driver (S8 File), an association previously detected and discussed in detail (Hill *et al.*, 2019a).

Combining the GWAS with the results from the selective sweep analysis, we then compared genes containing MTAs for all three traits (flowering time, grain yield, and plant height) with genes located within the 69 candidate regions under selection that are related to breeding in Australian barley (S3 File). The results show that 23, 7, and 23 genes with significant MTAs for all three traits, respectively, are located within breeding-related genomic regions (S8–10 Files) including the known phenology genes *HvPPD-H1*, *AGAMOUS-LIKE GENE 1 (HvAGL1)*, and *HvGA2ox3*. These results indicate that a subset of breeding loci are relevant for continued agronomic trait improvement and may have undergone additional selection to allow introduced European barleys to adapt to Australian growing conditions.

Discussion

The conservation of genetic diversity for the future breeding of crop varieties is particularly important in mitigating the adverse impacts of climate change on crop production. In this study, we assessed the contributions of both historical and recent breeding efforts towards local adaptation and crop improvement in a global barley panel of 632 genotypes. We used Australian barley as a model for the profound impact that breeding efforts had on developing a previously poorly adapted crop suitable to local environments. Most Australian and international cultivars in the global barley panel were varieties that led the market when they were released. Thus, this study panel represents the long-term breeding progress for the world's fourth-most, and Australia's second-most widely grown crop at peak agronomic performance.

The current study employed ~34,000 genetic markers—approximately 9 times more markers than used in previous studies of barley (Hill *et al.*, 2019a, b; He *et al.*, 2019)—to perform diversity,

selection footprint, and high-resolution GWAS analyses of agronomically relevant traits. By including sequencing data containing flowering-time related genomic regions, it was possible to detect low levels of variation in key candidate genes for flowering time that might have otherwise been missed. The diversity analyses revealed that the most recently released Australian cultivars exhibited more than 12% higher nucleotide diversity than earlier-released cultivars. Thus, our results show that modern Australian barley cultivars are not genetically depauperate in comparison to historical varieties, which counters the common perception that intensive breeding leads to the erosion of adaptive genetic diversity in modern cultivars (Van de Wouw *et al.*, 2010a, b). This notion is supported by several recent reports on cereal crops such as wheat, which demonstrate that genetic diversity has not been reduced in European wheat cultivars over the past five decades of progress in breeding (Göransson *et al.*, 2019; Voss-Fels *et al.*, 2019).

Moreover, the faster LD decay and lower level of long-range LD of more recently released cultivars indicate that recent breeding efforts have increasingly integrated lines with more diverse genetic backgrounds into the pool of Australian germplasm. This is different compared to a recent study of 429 chickpea genotypes (Varshney *et al.*, 2019) who found that LD decay in breeding lines was slower compared to landraces and elite cultivars, possibly due to a narrower genetic base of the domestic gene pool compared to barley. Taken together, these observations also highlight the importance of breaking large linkage blocks in future breeding efforts by increasing genetic diversity through new genetic crosses, such as unreleased breeding and research lines, landraces, and selected wild barley to eliminate the genetic hitchhiking of disadvantageous alleles within these LD blocks.

The selection footprint analyses for different Australian subpopulations detected substantial population differences (high F_{ST}) and genomic regions with substantially lower levels of diversity in more recently released cultivars than in historical cultivars (CatB and CatC groups, high ROD) as possible candidate regions that were under selection during breeding in the recent past (S3 Table). We estimate that 2.3% of barley genes (i.e. 922 genes) fall into the selected category and thus have been affected by breeding selection in Australian barley. An excess of rare alleles relative to expectation for five phenology-related genes in historical Australian barley groups and seven phenology-related genes in geographic subpopulations are consistent with an increase in population size following a bottleneck or a selective sweep, and could indicate strong selection during post-domestication (breeding) diversification. Notably, a lack of rare alleles for the

vernalisation response genes *VERNALISATION INSENSITIVE 3* (*HvVIN3*) and *HvZCCT-Hb* (the latter of which is a homologue of the *VERNALIZATION H2* (*VRN-H2*) locus) was detected for European and South American barley varieties, respectively. As the ancestor of domesticated barley is likely a winter-type wild barley (Saisho *et al.*, 2011), advantageous mutations in vernalisation response genes may have resulted in flowering time promotion in the absence of cold which facilitated the expansion of cultivable areas closer to the equator, pointing towards balancing selection for vernalisation requirements during past breeding efforts.

There are notable similarities between our study and a recent study by Pont *et al.* (2019), who explored how plant breeding has shaped the genome of modern bread wheats. They investigated selection footprints resulting from domestication breeding by comparing historical wheat groups, from old landraces to modern elite lines using exome sequencing of 500 bread wheat genotypes. They detected an initial ROD between old landraces and early cultivars (11.7% of ROD), indicating early breeding impacts, and a second stronger ROD between wheat varieties released prior to and after the green revolution (13.3% of ROD). They identified 5,089 polymorphic sites in the regions under selection, which also included phenology genes (e.g. photoperiod sensitivity, *Ppd*, and vernalization, *VRN*), as well as genes driving plant architecture (e.g. reduced height, *Rht*).

Next, we performed high-resolution GWAS analyses of agronomically relevant traits, and detected novel and highly significant MTAs for all three traits not detected in our previous study (Hill *et al.*, 2019a, b). For example, we detected highly significant MTAs for plant height, located within the phenology genes HORVU3Hr1G021140 (gigantea protein GI), and HORVU3Hr1G022170 (homeobox-leucine zipper protein ROC4). ROC4 influences the transcript levels of *GRAIN NUMBER*, *PLANT HEIGHT*, AND *HEADING DATE7* (*Ghd7*) and causes long day-dependent early flowering in rice (Wei *et al.*, 2016), whereas *GI* is a circadian clock gene responsible for fine-tuning plant developmental processes in response to photoperiod (Mishra and Panigrahi, 2015). For grain yield, highly significant MTAs were located within the genes HORVU2Hr1G125100 (peroxidase superfamily protein), HORVU7Hr1G002260 (disease resistance protein CC-NBS-LRR class family), and HORVU7Hr1G045290 (aluminum-activated malate transporter 9), which all play functional roles in pathogen and abiotic stress resistance (Almagro *et al.*, 2008; Zhang *et al.*, 2018). Interestingly, a gene encoding a CC-NBS-LRR-class

disease resistance protein was recently reported to be associated with grain yield in chickpea (Varshney *et al.*, 2019).

Finally, the diversity, selection footprint, and GWAS analyses performed in the present study demonstrate that several key loci, including major phenology genes such as *HvPPD-H1*, harbour coincident signals, supporting the view that a subset of breeding loci is relevant to the continued improvement of agronomic traits and have undergone additional selection in the adaptation of introduced European barleys to Australian growing conditions.

It will be interesting to follow-up these results with detailed genetic analyses on individual genes to characterize their functions in more detail. Gene manipulation in major crops to verify identified candidate genes is still a major bottleneck and requires the development of genomics-based tools, including CRISPR/Cas9 genome editing. Ideally, functional verification of the identified target genes should be performed in genetic backgrounds of commercial barley varieties as they are of agronomic value for breeding of future cultivars. The most widely used CRISPR/Cas9 genome editing method was optimized for the barley model cultivar Golden Promise; unfortunately, attempts to transform other barley varieties have been largely unsuccessful. A highly efficient and genotype-independent barley gene editing method based on anther culture was recently published, opening avenues for functionally verification of candidate genes in commercial barley varieties as well as precision barley breeding (Han *et al.*, 2020).

In summary, our combined variant dataset and germplasm collection provide a rich source of genetic information that can be applied to understanding and improving diverse traits, such as environmental adaptation and enhanced yield, and could accelerate genetic gains in future barley breeding.

Methods

Plant material

The barley panel consisted of 632 genotypes, including 250 cultivars and 382 breeding and research accessions from 37 countries throughout Europe, Asia, North and South America, Africa,

and Australia, and were selected from over 4,000 accessions preserved at the Western Barley Genetics Alliance at Murdoch University (Perth, Australia) to represent barley genotypes from major global barley breeding programmes. For a detailed description of all lines and varieties used in this study, see S1 File. This panel spanned the entire spectrum of cultivated barley, consisting of two- (92%) and six-row (8%) genotypes, and of winter (7%), spring (92%), and facultative (1%) growth habits. The selected germplasm also included 47 Australian cultivars released since 1903 as well as 180 Australian breeding and research lines. All cultivars of the global barley panel are highly productive and genetically uniform commercial varieties developed by professional plant breeders, whereas the all breeding and research lines include germplasm collections for potential use in developing future cultivars.

Field experiments and phenotypic data

A total of 16 field experiments were conducted in 2015, 2016, and 2017 across a variety of environments in Western Australia (South Perth, Geraldton, Katanning, two sites at Merredin and Esperance, respectively). The number of cultivars of the global barley panel tested at each field site are provided in S6 File. In Western Australia, Geraldton, South Perth, and Esperance are located along the coast of the Southern Ocean and all receive high annual rainfall but have very different daily maximum temperatures (the Geraldton site being the warmest, and the Esperance site being the coolest). The distance between Geraldton and Esperance is over 1,100 km. The Merredin site is located inland and receives little rainfall, while the Katanning site receives a medium amount of rainfall. The experimental design for field trial sites was performed as previously described (Hill *et al.*, 2019a). Briefly, all regional field trials (partially replicated design) were planted in a randomized complete block design using plots of 1 by 3 m² laid out in a row-column format. Field trials in South Perth were conducted using a hill plot technique with a 40-cm distance within and between rows due to space limitations. Seven control varieties were used for spatial adjustment of the experimental data.

Measurements were taken at each plot of each field experiment in the study to determine flowering time (days to ZS49), plant height, and grain yield as previously described (Hill *et al.*, 2019a). Briefly, plant maturity was recorded as the number of days from sowing to 50% awn emergence above the flag leaf (ZS49) (Zadoks *et al.*, 1974), as a proxy for flowering time (Alqudah and Schnurbusch, 2017). Plant height was determined by estimating the average height

from the base to the tip of the head of all plants in each plot. Grain yield (kg ha⁻¹) was determined by destructively harvesting all plant material from each plot to separate the grain and determining grain mass. Grain yield data collected in the 2015, 2016, and 2017 field experiments, as well as plant height and plant maturity data for the 2017 field trials, were analysed using linear mixed models (LMMs) in ASReml-R (<https://www.vsni.co.uk/software/asreml-r/>) to determine best linear unbiased predictions (BLUPs) or best linear unbiased estimations (BLUEs) for each trait for further analysis. Local best practices for fertilization and disease control were adopted for each trial site.

Evaluation of agronomic trait stability across field sites and genotypes

To evaluate the yield stability of global barley panel across all location and year scenarios in the main trials, we calculated the CV for flowering time, grain yield, and plant height across all location-by-year combinations, along with Z scores according to Equation 1:

$$Z = \frac{x_{ij} - \mu_j}{\sigma} \quad (1)$$

where x_{ij} is the trait value of variety i for year-by-location combination j , μ_j is the mean of the trait value for all plants of variety i in j , and σ is the standard deviation of the population mean.

Z scores were used to determine above- (positive Z score) and below-average (negative Z score)-yielding cultivars, as well as cultivars that flowered earlier (negative Z score) or later (positive Z score) or were shorter (negative Z score) or taller (positive Z score) than average for all year-by-location combinations. The critical Z score values for a 95% confidence level were -1.96 and $+1.96$ standard deviations, equal to a P-value of 0.05. Genotype trait characteristics (e.g. early flowering, high yielding, or short stature) were defined as ‘robust’ if they were consistently below or above the population mean in one location, ‘stable’ if they were significant (less than -1.96 or greater than $+1.96$ standard deviation) in more than one location, and ‘consistent’ if they were significant (less than -1.96 or greater than $+1.96$ standard deviation) across at least two years at one or more locations.

DNA extraction

Genomic DNA was extracted from the leaves of a single barley plant per variety using the cetyltrimethyl-ammonium bromide (CTAB) method as previously described (Hill *et al.*, 2019a, b).

DNA quality was assessed on 1% agarose gels and quantified using a NanoDrop spectrophotometer (Thermo Scientific NanoDrop Products, Wilmington, Delaware USA).

Sequencing, sequence alignment, genotype calling, variant discovery, and variant prediction

Target-enrichment sequencing

To assess the genetic diversity of phenology and phenology-related genes in the global collection of barley landraces and cultivars, we designed a custom target-enrichment sequencing assay for loci implicated in the flowering pathway in barley and related plant species, as previously described (Hill *et al.*, 2019a, b). In short, the target-enrichment sequencing of genomic DNA regions was performed by solution-based hybrid capture using a synthetic library consisting of 13,588 RNA probes (MYbaits, MYcroarray®, Ann Arbor, MI, USA) following the manufacturer's protocol (v.2.3.1). Post-capture DNA libraries were combined into 10 pools of approximately 96 samples each and sequenced on three lanes on an Illumina HiSeq 3000 (Illumina Inc., San Diego, CA, USA) to generate approximately 0.5 million 2x150-bp paired-end reads per sample. Genome sequencing was conducted at AgriBio (Centre for AgriBioscience, Bundoora, VIC, Australia). Sequence files were post-run filtered and aligned to the latest release of the barley reference genome assembly (Mascher *et al.*, 2017) (IBSC v2) using Nuclear software v.3.6.16 (GYDLE Inc., Montreal, Canada). SNP variant discovery and genotype calling were performed using custom Perl scripts to produce a variant call format (VCF) v.4.2 genotype file based on the alignment files as previously described (Hill *et al.*, 2019a). Only SNPs with <10% missing values and a MAF >1% (4,260 SNPs) were used for subsequent analyses.

Low-coverage whole genome sequencing

Each sample for low-coverage (1x) WGS consisted of a pool of 20 individual barley pre-capture DNA libraries from the target-enrichment sequencing experiment, dissolved in 10 mM Tris HCl (pH of 8.0). Thirty microliters of the pooled remainders of pre-capture DNA libraries were subjected to low-coverage WGS by the Beijing Genome Institute (BGI, Hong Kong) on an Illumina HiSeq 4000 to generate approximately 50 million 2x150-bp paired-end reads per sample.

The latest release of the barley reference genome assembly (IBSC v2) was used as a reference to map the clean reads with the alignment algorithm BWA-MEM (Li and Durbin, 2010) using default parameters. Duplicates were marked and removed using Picard v.1.129 (<http://broadinstitute.github.io/picard/>). Only reads with unique mapping positions in the reference genome were retained and used to detect genomic variations (SNPs and InDels).

InDels and SNPs were detected by running three rounds of SAMtools v.1.7 plus BCFtools v.1.7 and the Genome Analysis ToolKit (McKenna *et al.*, 2010) (GATK v.3.8) variant-calling pipeline. Briefly, the first round was performed with SAMtools plus BCFtools, with filtering based on both mapping quality and variant calling quality. The result of the first round was used as a guide for realignment around potential InDels and variant calling for the second round of GATK. The common variants detected by both the SAMtools plus BCFtools pipeline and the GATK pipeline were used to guide variant calling in the third round using GATK.

Genotyping-by-sequencing by DArTseq

In addition, DArTseq GBS was performed using the DArTseq platform (DArT PL, Canberra, NSW, Australia) according to the manufacturer's protocol (<https://www.diversityarrays.com/>). Briefly, 100 µl of 50 ng µL⁻¹ DNA was sent to DArT PL, and GBS was performed using complexity reduction followed by sequencing on a HiSeq Illumina platform as previously described (Akbari *et al.*, 2006). DArTseq marker sequences were aligned against the Morex barley genome assembly (Mascher *et al.*, 2017) IBSC v.2. The genetic position of each marker was determined based on the Morex physical reference assembly. Filtered DArTseq GBS (<10% missing values and minor allele frequency [MAF] >1%) yielded 14,032 SNPs across all 632 samples.

Stringent filtering steps were adopted to obtain clean data as previously described (Hill *et al.*, 2019a, b). All genotype data were combined, filtered based on duplicates and MAF >1%, and imputed using BEAGLE v.4.1 (Browning and Browning, 2013) to yield a final number of 33,486 filtered genetic markers (32,645 SNPs and 841 InDels) with a MAF > 1%.

Population structure and genotype data analyses

The PIC was calculated for each of the 33,486 filtered genetic markers according to Equation 2:

$$PIC = 1 - \sum_{i=1}^l P_i^2 - \sum_{i=1}^{l-1} \sum_{j=i+1}^l 2P_i^2 P_j^2 \quad (2)$$

where P_i and P_j are the population frequencies of the i^{th} and j^{th} alleles, respectively.

The ADMIXTURE v.1.3.0 model-based clustering algorithm (Alexander and Lange, 2011) was used to investigate the subpopulation structure of the global barley panel. Prior to subpopulation structure analysis in ADMIXTURE, the genotype dataset was LD pruned using Plink v1.9 (Purcell *et al.*, 2007) with a window size of 50 kb, step size of 5, and pairwise r^2 threshold of 0.5, yielding 18,869 genetic variants. A preliminary analysis was performed using 100 replicate runs by inputting successive values of K from 1 to 18, as previously described (Hill *et al.*, 2019a). A 10-fold cross-validation procedure was performed with 100 different fixed initial seeds in multi-threaded mode for each K-value. The most likely K-value was determined using ADMIXTURE cross-validation error values. CLUMPP (Jacobsson and Rosenberg, 2007) v.1.1.2 software was

used to obtain the optimal alignments of 100 replicates for each K-value. Membership proportions of each genotyped individual were averaged across runs according to the permutation with the greatest symmetric similarity coefficient as described previously (Hill *et al.*, 2019a). The output from CLUMPP for the optimal K-value was used to make plots using the cluster visualization package Pophelper v.2.2.3 (Francis, 2017) implemented in R v.3.5.1 (<http://www.R-project.org/>).

To summarize the genetic structure and variation present in the barley germplasm, PCA was also conducted using all 33,486 filtered genetic markers in TASSEL (Bradbury *et al.*, 2007) v.5.2.39. The first three PCs were plotted against each other using the ‘scatter plot’ function in Microsoft Excel 2016. NJ trees were constructed using the Java application Archaeopteryx v.0.9909 (Han *et al.*, 2009) based on genetic distances calculated in TASSEL v.5.2.39. The sub-structures in the collection inferred using different methodologies were compared, and the final K-value was ascertained using ADMIXTURE (Alexander and Lange, 2011).

Linkage disequilibrium

Genome-wide LD analysis was performed for the global barley panel and subgroups using all 33,486 filtered genetic markers using Plink v.1.9 (Purcell *et al.*, 2007). LD was estimated by using squared allele frequency correlations (r^2) between the intra-chromosomal pairs of loci (Weir, 1996). The loci were considered to be in significant LD when $P < 0.001$. To investigate the extent of and average LD decay in the panel, significant inter- and intra-chromosomal r^2 values within each 100-kb bin were plotted against the physical distance (kb) between markers. Curves were fitted by a second-degree LOESS function using R v.3.5.1 (<http://www.R-project.org/>).

Diversity parameter estimation and detection of selective sweeps

To detect genomic areas with selective sweeps driven by artificial (breeding) selection, we calculated F_{ST} , π , and ROD using VCFtools v.0.1.14 (Danecek *et al.*, 2011) and a window size of 10 Mb. F_{ST} estimates for pairs of subpopulations were calculated as previously described (Weir and Cockerham, 1984). Subpopulation-specific estimates of Tajima’s D were calculated using VCFtools v.0.1.14 (Danecek *et al.*, 2011) to compare the average number of pairwise differences and the number of segregating sites between samples within each of our geographic and historical subpopulations in Australia. The ROD index was calculated for each 10-Mb window based on the ratio of diversity between Australian subpopulation CatB to that of CatA according to Equation 3:

$$ROD=1 - (\pi \text{ CatA}/\pi \text{ CatB}) \quad (3)$$

and between Australian subpopulation CatC to that of CatA according to Equation 4:

$$ROD=1 - (\pi \text{ CatA}/\pi \text{ CatC}) \quad (4)$$

where the nucleotide diversity statistic π is the average number of nucleotide differences between any two DNA sequences. In addition, whole-genome screening of selected regions was performed using XP-CLR, a likelihood method for detecting selective sweeps that is based on the multilocus allele frequency differentiation between two populations (Chen *et al.*, 2010). XP-CLR tests were run with a window size and step size of 1 Mb, with CatA set as the reference and compared to CatB and CatC for each chromosome. Invariant or singleton SNPs were excluded, leaving on average ~45% of available variants for the analysis.

A total of 69 regions, which were in the highest 95th (F_{ST} , ROD) or 99th (XP-CLR) percentile of all regions identified, were considered to be under selection. Within the identified 69 regions under selection, 3,105 genetic variants were located within 922 genes. The 3,105 genetic variants were used for VEP using the Ensembl Variant Effect Predictor toolset (Ensembl Variant Effect Predictor web interface, <http://www.ensembl.org/vep>). VEP was performed to determine the effect of the genetic variants on genes, transcripts, and protein sequence, as well as regulatory regions. SIFT was estimated to predict the effects of amino acid substitutions on protein function based on sequence homology and the physical properties of amino acids. The results were filtered for missense only with SIFT scores (SIFT score <0.05) from tolerant to deleterious. Regions of genetic differentiation between subpopulations and genes within these regions were identified based on F_{ST} , π , ROD, and XP-CLR values of markers plotted linearly along each chromosome according to physical position. All visualizations were performed using the R packages yarr and ggplot2.

Gene ontology and pathway enrichment analysis

Gene ontology and pathway enrichment analysis of the 922 candidate genes under selection was performed as previously described (Hill *et al.*, 2016; Ho *et al.*, 2020). Briefly, singular enrichment

analysis (SEA) was performed using AgriGO v.2.0 (Tian *et al.*, 2017) with the following parameter settings: Fisher's test, 0.05 significance level, 5 minimum mapping entries, and complete gene ontology type.

Association analysis

GWASs were performed using TASSEL v.5.2.39 (Bradbury *et al.*, 2007) and a total of 33,486 filtered genetic variants with <10% missing values and a MAF >1%. Different statistical models were used to calculate P-values for putative MTAs as follows, which included population structure to avoid spurious associations. For the 2015 and 2016 data, a compressed MLM with a population structure (Q) matrix (PCs) and kinship (K) matrix (matrix of genetic similarities based on simple matching coefficients) was used to correct for population structure as previously described (Hill *et al.*, 2019a). According to the QQ plot, the MLM that incorporated Q and K was suitable for these datasets. Data from the 2015 and 2016 field trials were used in a previously published GWAS using only target-enrichment sequencing data (4,260 SNPs) (Hill *et al.*, 2019a). For the 2017 data, GLMs with PCs as a correction for population structure were tested for all associations, which, according to the QQ plots, were suitable for this study. For all MTAs, multiple testing using Storey's q-value method (Storey and Tibshirani, 2003) was performed to limit false discoveries and to assess statistical significance. As part of the q-value method, the smoother method, an extension of FDR correction, was employed. Lambda was set to 0, which estimates $\pi(0) = 1$, produces a list of significant tests equivalent to that obtained with the Benjamini and Hochberg (1995) procedure, and is considered to be a conservative case of Storey's q-value methodology. Only markers with a qFDR of <0.05 were considered to be significant. Manhattan plots were drawn with qman (Turner, 2014) v.0.1.4.

Broad-sense heritability (H^2) was calculated using the following equation by treating genotype and environment as random effects, applying an MLM according to Equation 5:

$$H^2 = \frac{\sigma_a^2}{\sigma_a^2 + \sigma_e^2} \quad (5)$$

where σ_a^2 and σ_e^2 represent the variance derived from genotypic and environmental effects, respectively.

MTAs were defined as 'robust' if they explained more than 5% of the phenotypic variation,

‘stable’ if they were identified in more than one location, and ‘consistent’ if they were identified in more than one year. Phenogram (Wolfe et al., 2013) was used to produce the graphical genotype map in Fig 6.

Data Availability Statement

All relevant data can be found within the manuscript and its supporting materials. The targeted resequencing data was previously published in the data descriptor (Hill et al., 2019b) and is publicly available in ArrayExpress under the identifier E-MTAB-7362. The SNP and InDel genotyping results for the phenology genes are available in Figshare: <https://doi.org/10.6084/m9.figshare.c.4357706> (Hill et al., 2019b).

Acknowledgements

We thank Ms Lee-Anne McFawn and Mr David Farleigh from DPIRD (South Perth, WA) for providing technical assistance in the field trials. This work was supported by funding from the Grains Research and Development Corporation (GRDC) of Australia (DAW00240/UMU00050, DAV00127 and DAW00233), Department of Primary Industries and Regional Development (WA, Australia), and Western Australian State Agricultural Biotechnology Centre (SABC).

Author contributions

C. H. and C. L. conceived the project. T. A. and C. L. collected the barley accessions, conducted initial phenological evaluation, and selected the accessions for this study. T. A., X. Z., and C. L. conducted the field experiments and phenotyping. K. C. conducted the spatial analysis of the agronomic data. C. H. and X. Z. collected samples and extracted DNA. C. H., P. W., G. Z., and C. T. prepared the low-coverage WGS and DArTseq data. C. H. prepared the target-enrichment sequencing data and generated the combined genetic marker data. C. H. performed all population genetic, GWAS, and statistical analyses. C. H. analysed all data and wrote the paper. C. L. supervised the project.

Conflict of interest statement

The authors declare no competing interests.

Short legends for Supporting Information

S1 Fig. Graphical genotype map of genetic variants detected using different genotyping technologies. Genetic markers detected via a) low-coverage whole genome sequencing, b) Genotyping-by-sequencing (GBS) by Diversity Arrays Technology sequencing (DArT-Seq), c) target capture sequencing, and d) all genetic markers combined (a- c). Plots drawn using the PhenoGram online software tool. Cat C: recent cultivars, released between 2006 and 2019; and Cat D: breeding and research lines).

S2 Fig. The extent of Linkage Disequilibrium (LD) across the seven chromosomes in a worldwide collection of domesticated barley varieties. Values are intra-chromosomal LD r^2 values for all intra-chromosomal pairs of genetic variants binned by distance. Curves were fitted by second-degree LOESS curve.

S3 Fig. Chromosomal distribution of nucleotide diversity (π) between different historic groups of domesticated Australian barley genotypes. Statistics based on 10 Mb windows. A legend is provided at the top of the figure. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005; CatC: Cultivars released between 2006 and 2019.

S4 Fig. Chromosomal distribution of nucleotide diversity (π) between different geographic groups of the barley diversity panel. Statistics based on 10 Mb windows. A legend is provided at the top of the figure.

S5 Fig. Chromosomal distribution of Tajima's D values between different historic groups of domesticated Australian barley genotypes. Statistics based on 10 Mb windows. Filled circles show values above the 99th percentile and are colour-coded according to the different historic groups. A legend is provided at the top of the figure. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005; CatC: Cultivars released between 2006 and 2019. CatD: unreleased breeding and research lines.

legend is provided at the top of the figure.

S6 Fig. Chromosomal distribution of Tajima's D values between different geographic groups of the barley diversity panel. Statistics based on 10 Mb windows. Filled circles show values

above the 99th percentile and are colour-coded according to the different geographic groups. A legend is provided at the top of the figure.

S7 Fig. Chromosomal distribution of Fixation Index (FST) values and XP-CLR scores between different historic groups of domesticated Australian barley genotypes. Highlighted regions (triangles coloured as per legend) are based on the 99th percentile (XP-CLR). Statistics based on 10 Mb windows. A legend is provided at the top of the figure. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005; CatC: Cultivars released between 2006 and 2019.

S8 Fig. Chromosomal distribution of Reduction of Diversity (ROD) values between CatA and CatB historic groups of domesticated Australian barley genotypes. Statistics based on 10 Mb windows. A legend is provided at the top of the figure. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005.
2005; CatC: Cultivars released between 2006 and 2019.

S9 Fig. Chromosomal distribution of Reduction of Diversity (ROD) values between CatA and CatC historic groups of domesticated Australian barley genotypes. Statistics based on 10 Mb windows. A legend is provided at the top of the figure. CatA: Cultivars released between 1903 and 1998; CatC: Cultivars released between 2006 and 2019.

S10 Fig. Manhattan and QQ plots of flowering time for all field trials with significant marker-trait associations. Days to ZS49 were used as an equivalent to flowering time (FT). Left panel: Manhattan plots, right panel: Quantile–quantile (QQ) plots. GWAS results are presented by negative log₁₀ of unadjusted p-values against position on each of the seven chromosomes. Horizontal dashed lines indicate the genome-wide significant threshold selected by local false discovery rate and a q-value cut-off at 0.05 (blue) and 0.01 (red).

S11 Fig. Manhattan and QQ plots of grain yield for all field trials with significant marker-trait associations. Other details as per legend to S18 Fig.

S12 Fig. Manhattan and QQ plots of plant height for all field trials with significant marker-trait associations. Other details as per legend to S18 Fig.

S1 Table. Barley genetic variants. Sequencing of the 632 genotypes delivered 33,486 filtered genetic markers detailed in the current table with the number of variants (SNPs and InDels) from low-coverage whole genome sequencing (LC), target capture sequencing (TC), and DArTseq (DArT), the number of genic and non-genic variants and associated ratio (genic/non-genic) with the number of targeted genes per chromosomes. ‘Genic variants’ columns count include those associated with a high confidence annotation. ‘Genes’ column includes genes with high-confidence annotation. ‘Non-genic’ variants include both ‘Upstream’ and ‘Downstream’ variants as well as variants without associated high-confidence annotation.

S2 Table. Nucleotide diversity index (π) and Tajima’s D summary statistics of the genetic variation observed within different subpopulations in the barley diversity panel. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005; CatC: Cultivars released between 2006 and 2019.

S3 Table. F_{ST} summary statistics in the genetic variation observed between different subpopulations in the barley diversity panel. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005; CatC: Cultivars released between 2006 and 2019.

S4 Table. Allele frequency of missense genetic variants in phenology-gene related candidate genomic regions that underwent selection during breeding of Australian barley cultivars. Shown are only variants within phenology-gene related gene regions (including 500bp flanking regions) and with >20% allele frequency (freq.) difference between CatA, CatB, and CatC barley cultivars. TF: transcription factor. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005; CatC: Cultivars released between 2006 and 2019.

S5 Table. Allele frequency of missense genetic variants in candidate genomic regions that underwent selection during breeding of in Australian barley cultivars. Shown are only variants with >20% allele frequency (freq.) difference of all missense variants between CatA, CatB, and CatC barley cultivars. All candidate genomic regions are located on chromosome 7H. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005; CatC: Cultivars released between 2006 and 2019.

S1 File: List of barley genotypes. The barley diversity panel consists of 632 genotypes sourced worldwide. Type: Res.: Research and breeding lines; Dom: Domestic cultivar. Breeder: DPI: Department of Primary Industries; Uni Adel: University of Adelaide; DAWA: Department of Agriculture, Western Australia; CIMMYT: International Maize and Wheat Improvement Center; TIA: Tasmanian Institute of Agriculture.

S2 File. Selected candidate regions from comparisons of Australian barley cultivars released between 1903 and 2019. There are 69 selected regions from comparing CatA with CatB and CatC, respectively, and 924 candidate genes falling within the regions. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005; CatC: Cultivars released between 2006 and 2019. Region ID: The identifier of a selected region; Region Coordinate: The range of the selected region; F_{ST} (CatA-CatB): Fixation Index (F_{ST}) values among CatA and CatB barley historic groups in the candidate region. Highlighted regions are above the 95th percentile. F_{ST} (CatA-CatC): Fixation Index (F_{ST}) values among CatA and CatC barley historic groups in the candidate region. Highlighted regions are above the 95th percentile. ROD (CatA-CatB): Reduction Of Diversity (ROD) values among CatA and CatB barley historic groups in the candidate region. Highlighted regions are above the 95th percentile. ROD (CatA-CatC): Reduction Of Diversity (ROD) values among CatA and CatC barley historic groups in the candidate region. Highlighted regions are above the 95th percentile. XP-CLR (CatA-CatB): Cross-Population Composite Likelihood Ratio Test (XP-CLR) values among CatA and CatB barley historic groups in the candidate region. Highlighted regions are above the 99th percentile. XP-CLR (CatA-CatC): Cross-Population Composite Likelihood Ratio Test (XP-CLR) values among CatA and CatC barley historic groups in the candidate region. Highlighted regions are above the 99th percentile.

S3 File. Candidate genes within selected candidate regions from comparisons of Australian barley cultivars released between 1903 and 2019. There are 69 selected regions from comparing CatA with CatB and CatC, respectively, and 924 candidate genes falling within the regions. Of these, 890 have gene-stable ID's and additional information is provided in this dataset. CatA: Cultivars released between 1903 and 1998; CatB: Cultivars released between 1999 and 2005; CatC: Cultivars released between 2006 and 2019. Gene ID: The candidate gene locus from barley annotation of IBSC v2 (Mascher *et al.*, 2017); Gene Type: A gene classification; Gene Coordinate: The range of the gene from annotation; Functional Annotation: Annotation from IBSC v2, Gene

name (TC): Abbreviated name given for genes selected and sequenced in the targeted re-sequencing (TC) phenology gene project (Hill et al., 2019a, b).

S4 File. Gene Ontology and pathway enrichment analysis for candidate genes under selection in Australian barleys. Singular Enrichment Analysis (SEA) was performed using AgriGO v.2.0 using Fisher test, 0.05 significance level, 5 minimum mapping entries and Complete GO gene ontology type.

S5 File. Variant Effect Predictor analysis for candidate genes under selection. Variant Effect Prediction (VEP) was performed using ensembl for candidate genes potentially under selection. Shown are results for missense only with Sorting Intolerant From Tolerated (SIFT) scores.

S6 File. Descriptive statistics for all investigated traits in the field trials. Minimal (Min) and Maximal (Max) values are shown next to the arithmetic mean (Mean), Standard deviation (SD) and coefficient of variation (CV). H^2 : Broad-sense heritability. N: Number of lines scored successfully in the field trial. *Plant development measured on 2 Oct as an estimation to ZS49. Esperance (1): Esperance field trial location 1. Esperance (2): Esperance field trial location 2 (EDRS). Merredin (1): non-irrigated. Merredin (2): irrigated. Perth (1): Perth, time of sowing 1 (early). Perth (2): Perth, time of sowing 2 (mid). Perth (3): Perth, time of sowing 3 (late).

S7 File. Z scores for all barley varieties and investigated traits in the field trials. The critical Z score values for a 95% confidence level were -1.96 and +1.96 standard deviations, equal to a P-value of 0.05. Genotype trait characteristics were defined as 'robust' if they were consistently below or above the population mean across one location (same value direction), 'stable' if they were significant (less than -1.96 or more than +1.96 standard deviations) for more than one location, and 'consistent' if they were significant (less than -1.96 or more than +1.96 standard deviations) across at least two years at one location or more. *Plant development measured on 2 Oct as an estimation to ZS49. Esperance (1): Esperance field trial location 1. Esperance (2): Esperance field trial location 2 (EDRS). Merredin (1): non-irrigated. Merredin (2): irrigated. Perth (1): Perth, time of sowing 1 (early). Perth (2): Perth, time of sowing 2 (mid). Perth (3): Perth, time of sowing 3 (late).

S8 File. Significant marker-trait associations for flowering time (FT, Days to ZS49) identified via genome-wide association mapping for all studied agronomic traits. MAF:

Minor allele frequency. R2: Contribution to phenotypic variation. P-value: Adjusted p-value after multiple comparisons false discovery rate testing using Storey's qvalue (Storey, 2002). Esperance (1): Esperance field trial location 1. Esperance (2): Esperance field trial location 2 (EDRS).

Merredin (1): non-irrigated. Merredin (2): irrigated. Perth (1): Perth, time of sowing 1 (early).

Perth (2): Perth, time of sowing 2 (mid). Perth (3): Perth, time of sowing 3 (late). Stable: Marker-trait association (MTA) detected at more than one location. Consistent: MTA detected for more than one year at the same location. Robust: More than 5% phenotypic variation explained.

S9 File. Significant marker-trait associations for grain yield identified via genome-wide association mapping for all studied agronomic traits. MAF: Minor allele frequency. R2:

Contribution to phenotypic variation. P-value: Adjusted p-value after multiple comparisons false discovery rate testing using Storey's qvalue (Storey, 2002). Esperance (1): Esperance field trial location 1. Merredin (1): non-irrigated. Stable: Marker-trait association (MTA) detected at more than one location. Consistent: MTA detected for more than one year at the same location. Robust: More than 5% phenotypic variation explained.

S10 File. Significant marker-trait associations for plant height identified via genome-wide association mapping for all studied agronomic traits. MAF: Minor allele frequency. R2:

Contribution to phenotypic variation. P-value: Adjusted p-value after multiple comparisons false discovery rate testing using Storey's qvalue (Storey, 2002). Esperance (1): Esperance field trial location 1. Merredin (1): non-irrigated. Stable: Marker-trait association (MTA) detected at more than one location. Consistent: MTA detected for more than one year at the same location. Robust: More than 5% phenotypic variation explained

References

- Alexander, D.H. and Lange, K. (2011) Enhancements to the ADMIXTURE algorithm for individual ancestry estimation. *BMC Bioinform.* **12**, 246.
- Akbari, M., Wenzl, P., Caig, V., Carling, J., Xia, L., Yang, S. *et al.* (2006) Diversity arrays technology (DArT) for high-throughput profiling of the hexaploid wheat genome. *Theor. Appl. Genet.* **113**, 1409–1420.

Almagro, L., Gómez Ros, L.V., Belchi-Navarro, S., Bru, R., Ros Barceló, A. and Pedreno, M.A. (2008) Class III peroxidases in plant defence reactions. *J. Exp. Bot.* **60**, 377–390.

Alqudah, A.M. and Schnurbusch, T. (2017) Heading date is not flowering time in spring barley. *Front. Plant Sci.* **8**, 896.

Benjamini, Y. and Hochberg, Y. (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. Series B Stat. Methodol.* **57**, 289–300.

Bradbury, P.J., Zhang, Z., Kroon, D.E., Casstevens, T.M., Ramdoss, Y. and Buckler, E.S. (2007) TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics*, **23**, 2633–2007.

Browning, B.L. and Browning, S.R. (2013) Improving the accuracy and efficiency of identity-by-descent detection in population data. *Genetics*, **194**, 459–471.

Bustos-Korts, D., Dawson, I.K., Russell, J., Tondelli, A., Guerra, D., Ferrandi, C. *et al.* (2019) Exome sequences and multi-environment field trials elucidate the genetic basis of adaptation in barley. *Plant J.* **99**, 1172–1191.

Chen, H., Patterson, N. and Reich, D. (2010) Population differentiation as a test for selective sweeps. *Genome Res.* **20**, 393–402.

Colen, L. and Swinnen, J. (2016) Economic growth, globalisation and beer consumption. *J. Agricult. Econ.* **67**, 186–207.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A. *et al.* (2011) The variant call format and VCFtools. *Bioinformatics*, **27**, 2156–2158.

Dempewolf, H., Baute, G., Anderson, J., Kilian, B., Smith, C. and Guarino, L. (2017) Past and future use of wild relatives in crop breeding. *Crop Sci.* **57**, 1070–1082.

Dreissig, S., Mascher, M. and Heckmann, S. (2019) Variation in recombination rate is shaped by domestication and environmental conditions in barley. *Mol. Biol. Evol.* **36**, 2029–2039.

Francis, R.M. (2017) pophelper: an R package and web app to analyse and visualize population structure. *Mol. Ecol. Res.* **17**, 27–32.

Friedt, W., Horsley, R.D., Harvey, B.L., Poulsen, D.M.E., Lance, R.C.M., Ceccarelli, S. *et al.* (2011) Barley Breeding History, Progress, Objectives, and Technology. In *Barley: Production, Improvement, and Uses* (Ullrich, S.E., ed.). New Jersey: Blackwell Publishing, pp. 160–220.

Gandhi, V.P. and Zhou, Z. (2014) Food demand and the food security challenge with rapid economic growth in the emerging economies of India and China. *Food Res. Int.* **63**, 108–124.

Göransson, M., Hallsson, J.H., Lillemo, M., Orabi, J., Backes, G., Jahoor, A. *et al.* (2019) Identification of ideal allele combinations for the adaptation of spring barley to Northern latitudes. *Front. Plant Sci.* **10**, 542.

Han, Y., Broughton, S., Liu, L., Zhang, X. Q., Zeng, J., He, X. and Li, C. (2020). Highly efficient and genotype-independent barley gene-editing based on anther culture. *Plant Commun.*, <https://doi.org/10.1016/j.xplc.2020.100082>

Han, M.V. and Zmasek, C.M. (2009) phyloXML: XML for evolutionary biology and comparative genomics. *BMC Bioinform.* **10**, 356.

He, T., Hill, C.B., Angessa, T.T., Zhang, X.Q., Chen, K., Moody, D., *et al.* (2019) Gene-set association and epistatic analyses reveal complex gene interaction networks affecting flowering time in a worldwide barley collection. *J. Exp. Bot.* **70**, 5603–5616.

Hill, C.B., Angessa, T.T., McFawn, L.A., Wong, D., Tibbits, J., Zhang, X.Q., *et al.* (2019a) Hybridisation-based target enrichment of phenology genes to dissect the genetic basis of yield and adaptation in barley. *Plant Biotechnol. J.* **17**, 932–944.

Hill, C.B., Cassin, A., Keeble-Gagnère, G., Doblin, M.S., Bacic, A. and Roessner, U. (2016) *De novo* transcriptome assembly and analysis of differentially expressed genes of two barley genotypes reveal root-zone-specific responses to salt exposure. *Sci. Reports*, **6**, 31558.

Hill, C.B. and Li, C. (2016) Genetic architecture of flowering phenology in cereals and opportunities for crop improvement. *Front Plant Sci.* **7**, 1906.

Hill, C.B., Wong, D., Tibbits, J., Forrest, K., Hayden, M., Zhang, X.Q. *et al.* (2019b) Targeted enrichment by solution-based hybrid capture to identify genetic sequence variants in barley. *Sci. Data*, **6**, 12.

Ho, W.W.H., Hill, C. B., Doblin, M. S., Shelden, M. C., van de Meene, A., Rupasinghe, T., *et al.* (2020). Integrative multi-omics analyses of barley rootzones under salinity stress reveal two distinctive salt tolerance mechanisms. *Plant Commun.* **1**, 100031.

Jakobsson, M. and Rosenberg, N.A. (2007) CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics*, **23**, 1801–1806.

Jia, Q., Li, C., Shang, Y., Zhu, J., Hua, W., Wang, J. *et al.* (2015) Molecular characterization and functional analysis of barley semi-dwarf mutant Riso no. 9265. *BMC Genom.* **16**, 927.

Komatsuda, T., Pourkheirandish, M., He, C., Azhaguvel, P., Kanamori, H., Perovic, D. *et al.* Six-rowed barley originated from a mutation in a homeodomain-leucine zipper I-class homeobox gene. *Proc. Natl. Acad. Sci. USA*, **104**, 1424–1429.

Li, H. and Durbin, R. (2010) Fast and accurate long-read alignment with Burrows–Wheeler transform. *Bioinformatics*, **26**, 589–595.

Liu, H.J. and Yan, J. (2019) Crop genome-wide association study: a harvest of biological relevance. *Plant J.* **97**, 8–18.

Malysheva-Otto, L.V., Ganai, M.W. and Röder, M.S. (2006) Analysis of molecular diversity, population structure and linkage disequilibrium in a worldwide survey of cultivated barley germplasm (*Hordeum vulgare* L.). *BMC Genet.* **7**, 6.

Mascher, M., Gundlach, H., Himmelbach, A., Beier, S., Twardziok, S.O., Wicker, T. *et al.* (2017) A chromosome conformation capture ordered sequence of the barley genome. *Nature*, **544**, 427–433.

Merchuk-Ovnat, L., Silberman, R., Laiba, E., Maurer, A., Pillen, K., Faigenboim, A. *et al.* (2018) Genome scan identifies flowering-independent effects of barley HsDry2. 2 locus on yield traits under water deficit. *J. Exp. Bot.* **69**, 1765–1779.

- Mishra, P. and Panigrahi, K.C. (2015) GIGANTEA—an emerging story. *Front. Plant Sci.* **6**, 8.
- McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A. *et al.* (2010) The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303.
- Nei, M. and Li, W.H. (1979) Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci. USA*, **76**, 5269–5273.
- Phan, H.T., Rybak, K., Bertazzoni, S., Furuki, E., Dinglasan, E., Hickey, L.T. *et al.* (2018) Novel sources of resistance to *Septoria nodorum* blotch in the Vavilov wheat collection identified by genome-wide association studies. *Theor. Appl. Genet.* **131**, 1223–1238.
- Pont, C., Leroy, T., Seidel, M., Tondelli, A., Duchemin, W., Armisen, D., Lang, D., Bustos-Korts, D., Goué, N., Balfourier, F. and Molnár-Láng, M. (2019) Tracing the ancestry of modern bread wheats. *Nat. Genet.* **51**, 905–911.
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D. *et al.* (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575.
- Saisho, D., Ishii, M., Hori, K. and Sato, K. (2011) Natural variation of barley vernalization requirements: implication of quantitative variation of winter growth habit as an adaptive trait in East Asia. *Plant Cell Physiol.* **52**, 775–784.
- Storey, J.D. and Tibshirani, R. (2003) Statistical significance for genomewide studies. *Proc. Natl. Acad. Sci. USA*, **100**, 9440–9445.
- Tian, T., Liu, Y., Yan, H., You, Q., Yi, X., Du, Z. *et al.* (2017) agriGO v2. 0: a GO analysis toolkit for the agricultural community, 2017 update. *Nucleic Acids Res.* **45**, W122–W129.
- Turner, S.D. (2014) qqman: an R package for visualizing GWAS results using Q-Q and manhattan plots. *J. Open Source Softw.* **3**, 731.
- Van de Wouw, M., Kik, C., van Hintum, T., van Treuren, R. and Visser, B. (2010a) Genetic erosion in crops: concept, research results and challenges. *Plant Genet. Resour.* **8**, 1–15.

Van de Wouw, M, van Hintum, T., Kik, C., van Treuren, R. and Visser, B. (2010b) Genetic diversity trends in twentieth century crop cultivars: a meta analysis. *Theor. Appl. Genet.* **120**, 1241–52.

Varshney, R.K., Thudi, M., Roorkiwal, M., He, W., Upadhyaya, H.D., Yang, W. *et al.* (2019) Resequencing of 429 chickpea accessions from 45 countries provides insights into genome diversity, domestication and agronomic traits. *Nat. Genet.* **51**, 857–864.

Voss-Fels, K.P., Stahl, A., Wittkop, B., Lichthardt, C., Nagler, S., Rose, T. *et al.* (2019) Breeding improves wheat productivity under contrasting agrochemical input levels. *Nat. Plants* **5**, 706–714.

Wei, J., Choi, H., Jin, P., Wu, Y., Yoon, J., Lee, Y.S. *et al.* (2016) GL2-type homeobox gene *Roc4* in rice promotes flowering time preferentially under long days by repressing *Ghd7*. *Plant Sci.* **252**, 133–143.

Weir, B.S. (1996) *Genetic Data Analysis II: Methods for Discrete Population Genetic Data*. 2nd ed. Sunderland: Sinauer Associates.

Weir, B.S. and Cockerham C.C. (1984) Estimating *F*-statistics for the analysis of population structure. *Evolution*, **38**, 1358–1370.

Wolfe, D., Dudek, S., Ritchie, M.D. and Pendergrass, S.A. (2013) Visualizing genomic information across chromosomes with PhenoGram. *BioData Min.* **6**, 18.

Zadoks, J.C., Chang, T.T. and Konzak, C.F. (1974) A decimal code for the growth stages of cereals. *Weed Res.* **14**, 415–421.

Zhang, L., Wu, X.X., Wang, J., Qi, C., Wang, X., Wang, G. *et al.* (2018) BoALMT1, an Al-induced malate transporter in cabbage, enhances aluminum tolerance in *Arabidopsis thaliana*. *Front. Plant Sci.* **8**, 2156.

Table legends

Table 1: Summary of molecular diversity and polymorphism information content for the whole panel and all subgroups. MAF: Minor allele frequency. PIC: Polymorphism information

content. N: Number of genotypes per group (where information available, see S1 File). CatA: Cultivars released between 1903 and 1998, CatB: cultivars released between 1999 and 2005; CatC: cultivars released between 2006 and 2019; Cat D: breeding and research lines.

Figure legends

Figure 1: Population structure of the global barley panel. a) Exploration of the optimal number of genetic subpopulations (K) using Δ cross-validation error and standard error values in the barley germplasm collection. A solid line denotes the choice of K=12 which represents the most likely number of subpopulations within the barley germplasm collection. b) Population structure of the entire barley panel was inferred by assuming twelve subpopulations (K). Each colour represents a different subpopulation as per the legend. Distribution of ADMIXTURE-defined populations based on c) seven geographical locations, d) three growth habits, and e) two row types. The neighbour-joining trees of 632 barley genotypes with clusters highlighted are based on f) geographic location or g) growth habit. h) Neighbour-joining trees of 47 selected Australian barley cultivars. Each colour represents a different historic group as per legend. The trees were constructed from simple matching distances of 33,486 common genetic variants in the barley population. Fac., facultative.

Figure 2: Principal component analysis (PCA) of the first two components of 632 barley varieties. a) PCA based on geographic region: The seven divergence groups are coloured respectively; b) PCA based on growth habit: The three divergence groups are coloured respectively; c) PCA based on row type: The two divergence groups are coloured respectively. PC1 and PC2 together explain about 54% of the total variation and partitioned the population into distinct clusters.

Figure 3: Genome-wide linkage disequilibrium (LD) decay in different historical groups of domesticated Australian barley genotypes. Values are reported as mean LD r^2 for all pairs of genetic variants binned by distance (100 kb). Curves were fitted by a LOESS function. CatA: Cultivars released between 1903 and 1998, CatB: cultivars released between 1999 and 2005, CatC: cultivars released between 2006 and 2019, Cat D: breeding and research lines, Total: total barley population of 632 varieties.

Figure 4: Genetic diversity and selection (breeding) signatures of different groups of domesticated barley genotypes. a) Plots of nucleotide diversity index (π) values and b) Tajima's D values to compare the average number of pairwise differences and the number of segregating sites between samples within each of our geographic and historical subpopulations in Australia (highlighted in light grey shading; a timescale is provided above the panel). Solid thin black horizontal lines indicate means, transparent horizontal bands of different colours indicate Bayesian 95% highest-density intervals (HDIs), black dots represent individual data points, full densities are shown as bean plots. c) Tajima's D distribution among the different historical groups of domesticated Australian barley genotypes and d) barley varieties from different geographic regions. Filled circles show values above the 99th percentile and are colour coded according to the different historical or geographic groups as indicated in the legends within the panels. Boxes point to data points above the 99th percentile that are located within phenology-related genes. Details are further described in the figure. All statistics are based on 10-Mb windows. CatA: Cultivars released between 1903 and 1998; CatB: cultivars released between 1999 and 2005; CatC: cultivars released between 2006 and 2019; Cat D: breeding and research lines.

Figure 5: Breeding selection signatures of domesticated Australian barley genotypes. a) Pirate plot of the genetic differentiation fixation index (F_{ST}) values between different historical groups of domesticated Australian barley genotypes (a timescale from 1903 to 2019 is provided above the panel). Solid thin black lines indicate means, black horizontal bands indicate Bayesian 95% highest-density intervals (HDIs), black dots represent individual data points, full densities are shown as bean plots. b) F_{ST} values and cross-population composite likelihood ratio (XP-CLR) test scores for CatA, CatB, and CatC historical barley groups on each chromosome (Chr.), illustrating the range of variation in diversity between these groups. c) Reduction of Diversity (ROD) distribution between the CatA and CatB historical barley, and d) distribution between the CatA and CatC historical barley groups. Highlighted regions (as per the legend for b and yellow bars for c and d) are above the 95th percentile (F_{ST} and ROD), or above the 99th percentile (XP-CLR). Boxes indicate regions located within phenology-related genes, with details further described in the figure. All statistics are based on 10-Mb windows. CatA: Cultivars released between 1903 and 1998, CatB: cultivars released between 1999 and 2005, CatC: cultivars released between 2006 and 2019.

Figure 6: Graphical genotype map of selected genetic variants associated with three agronomic traits. Significant genetic variants detected via genome-wide association studies for flowering time (FT, measured as days to Zadoks stage 49 [ZS49]), grain yield (GY), and plant height (PH). Only stable (S), consistent (C), and/or robust (R) markers are shown (Supplemental Files 8–10). Selected genetic variants (consistent and/or robust markers) that also fall within candidate regions for breeding selection are marked with orange asterisks. Plots drawn using the PhenoGram software tool. Marker categories are abbreviated as S: stable; SR: stable and robust; SCR: stable, robust and consistent; CR: consistent and robust; C: consistent; R: robust.

Tables

Table 1

Group	Average MAF	PIC	N
Whole panel	0.14	0.17	632
2-row group	0.14	0.17	579
6-row group	0.15	0.18	48
Spring	0.14	0.17	520
Winter/facultative	0.14	0.17	41
Australian (all)	0.13	0.16	227
Australian (CatA)	0.12	0.14	16
Australian (CatB)	0.13	0.15	14
Australian (CatC)	0.13	0.16	17
Australian (CatD)	0.13	0.16	180
European	0.13	0.16	141
North American	0.14	0.17	183
South American	0.13	0.16	34
Asian	0.15	0.18	24
African	0.13	0.15	15

tpj_15173_f1.png

a)

b)

c)

○ Flowering time △ Grain yield ◇ Plant height * Under breeding selection

■ S	■ SR	■ SCR
■ CR	■ C	■ R