

Fast response electromagnetic follow-ups from low latency GW triggers

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 J. Phys.: Conf. Ser. 716 012009

(<http://iopscience.iop.org/1742-6596/716/1/012009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 134.7.93.13

This content was downloaded on 13/09/2016 at 03:21

Please note that [terms and conditions apply](#).

You may also be interested in:

[Excimer laser annealing of perovskite thin films: morphology and gas-sensor properties](#)

X Du, J Tunney, M Post et al.

[Quantum regime of resonant-tunnelling diode oscillations](#)

I A Elant'ev, I U Bezotosny, M P Bezhko et al.

[GALAXY SURVEY ON THE FLY: PROSPECTS OF RAPID GALAXY CATALOGING TO AID THE](#)

[DETECTION OF GRAVITATIONAL WAVE OBSERVATIONS](#)

I. Bartos, A. P. S. Crotts and S. Márka

[Very low latency search for low mass CBC in the S6 and VSR2 data](#)

D Buskulic, the LIGO Scientific Collaboration and the Virgo Collaboration

[Electromagnetic follow-up of gravitational wave transient signal candidates](#)

Marica Branchesi and Ligo Scientific Collaboration and Virgo Collaboration

[An improved liquid nitrogen level sensor](#)

P W Nicholson

[Triangulation of gravitational wave sources with a network of detectors](#)

Stephen Fairhurst

Fast response electromagnetic follow-ups from low latency GW triggers

E J Howell¹, Q Chu¹, A Rowlinson², H Gao³, B Zhang⁴, S J Tingay^{5,6}, M Boër⁷ and L Wen¹

¹School of Physics, University of Western Australia, Crawley WA 6009, Australia

²CSIRO Astronomy and Space Science, Sydney, Australia

³Department of Astronomy, Beijing Normal University, Beijing 100875, China

⁴Department of Physics and Astronomy, University of Nevada Las Vegas, NV 89154, USA

⁵International Centre for Radio Astronomy Research, Curtin University, Perth, WA 6845, Australia

⁶ARC Centre of Excellence for All-sky Astrophysics (CAASTRO)

⁷CNRS - ARTEMIS, boulevard de l'Observatoire, CS 34229, 06304 Nice Cedex 04, France

E-mail: E-mail:eric.howell@uwa.edu.au

Abstract. We investigate joint low-latency gravitational wave (GW) detection and prompt electromagnetic (EM) follow-up observations of coalescing binary neutron stars (BNSs). Assuming that BNS mergers are associated with short duration gamma ray bursts (SGRBs), we evaluate if rapid EM follow-ups can capture the prompt emission, early engine activity or reveal any potential by-products such as magnetars or fast radio bursts. To examine the expected performance of extreme low-latency search pipelines, we simulate a population of coalescing BNSs and use these to estimate the detectability and localisation efficiency at different times before merger. Using observational SGRB flux data corrected to the range of the advanced GW interferometric detectors, we determine what EM observations could be achieved from low-frequency radio up to high energy γ -ray. We show that while challenging, breakthrough multi-messenger science is possible through low latency pipelines.

1. Introduction

The first observations by the advanced generation of interferometric gravitational wave (GW) detectors is set to ignite a new era of astronomy in which GW triggered follow-ups by facilities operating outside the GW spectrum will become common place. Complementary observations from electromagnetic (EM), neutrino or high energy particle facilities would not only improve the confidence of a GW detection, but also maximise the science return via a wealth of additional astrophysical information.

The advanced generation of detectors will be led by Advanced LIGO (aLIGO) [1] during the second half of 2015 and followed by Advanced Virgo (AdV) [2] a year later [3]. Coalescing systems of BNSs are one of the most promising sources of GWs in terms of detection [4]; at design sensitivity (around 2018), the expected detection rates of these sources are of order 0.4 to 400 BNS/year [5, 6]. One important aspect of these well-modeled inspiralling systems is that a detection can be made tens of seconds before the merger if enough waveform cycles can be detected to boost the signal-to-noise ratio [7, 8]. This could allow an *extreme low-latency* alert to be sent out to EM facilities as near real-time as possible to catch the fading EM signature. Such




Figure 1. A cartoon illustrating some of the possible EM pathways, timescales and end products of coalescing systems of NSs and BHs. Adapted from Chu et al. [9].

prompt follow-ups could provide valuable insight into the inner workings of these cataclysmic events.

Here we report on the results of a study conducted by Chu et al. [9] which examined the capability of a range of EM instruments for rapid follow-ups from extreme low-latency GW triggers. We present here only the main results assuming an aLIGO/AdV network at design sensitivity. We refer the interested reader to Chu et al. for a complete analysis including results from larger GW Network configurations.

2. The EM pathways for BNS mergers

Figure 1 summarises some of the popular scenarios for the expected EM signatures from BNS mergers. These are based on the proposed link between short duration gamma ray bursts (SGRBs) and the merger of compact objects [10, 11, 12, 13, 14]. The predictions are highly uncertain and could lead to various pathways and end products. The anticipated onset time, duration, and observable wavelengths are shown in the figure.

Figure 1 shows that SGRBs could be accompanied by an optical flash or a fast radio burst (FRB) [15]; an FRB could also result from the collapse of a merger product to a black hole [16]. If a stable or supra-massive magnetar is formed, the long lived X-ray plateaus observed in many SGRBs could be the signature of direct dissipation of magnetar wind energy [17, 18, 19]. At later times, predictions for an optical/IR kilonova [20] are supported though recent observations [21, 22]. For a relatively small opacity, a kilonova precursor may also appear soon after the merger [23].

SIGNAL TYPE	TARGET	TIME RANGE (s)	Gamma		X-ray	Optical/IR	Radio	
			High (> 10 MeV)	Low (< 10 MeV)			High (> 1 GHz)	Low (< 1 GHz)
Jet Related	SGRB Prompt	0-5	✓	✓	?	?		
	SGRB Afterglow	10s->	✓		✓	✓	✓	?
	Reverse Shock	60-10000	?	?	?	?	✓	?
Isotropic Ejecta Related	Kilonova	1000 ->			?	✓		
	Ejecta Afterglow	10000 ->			?	?	?	?
Other Signals	Magnetar Wind Dissipation	1-1000	?	✓	✓	?	?	?
	FRB (early)	-5 -10					?	?
	FRB (late)	10-10000					?	?

Figure 2. Some of the multimessenger observations that could be achieved for our follow-up scenario. The signals are grouped into jetted, isotropic or other (or unknown) signal types; the approximate emission time ranges around merger are also given. As shown by the legend, **ticks** indicate the EM observations with observational support; **question marks** indicate those that have been proposed through theoretical models; **dark cells** indicate there are EM instruments available in the band with sufficiently fast responses, exposure times and FoVs to cover or tile the GW error box within the emission time-range.

3. Accessing the low-latency GW detector response and EM follow-up

Since the initial LIGO era a number of low-latency GW trigger-generation pipelines have been proposed and tested [24, 25, 8, 26, 27]. The low latency localization pipeline BAYESTAR [28, 29] has shown that localization can be achieved in seconds with potential speed ups proportional to the number of CPU cores. For success, wide-field EM telescopes capable of rapid response within tens of seconds and with fields-of-view (FoV) comparable to the error area of GW source sky directions (of order hundreds of deg²) will be critical. To test the EM-follow up prospects we simulated a total of 200,000 BNS sources, drawing free parameters of the waveform and the sky location of each event from standard distributions. Around 10³ BNS detections were obtained and the localization accuracy of each was determined using the formalism of [30]. Detailed statistics of the detected sample are provided in Chu et al. [9]. For an aLIGO/AdV network the error region required to contain 50% of the detections, assuming sources are detected 40s before merger, was around 1000 deg². We assume this estimate, along with a 40s latency to send out the GW trigger.

To explore prompt EM follow-up observations, we estimate the fluxes at different wavelengths by extrapolating data from observed EM emissions associated with SGRBs and convert to 200 Mpc. We then consider the FoVs, sensitivities and response times for a range of EM telescopes to estimate their ability to capture a prompt EM signature.

4. Results and discussion

Figure 2 shows that at low frequency radio (MWA [31]; FoV:610 deg²@150 MHz; response in tens of secs) and both high and low-energy γ -ray (e.g. Fermi [32]; GBM FoV:30000 deg², LAT FoV:6564 deg²) the large FoVs and the fast response potential of MWA show that post-prompt (> 5s) breakthrough observations could be realised [33]. In the X-ray, small FoV instruments such as *Swift* are hindered by the large GW error region; a requirement for reconstructed sky locations to overlap with nearby galaxies [34] could improve this situation.

In the optical, even for a wide field, fast response telescope like ZTF ([35]; FoV:47 deg²; exposure 30s; 20.5-21 mag), to cover the 1000 deg² error region with an exposure times of 30s would require hundreds of seconds. Similar instruments will require sophisticated tiling strategies and would benefit from a co-ordinated observational scheme (see [36] for search strategies using the iPTF). A wide FoV X-ray instrument capable of rapid pointing, such as the NASA proposed 900 deg² FoV soft X-ray instrument ISS-Lobster [37], would be invaluable for such an approach.

ACKNOWLEDGMENTS

EJH acknowledges support from a UWA Research Fellowship and LW support from the Australian Research Council.

- [1] Aasi J *et al.* 2015 *Classical and Quantum Gravity* **32** 074001 (*Preprint* 1411.4547)
- [2] Acernese F *et al.* (VIRGO) 2015 *Class.Quant.Grav.* **32** 024001 (*Preprint* 1408.3978)
- [3] Aasi J *et al.* 2013 *astro-ph: 1304.0670* (*Preprint* 1304.0670)
- [4] Thorne K S 1987 *Three Hundred Years of Gravitation* ed Hawking, S W & Israel, W (Cambridge University Press) pp 330–458
- [5] Abadie J *et al.* 2010 *Classical and Quantum Gravity* **27** 173001 (*Preprint* 1003.2480)
- [6] Coward D M, Howell E J, Piran T, Stratta G, Branchesi M, Bromberg O, Gendre B, Burman R R and Guetta D 2012 *MNRAS* **425** 2668–2673 (*Preprint* 1202.2179)
- [7] Manzotti A and Dietz A 2012 *ArXiv e-prints* (*Preprint* 1202.4031)
- [8] Cannon K *et al.* 2012 *ApJ* **748** 136 (*Preprint* 1107.2665)
- [9] Chu Q, Howell E, Rowlinson A *et al.* 2015 *submitted to MNRAS*. (*Preprint* 1509.06876)
- [10] Paczynski B 1986 *ApJL* **308** L43–L46
- [11] Eichler D, Livio M, Piran T and Schramm D N 1989 *Nature* **340** 126–128
- [12] Gehrels N, Sarazin C L, O’Brien P T *et al.* 2005 *Nature* **437** 851–854 (*Preprint* arXiv:astro-ph/0505630)
- [13] Berger E *et al.* 2005 *Nature* **438** 988–990 (*Preprint* astro-ph/0508115)
- [14] Bloom J S *et al.* 2006 *ApJ* **638** 354–368 (*Preprint* astro-ph/0505480)
- [15] Totani T 2013 *PASJ* **65** L12 (*Preprint* 1307.4985)
- [16] Zhang B 2014 *ApJL* **780** L21 (*Preprint* 1310.4893)
- [17] Rowlinson A *et al.* 2010 *MNRAS* **409** 531–540 (*Preprint* 1007.2185)
- [18] Rowlinson A, O’Brien P T, Metzger B D, Tanvir N R and Levan A J 2013 *MNRAS* **430** 1061–1087 (*Preprint* 1301.0629)
- [19] Zhang B 2013 *ApJL* **763** L22 (*Preprint* 1212.0773)
- [20] Metzger B D, Martínez-Pinedo G, Darbha S, Quataert E, Arcones A, Kasen D, Thomas R, Nugent P, Panov I V and Zinner N T 2010 *MNRAS* **406** 2650–2662 (*Preprint* 1001.5029)
- [21] Berger E, Fong W and Chornock R 2013 *ApJL* **774** L23 (*Preprint* 1306.3960)
- [22] Tanvir N R, Levan A J, Fruchter A S, Hjorth J, Hounsell R A, Wiersema K and Tunnicliffe R L 2013 *Nature* **500** 547–549 (*Preprint* 1306.4971)
- [23] Metzger B D, Bauswein A, Goriely S and Kasen D 2015 *MNRAS* **446** 1115–1120 (*Preprint* 1409.0544)
- [24] Abadie J *et al.* 2012 *A&A* **541** A155 (*Preprint* 1112.6005)
- [25] Buskulic D *et al.* 2010 *Classical and Quantum Gravity* **27** 194013
- [26] Luan J, Hooper S, Wen L and Chen Y 2011 *ArXiv e-prints* 1108.3174 (*Preprint* 1108.3174)
- [27] Hooper S, Chung S K, Luan J, Blair D, Chen Y and Wen L 2012 *PRD* **86** 024012 (*Preprint* 1108.3186)
- [28] Singer L and Price L 2015 *in preparation*
- [29] Singer L P, Price L R, Farr B *et al.* 2014 *ApJ* **795** 105 (*Preprint* 1404.5623)
- [30] Wen L and Chen Y 2010 *Phys. Rev. D* **81** 082001
- [31] Tingay S J *et al.* 2013 *PASA* **30** e007 (*Preprint* 1206.6945)
- [32] Meegan C *et al.* 2009 *ApJ* **702** 791 URL <http://stacks.iop.org/0004-637X/702/i=1/a=791>
- [33] Howell E J, Rowlinson A, Coward D M *et al.* 2015 *PASA* **32** e046 (*Preprint* 1511.02959)
- [34] Evans P A *et al.* 2012 *ApJS* **203** 28 (*Preprint* 1205.1124)
- [35] Bellm E 2014 *The Third Hot-wiring the Transient Universe Workshop* ed Wozniak P R, Graham M J, Mahabal A A and Seaman R pp 27–33 (*Preprint* 1410.8185)
- [36] Singer L P, Kasliwal M M, Cenko S B *et al.* 2015 *ApJ* **806** 52
- [37] Camp J, Barthelmy S, Blackburn L, Carpenter K G, Gehrels N, Kanner J, Marshall F E, Racusin J L and Sakamoto T 2013 *Experimental Astronomy* (*Preprint* 1304.3705)