

Complete Genome Sequence of *Mesorhizobium ciceri* Strain CC1192, an Efficient Nitrogen-Fixing Microsymbiont of *Cicer arietinum*

Timothy Haskett,^a Penghao Wang,^a Joshua Ramsay,^b Graham O'Hara,^a Wayne Reeve,^a John Howieson,^a Jason Terpolilli^a

Centre for Rhizobium Studies, Murdoch University, Perth, Australia^a; School of Biomedical Sciences and Curtin Health Innovation Research Institute, Curtin University, Perth, Australia^b

We report the complete genome sequence of *Mesorhizobium ciceri* strain CC1192, an efficient nitrogen-fixing microsymbiont of *Cicer arietinum* (chickpea). The genome consists of 6.94 Mb distributed between a single chromosome (6.29 Mb) and a plasmid (0.65 Mb).

Received 22 April 2016 Accepted 28 April 2016 Published 9 June 2016

Citation Haskett T, Wang P, Ramsay J, O'Hara G, Reeve W, Howieson J, Terpolilli J. 2016. Complete genome sequence of *Mesorhizobium ciceri* strain CC1192, an efficient nitrogen-fixing microsymbiont of *Cicer arietinum*. *Genome Announc* 4(3):e00516-16. doi:10.1128/genomeA.00516-16.

Copyright © 2016 Haskett et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jason Terpolilli, J.Terpolilli@murdoch.edu.au.

Cicer arietinum (chickpea), a globally important grain legume, forms a N₂-fixing symbiosis with soil bacteria (rhizobia) in the genus *Mesorhizobium* (1) that can be harnessed in agriculture to enable *C. arietinum* to be grown without nitrogen fertilizer (2). When cultivated in soil lacking compatible rhizobia, *C. arietinum* can be inoculated with a symbiotically effective strain of *Mesorhizobium*. In Australia, *Mesorhizobium ciceri* strain CC1192 has been the commercial inoculant for this crop for several decades (3, 4). Although CC1192 has been the only *C. arietinum* inoculant used, there is evidence in the field that *C. arietinum* is nodulated by genetically distinct strains (5), suggesting that novel *C. arietinum*-nodulating rhizobia have evolved during the past 30 years. *Mesorhizobium* spp. can harbor symbiotic genes on mobile chromosomal regions called "symbiosis islands" or integrative and conjugative elements (ICEs) (6, 7). Thus, the complete genome sequence of CC1192 will enable the investigation of horizontal gene transfer of symbiosis genes from this strain to soil rhizobia.

Strain CC1192 was sourced from the Australian inoculant industry mother culture (<http://www.dpi.nsw.gov.au/content/agriculture/resources/soils/australian-inoculants-research-group>) and confirmed to fix N₂ effectively with *C. arietinum*, using established methods (8). CC1192 genomic DNA was extracted from a tryptone-yeast-grown culture (9) using a phenol-chloroform method as previously described (10). Whole-genome sequencing was performed by MacroGen (South Korea), using both Pacific BioSciences (PacBio) single-molecule real-time sequencing and Illumina HiSeq 2500 technology. Post filter, PacBio sequencing generated 96,824 trimmed reads with a mean read length of 11,579 bp, with Illumina sequencing generating an additional 19,850,308 paired-end reads. Raw Illumina reads were analyzed using FastQC version 0.10.1 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc>), and adaptors were removed by comparison against a comprehensive in-house adaptor sequence library. PacBio subreads were assessed using in-house software, and reads were automatically error-corrected in the assembly process.

The filtered Illumina and PacBio reads were assembled *de novo* using the hybrid approach of SPAdes assembler version 3.6.2 (11), with the number of mismatches and short indels reduced by incurring SPAdes's postprocessing module MismatchCorrector, utilizing the BWA tool (12). The assembly obtained was scaffolded using SSPACE version 3.0 (13) and annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (<http://www.ncbi.nlm.nih.gov/genomes/static/Pipeline.html>). The genome consists of 6,943,628 bp with an average GC content of 62.49%. There are 6,642 coding sequences distributed between a single circular chromosome of 6,295,397 bp and a single plasmid of 648,231 bp.

As is the case with *M. loti* R7A (14), genes potentially involved in nodulation (*nod*) and nitrogen fixation (*nif* and *fix*) appear to be located on a 419-kb symbiosis island integrated within the chromosome of CC1192 (base pairs 4,216,431 to 4,635,338). This region also harbors a type IV secretion system, conjugative relaxase, biotin and nicotinate biosynthetic clusters, proteins known to control excision and transfer of ICEMISym^{R7A} (15–17) and is flanked by direct repeat sequences. These characteristics are consistent with CC1192 carrying a mobile ICE. However, unlike ICEMISym^{R7A}, the CC1192 putative ICE is located adjacent to one of five serine tRNA genes and not a phenylalanine tRNA gene. Work is underway to investigate whether CC1192 can transfer its symbiotic capacity to non-*C. arietinum*-nodulating *Mesorhizobium* spp.

Nucleotide sequence accession numbers. The nucleotide sequence of the complete genome of CC1192 has been deposited in GenBank under the accession numbers CP015062 (chromosome) and CP015063 (plasmid pMc1192).

ACKNOWLEDGMENTS

This work was supported by grants UMU00040, UMU00044, and GRS10939 from the Grains Research and Development Corporation of Australia.

FUNDING INFORMATION

This work, including the efforts of Timothy L. Haskett, Graham W. O'Hara, John G. Howieson, and Jason J. Terpolilli, was funded by Grains Research and Development Corporation (GRDC) (UMU00040, UMU00044, and GRS10939).

REFERENCES

- Laranjo M, Alexandre A, Oliveira S. 2014. Legume growth-promoting rhizobia: an overview on the *Mesorhizobium* genus. *Microbiol Res* 169: 2–17. <http://dx.doi.org/10.1016/j.micres.2013.09.012>.
- Howieson JG, O'Hara GW, Carr SJ. 2000. Changing roles for legumes in Mediterranean agriculture: developments from an Australian perspective. *Field Crops Res* 65:107–122. [http://dx.doi.org/10.1016/S0378-4290\(99\)00081-7](http://dx.doi.org/10.1016/S0378-4290(99)00081-7).
- Drew EA, Herridge DF, Ballard RA, O'Hara GW, Deaker R, Denton MD, Yates RJ, Gemell G, Hartley E, Phillips L, Seymour N, Howieson JG, Ballard N. 2012. Inoculating legumes: a practical guide. Grains Research and Development Corporation, Kingston, Australia.
- Bullard GK, Roughley RJ, Pulsford DJ. 2005. The legume inoculant industry and inoculant quality control in Australia: 1953–2003. *Aust J Exp Agric* 45:127–140. <http://dx.doi.org/10.1071/EA03159>.
- Elias NV, Herridge DF. 2015. Naturalised populations of mesorhizobia in chickpea (*Cicer arietinum* L.) cropping soils: effects on nodule occupancy and productivity of commercial chickpea. *Plant Soil* 387:233–249. <http://dx.doi.org/10.1007/s11104-014-2298-z>.
- Sullivan JT, Ronson CW. 1998. Evolution of rhizobia by acquisition of a 500-kb symbiosis island that integrates into a phe-tRNA gene. *Proc Natl Acad Sci USA* 95:5145–5149. <http://dx.doi.org/10.1073/pnas.95.9.5145>.
- Nandasena KG, O'Hara GW, Tiwari RP, Sezmiş E, Howieson JG. 2007. *In situ* lateral transfer of symbiosis islands results in rapid evolution of diverse competitive strains of mesorhizobia suboptimal in symbiotic nitrogen fixation on the pasture legume *Biserrula pelecinus* L. *Environ Microbiol* 9:2496–2511. <http://dx.doi.org/10.1111/j.1462-2920.2007.01368.x>.
- Yates RJ, Howieson JG, Hungria M, Bala A, O'Hara GW, Terpolilli JJ. 2016. Authentication of rhizobia and assessment of the legume symbiosis in controlled plant growth systems, p 73–108. *In* Howieson JG and Dilworth MJ (ed.), *Working with rhizobia*. Australian Centre for International Agricultural Research, Canberra, Australia.
- Beringer JE. 1974. R factor transfer in *Rhizobium leguminosarum*. *J Gen Microbiol* 84:188–198. <http://dx.doi.org/10.1099/00221287-84-1-188>.
- Reeve WG, Tiwari RP, Melino V, Poole PS. 2016. Fundamental molecular techniques for rhizobia, p 221–244. *In* Howieson JG and Dilworth MJ (ed.), *Working with rhizobia*. Australian Centre for International Agricultural Research, Canberra, Australia.
- Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Prjibelski AD, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, Clingenpeel SR, Woyke T, McLean JS, Lasken R, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J Comput Biol* 20: 714–737. <http://dx.doi.org/10.1089/cmb.2013.0084>.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <http://dx.doi.org/10.1093/bioinformatics/btp324>.
- Boetzer M, Pirovano W. 2014. SSPACE-LongRead: scaffolding bacterial draft genomes using long read sequence information. *BMC Bioinformatics* 15:211. <http://dx.doi.org/10.1186/1471-2105-15-211>.
- Sullivan JT, Trzebiatowski JR, Cruickshank RW, Gouzy J, Brown SD, Elliot RM, Fleetwood DJ, McCallum NG, Rossbach U, Stuart GS, Weaver JE, Webby RJ, De Bruijn FJ, Ronson CW. 2002. Comparative sequence analysis of the symbiosis island of *Mesorhizobium loti* strain R7A. *J Bacteriol* 184:3086–3095. <http://dx.doi.org/10.1128/JB.184.11.3086-3095.2002>.
- Ramsay JP, Major AS, Komarovskiy VM, Sullivan JT, Dy RL, Hynes MF, Salmond GPC, Ronson CW. 2013. A widely conserved molecular switch controls quorum sensing and symbiosis island transfer in *Mesorhizobium loti* through expression of a novel antiactivator. *Mol Microbiol* 87:1–13. <http://dx.doi.org/10.1111/mmi.12079>.
- Ramsay JP, Sullivan JT, Jambari N, Ortori CA, Heeb S, Williams P, Barrett DA, Lamont IL, Ronson CW. 2009. A LuxRI-family regulatory system controls excision and transfer of the *Mesorhizobium loti* strain R7A symbiosis island by activating expression of two conserved hypothetical genes. *Mol Microbiol* 73:1141–1155. <http://dx.doi.org/10.1111/j.1365-2958.2009.06843.x>.
- Ramsay JP, Sullivan JT, Stuart GS, Lamont IL, Ronson CW. 2006. Excision and transfer of the *Mesorhizobium loti* R7A symbiosis island requires an integrase IntS, a novel recombination directionality factor RdfS, and a putative relaxase RlxS. *Mol Microbiol* 62:723–734. <http://dx.doi.org/10.1111/j.1365-2958.2006.05396.x>.