



September - November 2008  
Volume 26 Number 1

# AJAN

**australian journal of advanced nursing**

An international peer reviewed journal of nursing  
research and practice

**26:1**

## IN THIS ISSUE

### RESEARCH PAPERS

How general practice nurses view their expanding role

The changing role of practice nurses in Australia: an action research study

Nurses' descriptions of changes in cognitive function in the acute care setting

Nurses' views and practices regarding use of validated nutrition screening tools

Exploring women's reactions to a diagnosis of endometrial carcinoma: a pilot study of Australian women

Nurses' knowledge and practice involving patients' resuming sexual activity following myocardial infarction: Implications for training

Critical care nurses' knowledge about the care of deceased adult patients in an intensive care unit

The impact of therapeutic relationship on preoperative and postoperative patient anxiety

### SCHOLARLY PAPERS

Assessing leadership in nurse practitioner candidates

Legality, morality and reality - the role of the nurse in maintaining standards of care

Sampling methods: methodological issues involved in the recruitment of older people into a study of sexuality

Where have all the nurses gone?

### POINTS OF VIEW

A nurses' guide to the critical reading of research

## THE AUSTRALIAN JOURNAL OF ADVANCED NURSING

The Australian Journal of Advanced Nursing aims to provide a vehicle for nurses to publish original research and scholarly papers about all areas of nursing. Papers will develop, enhance, or critique nursing knowledge and provide practitioners, scholars and administrators with well-tested debate.

The AJAN will:

- publish original research on all nursing topics
- publish original scholarly articles on all nursing topics
- process manuscripts efficiently
- encourage evidence-based practice with the aim of increasing the quality of nursing care
- provide an environment to help authors to develop their research and writing skills
- provide an environment for nurses to participate in peer review

### Publisher & Editorial Office

Australian Nursing Federation  
PO Box 4239  
Kingston ACT, Australia 2604  
tel +61 2 6232 6533  
fax +61 2 6232 6610  
email: [ajan@anf.org.au](mailto:ajan@anf.org.au)  
<http://www.ajan.com.au>

ISSN 1447-4328

### Copyright

This journal is published in Australia and is fully copyrighted. All rights reserved. All material published in the Australian Journal of Advanced Nursing is the property of the Australian Nursing Federation and may not be reproduced, translated for reproduction or otherwise utilised without the permission of the publisher.

### Indexing

The AJAN is indexed in the CINAHL (Cumulative Index to Nursing and Allied Health Literature) Database, Current Contents, International Nursing Index, UnCover, University Microfilms, British Nursing Index, Medline, Australasian Medical Index and TOC Premier.

## PRODUCTION

### Editor

Jill Iliffe

### Journal Administrator

Rebecca Shaw

## EDITORIAL ADVISORY BOARD

### David Arthur, RN, PhD, FANZCMHN

Professor and Head, Alice Lee Centre for Nursing Studies, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

### Dr Joy Bickley Asher, RN, RM, Teaching Cert(Sec), BA, Ophthalmic N Dip(Hons), PG Dip(Nurs), PG Dip(Soc), PhD

Research Advisor, Royal New Zealand Plunket Society, Wellington, New Zealand

### Yu-Mei (Yu) Chao, RN, PhD

Adjunct Professor, Department of Nursing, College of Medicine, National Taiwan University, Taipei, Taiwan

### Mary Courtney, RN, BAdmin(Acc), MHP, PhD, FRCNA, AFCHSE

Assistant Dean (Research) Faculty of Health, Queensland University of Technology, Brisbane, Queensland, Australia

### Karen Francis, RN, PhD, MHlthSc, MEd, Grad Cert Uni Teach/Learn, BHlth Sc Nsg, Dip Hlth Sc Nsg

Professor and Head of School, School of Nursing and Midwifery, Monash University, Gippsland Campus, Churchill, Victoria, Australia

### Desley Hegney, RN, RM, CANN, COHN, DNE, BA(Hons), PhD, FRCNA, FAIM, FCN(NSW)

Professor, University of Queensland, Ipswich, Queensland, Australia

### Linda Kristjanson, RN, BN, MN, PhD

School of Nursing, Midwifery and Postgraduate Medicine, Edith Cowan University, Churchlands, Western Australia, Australia

### Anne McMurray, RN, BA(psych), Med, PhD, FRCNA

Peel Health Campus, Chair in Nursing, Murdoch University, Burleigh Heads, Queensland, Australia

### Colin Torrance, RN, BSc(Hon), PhD

Professor in Health Professional Education Department of Professional Education and Service Delivery Faculty of Health, Sport and Science University of Glamorgan Pontypridd, United Kingdom

### Lesley Wilkes, RN, CM RenalCert, BSc(Hons), GradDipEd(Nurs), MHPed, PhD

Professor of Nursing, Sydney West Area Health Service and the University of Western Sydney, Sydney, New South Wales, Australia

# Nurses' descriptions of changes in cognitive function in the acute care setting

## AUTHORS

### Malcolm Hare

RN, BSc (Nursing) (Honours),  
Nurse Manager, Informatics Support, Fremantle Hospital;  
Research Associate, School of Nursing and Midwifery,  
Curtin University of Technology, Western Australia  
malcolm.hare@health.wa.gov.au

### Sunita McGowan

RN, M.Sc (Nursing),  
Director Nursing Research and Evaluation, Fremantle  
Hospital; Adjunct Research Fellow, School of Nursing  
and Midwifery, Curtin University of Technology, Western  
Australia

### Dianne Wynaden

RN, RMHN, PhD,  
Associate Professor, School of Nursing and Midwifery,  
Curtin University of Technology; Research Consultant  
(Mental Health), Adjunct Senior Research Consultant,  
Department of Nursing Research, Evaluation and  
Informatics, Fremantle Hospital, Western Australia

### Gaye Speed

RN, B.AppSc (Nursing), PGradDip HAdmin,  
Nurse Consultant Clinical Improvement, Fremantle  
Hospital, Fremantle, Western Australia

### Ian Landsborough

RN, RMHN, M.Ed,  
Lecturer, School of Nursing and Midwifery, Curtin  
University of Technology, Western Australia

## KEY WORDS

confusion, delirium, quality nursing care, dementia,  
cognitive functioning, clinical audit

## ABSTRACT

### Objective

To describe nurses' documentation of cognition and behavioural changes in patients in acute care settings. Hospitalised patients often present with multiple co-morbidities including declining levels of cognitive functioning and this is particularly so for older people.

Many older people will experience a delirium during hospitalisation. While prevention or prompt management of delirium is paramount to providing quality care, research suggests that health professionals regularly fail to differentiate between delirium and other cognitive changes in hospitalised patients.

### Design

Four audits of progress notes were completed over a four week period at a Western Australian tertiary hospital to identify, quantify and categorise cognitive and behavioural changes in hospitalised patients. This paper describes data on nurses' documentation collected in the course of those audits. On four consecutive Thursdays, the medical records of all patients identified by nursing staff as being confused were reviewed. Where no definitive cause for the confusion was documented, the case notes were examined for evidence of risk factors to determine a probable cause.

### Setting

A Western Australian tertiary hospital.

### Subjects

The medical records of all patients identified by nursing staff as being confused.

### Main outcome measures

Documentation in patient medical records of a patient having cognitive or behavioural changes or being confused; use of a cognitive screening tool; or a diagnosis of delirium.

### Results

A total of 1209 patients were surveyed over the four audit days with 183 patients (15%) being identified as confused. 'Confusion' was the most common descriptor used by nurses to describe cognitive and behavioural changes; in many cases it was the only term used. Many of these changes were indicative of delirium. Little use by any health professional of cognitive screening tools was found.

### Conclusion

Cognitive and behavioural changes are a common problem in hospitalised patients who are elderly. The use of the term confusion to describe a range of cognitive and behavioural changes is a barrier to accurate identification of delirium, which is often the first indicator of serious underlying illness.

## INTRODUCTION

The international literature is critical of the widespread lack of recognition of delirium (Hustey et al 2003; Inouye et al 1999; Flacker and Marcantonio 1998). Nurses' inability to recognise delirium was illustrated by a recent study on nurses' level of knowledge of delirium and associated risk factors. The study reported that only 42% of nurses scored 70% or better on delirium knowledge questions and only 3.5% of nurses scored 70% or better for knowledge of delirium risk factors (Hare 2006). It has also been suggested that knowledge deficits may be perpetuated because health professionals consistently and routinely document cognitive and behavioural changes under the encompassing term 'confusion', which does not allow for qualification or quantification of significant changes in a patient's functioning (Milisen et al 2002). This practice continues even though prevention, or early intervention, for delirium has been shown to improve outcomes for this group of patients (Inouye et al 2005a).

Delirium presents as an acute short-term disturbance of consciousness lasting from a few hours to a few months characterised by disorganised thinking, perceptual disturbances and inattention (American Psychiatric Association 1999), and is often the first or only presenting sign of an underlying medical problem (Meagher 1998). Delirium causes cognitive and behavioural changes in a patient that fluctuate over the course of the day, can usually be attributed to a physiological cause and/or the use of substance(s), and is not better explained by a pre-existing or evolving dementia (American Psychiatric Association 1999). Three main variants of delirium exist: hyperactive, hypoactive and mixed. The hyperactive variant involves hyper-vigilance with associated agitation and hallucinations (American Psychiatric Association 1999). Patients experiencing the hyperactive variant are more likely to have their condition recognised and receive early intervention due to their presenting behaviours (Breitbart et al 2002; O'Keeffe 1999). Patients with the hypoactive variant often present as lethargic, drowsy and have difficulty focussing attention, and this form is more

likely to go unrecognised. The mixed variant fluctuates between features of both.

In older people, delirium is a predictor of mortality and morbidity, but delirium and its underlying causes are commonly under-diagnosed and as a consequence under-treated (Douglas et al 2005; Cole 2004; Inouye et al 1999; O'Keeffe 1999). This lack of recognition occurs in 25% of cases (Young and George 2003) increasing to 87.5% when the patient has a pre-existing dementia (Milisen et al 2002). In addition, delirium in elderly patients often results in a decreased level of functioning post discharge from hospital, which impacts on the individual's future quality of life (Inouye et al 1998). Estimates suggest that delirium is present in up to 62% of hospitalised orthopaedic patients who are elderly (Olofsson et al 2005) rising to as high as 89% in patients who have a pre-existing dementia (Fick et al 2002).

Nurses are well placed to identify, assess and document cognitive and behavioural changes as they spend more time with patients than most other health professionals. This is especially important for older patients, who are at high risk of developing delirium (McCusker et al 2003; O'Keeffe and Lavan 1996; Cole et al 1994), and whose care would be enhanced by prompt and accurate assessment and documentation of cognitive and behavioural changes which may be indicative of a serious underlying medical problem warranting urgent further investigation. However as previously noted, nurses' documentation of cognition is reportedly poor (Inouye et al 2005c). To determine whether the standard of documentation in a Western Australian teaching hospital was consistent with that reported in the literature, data on descriptors used by nurses to describe changes in cognition and behaviour were collected in the context of a point prevalence audit for delirium (Speed et al 2006).

## DESIGN

The Chair of the Human Research Ethics Committee was informed of the proposed audit and recommended that the audit be registered as a quality improvement activity. The audit was planned for one day in each of four consecutive weeks to provide a large sample size

and to facilitate inter-rater checking, as some patients were present during two or more audits. On each of the audit days, nurse coordinators on the medical and surgical wards at the hospital were asked to identify patients who displayed confusion, disorientation, fluctuating behaviours or who were behaving in an unusual manner. To identify patients who may have the hypoactive presentation of delirium, information was also sought about patients who appeared withdrawn, quiet or depressed. All inpatient records of the identified patients were then reviewed.

### DATA COLLECTION METHODS

Initially, a comprehensive review of the literature was conducted to identify the common causes, risk factors and presenting symptoms associated with delirium. Based on the literature review, the audit tool was piloted from the progress notes of several patients prior to the audit. The tool collected demographic information and data on behavioural descriptors identified during the literature review. Behavioural descriptors gathered from progress notes during the pilot were included as a tick-box list, with space for the auditors to add additional descriptors. Data were also gathered on whether the cognitive and behavioural changes fluctuated or were consistent, whether the patient had a pre-existing, confirmed diagnosis of dementia and whether there was evidence that the patient was confused on admission. All data were derived exclusively from the patients' progress notes. The five auditors were senior nurses, and inter-rater reliability was established by consensus in data collected during the pilot.

When a patient was identified with 'confusion', evidence was then sought from the patient record for any definitive medical diagnosis associated with the behaviour, such as pre-existing dementia, delirium or an organic disorder. Any data that assisted in confirming the cause of the behaviour, for example, 'delirium secondary to pneumonia', were also recorded. The audit team also looked for evidence of the use of formal cognitive assessment tools, such as the Mini Mental State Examination (MMSE) (Folstein et al 1975), Delirium Rating Scale (DRS)

(Trzepacz et al 2001) and Confusion Assessment Method (CAM) (Inouye et al 1990). Lastly, the audit team recorded behavioural descriptors associated with confusion if they were present in the nursing or integrated progress notes. Where information obtained suggested that the patient was likely to be suffering a delirium, the variety of descriptors used to document the patient's behaviour was identified (Speed et al 2006).

### RESULTS

Data from the audit showed that 183 (15%) of 1209 patients were identified by staff as being confused. Of the 183, 132 (72%) had documented cognitive and behavioural changes that were deemed to be associated with the presence of a delirium, while in the remaining 51 (28%) patients their behaviour was typical of their established dementia or organic disorder. Seventy-seven (58%) of the 132 patients deemed to have a delirium were documented as being confused on admission. Fourteen (11%) of the 132 had no documentation of their cognitive status on admission.

Of the 132 patients with cognitive and behavioural changes suggestive of delirium 78 (59%) were females and 54 (41%) were males. The age range was from 45 years to 95 years with a mean age of 79.9 (SD 10.21) years. A majority of these patients ( $n=78$  59%) had been admitted from home, 34 (26%) from a hostel, 14 (11%) from a nursing home and 6 (4%) transferred from other hospitals. Eighty-one (62%) patients were admitted as medical inpatients, 32 (24%) as orthopaedic inpatients, 11 (8%) as surgical inpatients and 8 (6%) were vascular inpatients.

Only 48 (36%) of the 132 patients with documented behavioural and cognitive changes consistent with delirium had a diagnosed delirium documented in their patient record. The records of the remaining 84 (64%) patients had descriptions of cognitive and behavioural changes suggestive of delirium but no formal diagnosis of delirium had been recorded. This may have occurred in part because the delirium was overlooked due to the presence of an established dementia ( $n = 58$ ).

A total of 364 descriptors (mean 2.76 per patient, SD 1.568) were found in the documentation audit. The number of descriptors in the notes ranged from one (n=33) to eight (n=1). Notably, one in eight cases (12.88%, n=17) had the term 'confused' as the only descriptor. The most common descriptors were 'confused' (n=99), 'disoriented' (n=54) and 'agitated' (n=46) (see table 1).

**Table 1: Descriptors of cognitive and behavioural changes**

Descriptor	Count	Percent
Confused	99	27.20
Disoriented	54	14.84
Agitated	46	12.64
Hallucinating	22	6.04
Drowsy	21	5.77
Plucking/pulling	17	4.67
Physically aggressive	15	4.12
Wandering	14	3.85
Vague	13	3.57
Verbally Abusive	10	2.75
Noisy	9	2.47
Forgetful	8	2.20
Insomnia	8	2.20
Lethargic	8	2.20
Non compliant	7	1.92
Fearful	5	1.37
Angry	4	1.10
Anxious	2	0.55
Threatening	2	0.55
Total	364	100.00

Only 43 (32.5%) of the 132 patients had evidence of the use of MMSE and only one (0.7%) had a DRS completed.

## DISCUSSION

The rate of delirium in hospitalised patients recorded during the audits (11%) is consistent with international rates (Milisen et al 2002; Dupplis and Wikblad 1999; Lynch et al 1998). Furthermore, the rates of undiagnosed delirium are also similar to that reported in the literature (Hustey et al 2003; Milisen et al 2002). Although delirium was not formally documented as a diagnosis in all of the 132 patients,

behavioural and cognitive changes consistent with delirium were recorded in the patients' records.

Many patients in this study had a pre-existing dementia. These patients are a vulnerable group, and are at high risk of developing delirium. Assessment and knowledge of baseline data on the patient's usual level of cognitive functioning is vital for early recognition and investigation of delirium, and crucial to their future quality of life (Rigney 2006). Health professionals should routinely assess elderly patients' cognitive and behavioural status, with the same priority afforded to the measurement of other vital signs, as recommended by Flaherty et al (2007). This would facilitate the early recognition of delirium, reducing adverse outcomes such as falls (O'Keeffe 1999), mortality, morbidity, length of stay and other long term or permanent sequelae (McCusker et al 2003; Inouye et al 1999).

This study provides additional confirmation of nurses' lack of knowledge of delirium and its associated risk factors and has provided a basis for ongoing research in this area. A further strength was the large audit sample size, which allowed for confirmation of inter-rater reliability, as some patients were audited more than once over the four weeks. A limitation of the study was the reliance on the nurse coordinator to identify confused patients. This is an important limitation, given that nurses are poor at identifying delirium (Inouye, Leo-Summers et al 2005) and some confused patients were likely to have been unreported.

## CONCLUSION

Nurses have a pivotal role in the accurate assessment of cognition and early detection of delirium in acute care settings and therefore the reduction of associated adverse events, such as falls (O'Keeffe 1999). Nevertheless, cognitive assessment has not been routinely included as a key component in nursing curricula and therefore has not translated into nursing practice.

To facilitate early recognition of any change in cognition and behaviour, it is recommended that nurses are educated and trained to conduct routine

cognitive assessment using validated tools in all hospitalised patients and to differentiate between acute and chronic cognitive conditions. Improved knowledge and assessment techniques will lead to improved documentation.

## REFERENCES

- American Psychiatric Association. *Practice guideline for the treatment of patients with delirium*. 1999. Available from [http://www.psych.org/psych\\_pract/treatg/pg/pg\\_delirium.cfm](http://www.psych.org/psych_pract/treatg/pg/pg_delirium.cfm) (accessed March 2006).
- Breitbart, W., Gibson, C. and Tremblay, A. 2002. The delirium experience: delirium recall and delirium related distress in hospitalised patients with cancer, their spouses/caregivers, and their nurses. *Psychosomatics*, 43(3):183-194.
- Cole, M. 2004. Delirium in elderly patients. *American Journal of Geriatric Psychiatry*, 12(1):7-21.
- Cole, M., Primeau, F., Bailey, R., Bonnycastle, M., Masciarelli, F., Engelsmann, F., Pepin, M. and Ducic, D. 1994. Systematic intervention for elderly inpatients with delirium: a randomised trial. *Canadian Medical Association Journal*, 151(7):965-970.
- Douglas, D., Zhang, Y., Holford, T., Bogardus, S., Leo-Summers, L. and Inouye, S. 2005. Premature death associated with delirium at 1-year follow-up. *Archives of Internal Medicine*, 165(14):1657-1661.
- Dupplis, G., and Wikblad, K. 1999. Acute confusional states in patients undergoing hip surgery. *Gerontology*, 46(1):36-43.
- Fick, D., Agostini, J. and Inouye, S. 2002. Delirium superimposed on dementia: a systematic review. *Journal of the American Geriatrics Society*, 50(10):1723-1732.
- Flacker, J., and Marcantonio, E. 1998. Delirium in the elderly: optimal management. *Drugs and Ageing*, 13(2):119-130.
- Flaherty, J., Rudolph, J., Shay, K., Kamholz, B., Boockvar, K., Shaughnessy, M., Shapiro, R., Stein, J., Weir, C. and Edes, T. 2007. Delirium is a serious and under-recognized problem: why assessment of mental status should be the sixth vital sign. *Journal of the American Medical Directors Association*, 8(5):273-275.
- Folstein, M., Folstein, S. and McHugh, P. 1975. 'Mini Mental State': a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12(3):189-98.
- Hare, M. 2006. A questionnaire to determine nursing staff at Fremantle Hospital level of knowledge of delirium and its risk factors. Honours Thesis, School of Nursing and Midwifery, Curtin University of Technology, Perth.
- Hustey, F., Meldon, S., Smith, M. and Lex, C. 2003. The effect of mental status screening on the care of elderly emergency department patients. *Annals of Emergency Medicine*, 41(5):678-684.
- Inouye, S., Bogardus, S., Charpentier, P., Leo-Summers, L., Acampora, D., Holford, T. and Cooney, L. 2005a. A multicomponent intervention to prevent delirium in hospitalised older patients. *The New England Journal of Medicine*, 340(9):669-676.
- Inouye, S., Leo-Summers, L., Zhang, Y., Bogardus, S., Leslie, D. and Agostini, J. 2005b. A chart-based method for identification of delirium: validation compared with interviewer ratings using the confusion assessment method. *Journal of the American Geriatrics Society*, 53(2):312-318.
- Inouye, S., Schlesinger, M. and Lydon, T. 1999. Delirium: a symptom of how hospital care is failing older persons and a window to improve quality of hospital care. *The American Journal of Medicine*, 106(5):565-573.
- Inouye, S., Rushing, J., Foreman, M., Palmer, R. and Pompei, P. 1998. Does delirium contribute to poor hospital outcomes? A three-site epidemiologic study. *Journal of General Internal Medicine*, 13(4):234-242.
- Inouye, S., van Dyck, C., Alessi, C., Balkin, S., Siegel, A. and Horwitz, R. 1990. Clarifying confusion: the confusion assessment method. *Annals of Internal Medicine*, 113(12):941-948.
- Lynch, E., Lazor, M., Gellis, J., Orav, J., Goldman, L. and Marcantonio, E. 1998. The impact of postoperative pain on the development of postoperative delirium. *Anesthesia and Analgesia*, 86(4):781-785.
- McCusker, J., Cole, M., Dendukuri, N. and Belzile, E. 2003. Does delirium increase hospital stay? *Journal of the American Geriatrics Society*, 51(11):1539-1546.
- Meagher, D. 1998. Delirium: the role of psychiatry. *Advances in Psychiatric Treatment*, 7(6):433-443.
- Milisen, K., Foreman, M., Wouters, B., Driesen, R., Godderis, J., Abraham, I. and Broos, P. 2002. Documentation of delirium in elderly patients with hip fracture. *Journal of Gerontological Nursing*, 28(11):23-29.
- O'Keeffe, S. 1999. Clinical subtypes of delirium in the elderly. *Dementia and Geriatric Cognitive Disorders*, 10(5):380-385.
- O'Keeffe, S. and Lavan, J. 1996. Predicting delirium in elderly patients: development and validation of a risk-stratification model. *Age and Ageing*, 25(4):317-321.
- Olofsson, B., Lundström, M., Borssén, B., Nyberg, L. and Gustafson, Y. 2005. Delirium is associated with poor rehabilitation outcome in elderly patients treated for femoral neck fractures. *Scandinavian Journal of Caring Sciences*, 19(2):119-127.
- Rigney, T. 2006. Delirium in the hospitalised elder and recommendations for practice. *Geriatric Nursing*, 27(3):151-157.
- Speed, G., Wynaden, D., McGowan, S., Hare, M. and Landsborough, I. 2006. Prevalence rate of delirium at two hospitals in Western Australia. *Australian Journal of Advanced Nursing*, 25(1):38-43.
- Trzepacz, P., Mittal, D., Torres, R., Kanary, K., Norton, J. and Jimerson, N. 2001. Validation of the delirium rating scale revised 98: comparison with the delirium rating scale and the cognitive test for delirium. *Journal of Neuropsychiatry and Clinical Neuroscience*, 13(2):229-242.
- Young, L. and George, J. 2003. Do guidelines improve the process and outcomes of care in delirium. *Age and Ageing*, 32(5):525-528.