

This is the peer reviewed version of the following article: Fougrouse, D. and Micklethwaite, S. and Halfpenny, A. and Reddy, S. and Cliff, J. and Martin, L. and Kilburn, M. et al. 2016. The golden ark: Arsenopyrite crystal plasticity and the retention of gold through high strain and metamorphism. *Terra Nova*. 28 (3): pp. 181-187, which has been published in final form at <http://doi.org/10.1111/ter.12206> This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving at <http://olabout.wiley.com/WileyCDA/Section/id-828039.html>

1 The golden ark: Arsenopyrite crystal plasticity and the
2 retention of gold through high strain and metamorphism.

3 **Denis Fougereuse^{1,2}, Steven Micklethwaite^{2,3}, Angela Halfpenny^{4,5}, Steven Reddy¹, John
4 B. Cliff⁶, Laure A.J. Martin⁶, Matt Kilburn⁶, Paul Guagliardo⁶, and Stanislav Ulrich⁷**

5 *¹Department of Applied Geology, Curtin University, GPO Box U1987, Perth, WA 6845,
6 Australia*

7 *²Centre for Exploration Targeting, School of Earth and Environment, The University of
8 Western Australia, M006, 35 Stirling Highway, Crawley, WA 6009, Australia*

9 *³School of Earth, Atmosphere & Environment, Monash University, Melbourne, Victoria,
10 Australia*

11 *⁴Microscopy & Microanalysis Facility, John de Later Centre, Curtin University, GPO BOX
12 U1987, Perth, WA 6845, Australia*

13 *⁵CSIRO Earth Science and Resource Engineering, Minerals Down Under Flagship, ARRC, 26
14 Dick Perry Avenue, Kensington, WA 6151, Australia*

15 *⁶Centre for Microscopy, Characterisation and Analysis, The University of Western Australia,
16 35 Stirling Highway, Crawley, WA 6009, Australia*

17 *⁷AngloGold Ashanti Ltd, Asset Development, 44 St George Terrace, Perth, WA 6000, Australia*

18

19 **ABSTRACT**

20 Quantitative electron backscatter diffraction analysis and ion microprobe imaging of gold-rich
21 arsenopyrites provide the first insights into the crystal plasticity and element mobility
22 behaviour of arsenopyrites through metamorphism (340° - 460° and 2 kbars). Remarkably, the
23 gold-rich arsenopyrites remained structurally and chemically robust during high strain
24 deformation. It is only during a superimposed lower strain deformation event, at a high angle

25 to the preferred orientation of the arsenopyrites, that small amounts of crystal plasticity affected
26 the arsenopyrites. During the low strain event a dissolution-reprecipitation reaction resulted in
27 loss of gold from the crystal lattice, facilitated by localised domains of recrystallization, most
28 likely due to fluid percolation along sub- and new grain boundaries. We suggest the abundance
29 and rheologically robust nature of gold-rich arsenopyrite in giant gold deposits, affected by
30 greenschist-amphibolite metamorphism, is actually a critical factor in the preservation of those
31 deposits.

32

33 **INTRODUCTION**

34 Although rarely acknowledged, arsenopyrite is one of the world's most important
35 sulphide minerals. It is ubiquitous as a gold bearing phase in many giant deposits, particularly
36 in sediment hosted orogenic gold deposits (Groves et al., 1998) as documented at Obuasi,
37 Ghana, (Oberthür et al., 1994), Sunrise Dam, Australia (Sung et al., 2009), Muruntau,
38 Uzbekistan (Morelli et al., 2007) or Red Lake, Canada (Dubé et al., 2004). The presence of
39 arsenic and arsenopyrite also provide important vectors to mineral resources (Boyle and
40 Jonasson, 1973; Eilu and Groves, 2001; Le Vaillant et al., 2015). For these reasons it is
41 surprising that almost nothing is known about the crystal plasticity of arsenopyrite, nor the
42 influence of microstructure on the ability of arsenopyrite to retain or release its major and trace
43 element content.

44 In contrast to arsenopyrite, the crystal plasticity of other sulphide phases such as pyrite,
45 pyrrhotite, sphalerite, chalcopyrite, galena, stibnite and pentlandite have been investigated in
46 detail (Barrie et al., 2011; Boyle et al., 1998; Cox, 1987; Kelly and Clark, 1975; Vukmanovic
47 et al., 2014). These studies conclude that crystal plasticity of phases, such as pyrite, occurs
48 relatively easily at temperatures as low as 260 °C and strain rates of approximately 10^{-12} - 10^{-

49 16 s^{-1} (Barrie et al., 2011) and can have a significant impact on trace element mobility (Reddy
50 and Hough, 2013).

51 Here we provide the first documentation of arsenopyrite crystal plastic behaviour under
52 high strain and greenschist facies metamorphic conditions. We examine the impact on trace
53 element modification, with important implications for both resource preservation and the
54 release of elements from arsenopyrite back to the natural environment.

55

56 **SAMPLES AND METHODS**

57 In order to examine the crystal-plastic behaviour of arsenopyrite and its impact on trace
58 element mobility, arsenopyrites with large enrichments of trace elements (Au, Ni) and well-
59 constrained metamorphic-strain conditions were selected from the economic refractory ores of
60 the giant Obuasi deposit. The mineralized arsenopyrites are hosted in paleoproterozoic sheared
61 phyllites (2155 Ma, zircon U/Pb maximum age of deposition; Oberthür et al., 1998) composed
62 of micas, quartz, carbonates (ankerite and siderite), chlorites, rutile and graphite. Arsenopyrite
63 growth and mineralization occurred during continuous cleavage development and peak upper
64 greenschist metamorphism at 340° - 460° and 2 kbars as shown by the development of strain
65 shadows parallel with the continuous cleavage surrounding the arsenopyrites (the Eburnean
66 orogeny: Schwartz et al. 1992; Oberthür et al. 1994; Fougrouse et al. in press). A large sample
67 suite (>200 samples) was collected and studied from mineralised exposures of the Obuasi mine
68 to provide representative petrographical and geochemical results across the deposit. Two grains
69 from sample 215-20 are reported in this study, collected ~950 m below the surface from a 10
70 m wide high strain domain associated with thick quartz veins, between two graphitic shear
71 zones. The arsenopyrites have been subject to two strain events (Fougrouse et al., in press-b),
72 mostly accommodating pure shear. The first event is high strain, at the peak of metamorphism,
73 which occurred during the Eburnean orogeny (D2_{Ob}; Fougrouse et al., in press). A pervasive

74 continuous cleavage developed while the quartz veins underwent boudinage and strain shadows
75 developed around the arsenopyrite grains (Fig. 1). The long-axes of the acicular arsenopyrites
76 (up to 0.1x1 mm) are aligned parallel to the cleavage. A large minimum finite strain of 48%
77 shortening and 235% stretch, perpendicular to the shortening, was estimated for the hosting
78 phyllites from strain shadow length and fold periodicity (Fig. 1; p. 271 Ramsay and Huber,
79 1983), using oriented thin sections cut orthogonal to the cleavage and parallel to the lineation.
80 Nonetheless, the arsenopyrite grains themselves remain largely euhedral and relatively
81 undeformed (Fig. 2). In the second event (D3_{Ob}), a low strain overprinting crenulation cleavage
82 developed, representing a minimum shortening estimated at 15% with no associated stretch.
83 The long-axes of the arsenopyrites are oriented at a high angle to the later crenulation cleavage.

84 Backscattered Scanning Electron Microscopy imaging (BSE) and Secondary Ion Mass
85 Spectrometer (SIMS and NanoSIMS) investigations were carried out at the Centre for
86 Microscopy, Characterisation and Analysis, University of Western Australia. BSE images were
87 collected using a TESCAN VEGA3 SEM and large area ion imaging of ¹⁹⁷Au were performed
88 on a Cameca IMS 1280 ion probe on 150x150 micron areas using a ~1 micron, Cs⁺ primary
89 beam. Ions were collected in peak-hopping mode in an ion-counting electron multiplier at a
90 mass resolution of ~3000. High-resolution elemental mapping was carried out using the
91 Cameca NanoSIMS 50 with a spot size of approximately 100 nm and a Cs⁺ primary beam. The
92 experiment allowed simultaneous detection of ¹⁹⁷Au and ³⁴S across areas measuring 100 x 100
93 μm with a 120 nm/pixel resolution. Quantitative orientation maps of arsenopyrite crystal lattice
94 were recorded using EBSD (Prior et al., 1999) at the CSIRO Earth Science and Resource
95 Engineering in Kensington (Australia) using a Zeiss Ultraplus FEG SEM and at Curtin
96 University using a TESCAN MIRA3 SEM. Systematic mis-indexing due to pseudosymmetry
97 within the arsenopyrite grains was removed by identifying the axis and angle relationships to
98 rotate the points and produce a correctly indexed grain. It should be noted that pseudosymmetry

99 is a problem in pyrite and arsenopyrite. Correcting for pseudosymmetry can produce internally
100 consistent datasets, although the actual solution may still be wrong (Nolze et al., 2016),
101 however in this study such issues do not change our conclusions.

102

103 **RESULTS**

104 Arsenopyrites from the ore zone of the Obuasi deposit are composed of two primary
105 domains: rims that appear homogenous in BSE images (A-rims) and inclusion-rich internal
106 cores. The cores contain inclusions of pyrrhotite, chalcopyrite or rutile. There is then a
107 discontinuously developed third domain (B-rims) crosscutting all others (Fig. 3). B-rims are
108 characterised by higher arsenic concentrations (~46 wt% as opposed to ~44 wt% in A-rims;
109 Fougere et al. in press) and preserve the original shape of the crystals. The B-rim contacts
110 are sharp, planar to curvilinear (bulbous) and penetrate in from microcracks and grain
111 boundaries (Fig. 3).

112 The EBSD data show a relative change in crystallographic orientation for each grain (Fig.
113 2 and 3). Consistent with other studies investigating sulphides crystal-plasticity (Reddy and
114 Hough 2013), we define high-angle grain boundaries to have misorientations $\geq 10^\circ$. All
115 boundaries $< 10^\circ$ are low-angle grain boundaries, representing subgrains. In most cases the
116 intragrain variation does not reach 5° , though new grains are formed at a few localised points.

117 The analysis reveals a difference in the deformation intensity between grain 1 and 2 (Figs.
118 2 and 3). Grain 1 is parallel to the main cleavage (S_{2Ob}), in a domain unaffected by the later
119 crenulation cleavage (S_{3Ob}), and contains only weak internal deformation (maximum of 3°
120 misorientation). Conversely, grain 2 is located on a fold flexure of the S_{3Ob} crenulation. In this
121 crystal, high densities of misorientations occur in a narrow localised domain, adjacent to a
122 fracture filled with pyrite (Fig. 3D). In this domain, the misorientations define low- and high-
123 angle boundaries creating new small sized grains. In the centre top of the grain an incipient

124 domain of less than 3° misorientation is developed. The cumulative orientation profiles (Fig.
125 3H) along the short axis of grain 2 show minor misorientations ($<2^\circ$), but much greater
126 misorientations are achieved parallel to the long axis of the arsenopyrite (up to 7° from the tip
127 of the grain to the core). Pole figure data indicates a dispersion of the orientation (Fig. 3). In
128 addition, the pyrite crystal infilling the fracture of grain 2 is virtually undeformed
129 (misorientation up to 2° ; Fig. 3C).

130 The SIMS and NanoSIMS ^{197}Au ion microprobe imaging reveals that the distribution of
131 the gold is zoned within the crystals, with gold-rich A-rims around gold-poor cores (Fig. 2 and
132 3). The A-rims are in turn crosscut by gold-poor B-rims (Fig. 3F). The A-rims represent
133 epitaxial zones with up to 1000 ppm Au (Fisher et al., 2014; Oberthür et al., 1994), although
134 these concentrations decrease on the outer edges of the grain as shown on the left side of grain
135 2 (Fig. 3). An A-rim is well developed in grain 1 where minimal internal deformation occurred.
136 In grain 2, a large A-rim is cut by discontinuous B-rims, characterized by low Au signal.
137 Internally, the A-rims are finely zoned with submicron-scale concentric bands with sharp
138 boundaries (Fig. 3F; Fougrouse et al. in press GCA). However, this fine epitaxial zoning in
139 the domain of high misorientation identified in grain 2, appears blurry, with diffuse boundaries
140 between bands and locally, small spherical high Au concentrations. The fuzzy gold signal in
141 this domain is also dissected by numerous small scale, sharply defined B-rims with low Au
142 signals. Critically, the overall B-rim width in this region of high densities of misorientations is
143 up to three times greater in comparison with the domains of undeformed arsenopyrite.

144

145 **CRYSTAL-PLASTIC EVOLUTION OF ARSENOPYRITES AND CONSEQUENCES** 146 **FOR GOLD RETENTION**

147 The crystallographic misorientations and localised intragrain textures, revealed by EBSD,
148 are consistent with dynamic recrystallisation by dislocation creep (Hirth and Tullis, 1992; Urai

149 et al., 1986). In grain 2, this led to the development of subgrains and neoblasts accommodating
150 crystal-plastic strain in a narrow localised domain, on the top right-hand side of the crystal.
151 This crystal-plastic strain domain is coincident with a fracture filled with undeformed pyrite
152 (Fig. 1). Overall, several significant features emerge from our analysis.

153 Firstly, grain 1 and grain 2 formed during high strain (D2_{Ob}, 50% flattening, >200%
154 stretch) and upper greenschist metamorphism (John et al., 1999; Schwartz et al., 1992), then
155 were subject to a second shortening event (D3_{Ob}, ~15% shortening), yet remained euhedral and
156 underwent only minor crystal-plastic deformation. Secondly, grain 1 has minimal lattice
157 distortion ($\leq 3^\circ$ of misorientation from grain tip to centre) in comparison to grain 2, which
158 contains a localized domain of subgrains, neoblasts and microfracture. In this regard, grain 1
159 is located parallel to the S2_{Ob} cleavage, in a microlithon between two crenulation planes (S3_{Ob}),
160 while grain 2 is located on a fold flexure of the S3_{Ob} cleavage. These relationships indicate that
161 crystal plastic microstructures developed not during the high strain peak metamorphic event
162 but during microfolding in the lower strain D3_{Ob} event, which only affected grain 2. Consistent
163 with this interpretation, the orientation profiles (Fig. 3H) show that crystal-plasticity and the
164 largest amount of lattice misorientation is recorded parallel to the D3 shortening direction. As
165 the fold grew, the arsenopyrite firstly underwent crystal-plastic deformation, then a fracture
166 propagated through the arsenopyrite in the hinge, perhaps facilitated by embrittlement related
167 to crystal-plastic strain hardening (Fleck et al., 1994). The fracture is infilled with undeformed
168 pyrite (misorientation $< 2^\circ$), again indicating the arsenopyrite crystal-plasticity occurred early
169 during D3_{Ob} microfolding.

170 We conclude arsenopyrite is remarkably resilient to crystal plasticity at the recorded
171 temperatures of 340° - 460° and pressures of 2 kbars. Two factors likely contribute to this
172 resilience. Arsenopyrite is monoclinic (Bindi et al., 2012), which means only a limited number
173 of slip systems are available for activation. In addition, arsenopyrites are acicular with a strong

174 shape preferred orientation (SPO). Because there is a competency contrast between
175 arsenopyrite and the quartz-phyllsilicate rich wall rock, at upper greenschist conditions, the
176 SPO allows for accumulation of large strains in the wall rock, while grains aligned with the
177 cleavage remain undeformed (Jessell et al., 2009). We acknowledge that crystal plasticity could
178 be more advanced where arsenopyrites impinge on one another (Barrie et al., 2010), but in this
179 case the percentage of such arsenopyrites was >1% at Obuasi.

180 Furthermore, our results show B-rims with sharp boundaries cutting primary gold
181 zonation, enriched in As and depleted in Au, but preserving the morphology and
182 crystallographic orientation of the parent arsenopyrite. These relationships are indicators that
183 a pseudomorphic dissolution-reprecipitation reaction (Harlov et al., 2011; Putnis, 2009) has
184 resulted in loss of trace element content (gold) from the crystal lattice. The compositional
185 difference between A- and B-rims indicates that re-equilibration to a more stoichiometric
186 composition was the driving force behind the replacement reaction (Geisler et al., 2007).

187 The texture of the gold zoning in the recrystallized domains of grain 2 (Fig. 3), is blurry
188 and diffuse, in apparent contradiction to the sharp oscillatory zoning of the undeformed
189 domains. This texture can be explained by volume diffusion and migration of dislocations
190 during crystal lattice deformation, i.e. dislocation–impurity pair (DIP) diffusion model
191 (Vukmanovic et al., 2014). Critically, in this same domain, the B-rim gold-poor replacement
192 zones penetrate further (up to three times greater), highlighting a link between crystal-plastic
193 microstructure and facilitation of reaction. We suggest that domains of crystal-plastic
194 microstructure (Fig. 3) increased the porosity along high and low angle crystallographic
195 boundaries (Timms et al., 2012), facilitating the reaction by enhanced mobility of components
196 to and from the reaction front (Fig. 4).

197

198 **CONCLUSIONS**

199 The implications of our observations for trace element retention and loss in arsenopyrites
200 are profound.

201 • Arsenopyrite is robust, maintaining its trace element content through conditions of high
202 wall rock strain and metamorphism. This behaviour stands in contrast to other sulphides
203 such as pyrite, characterized by multiple slip systems that can be activated at temperatures
204 as low as 260 °C (Barrie et al., 2011; Reddy and Hough, 2013).

205 • Such robust properties for arsenopyrite contribute to our understanding of its common
206 association with world-class gold deposits. The ability of arsenopyrite to absorb gold into
207 the crystal lattice and as nanoparticles (Cabri et al., 2000) establishes a clear link with
208 mineralisation. Perhaps more importantly, its ability to maintain such trace element content
209 through high strains means that precious metal resources are then preserved in the rock
210 record. In this manner, arsenopyrite acts like an ark, encapsulating and preserving precious
211 metals or trace elements.

212 • Figure 4 presents our model of the microstructural response of arsenopyrite to strain, and
213 associated loss of trace elements (gold). Internal deformation only occurs during
214 crenulation of rock fabrics, when arsenopyrites become aligned with the principal
215 contraction direction. Microfolding is accommodated by localised zones of crystal-plastic
216 deformation, strain-hardening and subsequent microfracture. Where crystal-plastic
217 microstructures are developed, they enhance the release of gold from arsenopyrites
218 undergoing reaction back to the natural environment.

219 • Although the analyses presented here from two arsenopyrites are representative of a large
220 population of arsenopyrites from Obuasi, which we examined petrographically, we
221 recommend future studies of arsenopyrite and other sulphides incorporate EBSD analysis
222 to confirm these results and better understand the role of microstructure in modifying gold
223 deposits.

224

225 **ACKNOWLEDGMENTS**

226 This study derives from Ph.D. research by DF. AngloGold Ashanti is acknowledged
227 for excellent financial support and field assistance. The Centre for Microscopy,
228 Characterisation & Analysis, University of Western Australia, is funded by the University,
229 State and Commonwealth Governments. SM was funded by the Hammond-Nisbet Endowment.

230

231 **REFERENCES**

232 Barrie, C. D., Boyle, A. P., Cook, N. J., and Prior, D. J., 2010, Pyrite deformation textures in
233 the massive sulfide ore deposits of the Norwegian Caledonides: *Tectonophysics*, v.
234 483, no. 3, p. 269-286.

235 Barrie, C. D., Pearce, M. A., and Boyle, A. P., 2011, Reconstructing the pyrite deformation
236 mechanism map: *Ore Geology Reviews*, v. 39, no. 4, p. 265-276.

237 Bindi, L., Moelo, Y., Léone, P., and Suchaud, M., 2012, Stoichiometric arsenopyrite, FeAsS,
238 from La Roche-Balue Quarry, Loire-Atlantique, France: crystal structure and
239 Mössbauer study: *The Canadian Mineralogist*, v. 50, no. 2, p. 471-479.

240 Boyle, A., Prior, D., Banham, M., and Timms, N. E., 1998, Plastic deformation of
241 metamorphic pyrite: new evidence from electron-backscatter diffraction and
242 forescatter orientation-contrast imaging: *Mineralium Deposita*, v. 34, no. 1, p. 71-81.

243 Boyle, R., and Jonasson, I. R., 1973, The geochemistry of arsenic and its use as an indicator
244 element in geochemical prospecting: *Journal of Geochemical Exploration*, v. 2, no. 3,
245 p. 251-296.

246 Cabri, L. J., Newville, M., Gordon, R. A., Crozier, E. D., Sutton, S. R., McMahon, G., and
247 Jiang, D.-T., 2000, Chemical speciation of gold in arsenopyrite: *The Canadian*
248 *Mineralogist*, v. 38, no. 5, p. 1265-1281.

249 Cox, S., 1987, Flow mechanisms in sulphide minerals: *Ore Geology Reviews*, v. 2, no. 1, p.
250 133-171.

251 Dubé, B., Williamson, K., Mcnicoll, V., Malo, M., Skulski, T., Twomey, T., and Sanborn-
252 Barrie, M., 2004, Timing of Gold Mineralization at Red Lake, Northwestern Ontario,
253 Canada: New Constraints from U-Pb Geochronology at the Goldcorp High-Grade
254 Zone, Red Lake Mine, and the Madsen Mine: *Economic Geology*, v. 99, no. 8, p.
255 1611-1641.

256 Eilu, P., and Groves, D. I., 2001, Primary alteration and geochemical dispersion haloes of
257 Archaean orogenic gold deposits in the Yilgarn Craton: the pre-weathering scenario:
258 *Geochemistry: Exploration, Environment, Analysis*, v. 1, no. 3, p. 183-200.

259 Fisher, L. A., Fougrouse, D., Cleverley, J. S., Ryan, C. G., Micklethwaite, S., Halfpenny, A.,
260 Hough, R. M., Gee, M., Paterson, D., and Howard, D. L., 2014, Quantified, multi-
261 scale X-ray fluorescence element mapping using the Maia detector array: application
262 to mineral deposit studies: *Mineralium Deposita*, p. 1-10.

263 Fleck, N. A., Muller, G. M., Ashby, M. F., and Hutchinson, J. W., 1994, Strain gradient
264 plasticity: Theory and experiment: *Acta Metallurgica et Materialia*, v. 42, no. 2, p.
265 475-487.

266 Fougrouse, D., Micklethwaite, S., Tomkins, A. G., Mei, Y., Kilburn, M. R., guagliardo, P.,
267 Fisher, L. A., Halfpenny, A., Gee, M., Paterson, D., and Howard, D. L., In press-a,
268 Gold remobilisation and formation of high grade ore shoots driven by dissolution-
269 reprecipitation replacement and Ni substitution into auriferous arsenopyrite
270 *Geochimica et Cosmochimica Acta*.

271 Fougrouse, D., Micklethwaite, S., Ulrich, S., Miller, J., McCuaig, T. C., Godel, B., and
272 Adams, D., in press-b, Evidence for Two Stages of Mineralization in West Africa's
273 Largest Gold Deposit: Obuasi, Ghana: *Economic Geology*.

274 Geisler, T., Schaltegger, U., and Tomaschek, F., 2007, Re-equilibration of zircon in aqueous
275 fluids and melts: *Elements*, v. 3, no. 1, p. 43-50.

276 Groves, D. I., Goldfarb, R. J., Gebre-Mariam, M., Hagemann, S., and Robert, F., 1998,
277 Orogenic gold deposits: a proposed classification in the context of their crustal
278 distribution and relationship to other gold deposit types: *Ore Geology Reviews*, v. 13,
279 no. 1, p. 7-27.

280 Harlov, D. E., Wirth, R., and Hetherington, C. J., 2011, Fluid-mediated partial alteration in
281 monazite: the role of coupled dissolution–reprecipitation in element redistribution and
282 mass transfer: *Contributions to Mineralogy and Petrology*, v. 162, no. 2, p. 329-348.

283 Hirth, G., and Tullis, J., 1992, Dislocation creep regimes in quartz aggregates: *Journal of*
284 *Structural Geology*, v. 14, no. 2, p. 145-159.

285 Jessell, M. W., Bons, P. D., Griera, A., Evans, L. A., and Wilson, C. J., 2009, A tale of two
286 viscosities: *Journal of Structural Geology*, v. 31, no. 7, p. 719-736.

287 John, T., Klemd, R., Hirdes, W., and Loh, G., 1999, The metamorphic evolution of the
288 Paleoproterozoic (Birimian) volcanic Ashanti belt (Ghana, West Africa): *Precambrian*
289 *Research*, v. 98, no. 1–2, p. 11-30.

290 Kelly, W. C., and Clark, B. R., 1975, Sulfide deformation studies; III, Experimental
291 deformation of chalcopyrite to 2,000 bars and 500 degrees C: *Economic Geology*, v.
292 70, no. 3, p. 431-453.

293 Le Vaillant, M., Barnes, S. J., Fiorentini, M. L., Miller, J., McCuaig, T. C., and Muccilli, P.,
294 2015, A Hydrothermal Ni-As-PGE Geochemical Halo Around the Miitel Komatiite-
295 Hosted Nickel Sulfide Deposit, Yilgarn Craton, Western Australia: *Economic*
296 *Geology*, v. 110, no. 2, p. 505-530.

297 Morelli, R., Creaser, R. A., Seltmann, R., Stuart, F. M., Selby, D., and Graupner, T., 2007,
298 Age and source constraints for the giant Muruntau gold deposit, Uzbekistan, from
299 coupled Re-Os-He isotopes in arsenopyrite: *Geology*, v. 35, no. 9, p. 795-798.

300 Nolze, G., Winkelmann, A., and Boyle, A. P., 2016, Pattern matching approach to
301 pseudosymmetry problems in electron backscatter diffraction: *Ultramicroscopy*, v.
302 160, p. 146-154.

303 Oberthür, T., Vetter, U., Davis, D. W., and Amanor, J. A., 1998, Age constraints on gold
304 mineralization and Paleoproterozoic crustal evolution in the Ashanti belt of southern
305 Ghana: *Precambrian Research*, v. 89, no. 3–4, p. 129-143.

306 Oberthür, T., Vetter, U., Schmidt Mumm, A., Weiser, T., Amanor, J., Gyapong, W., Kumi,
307 R., and Blenkinsop, T., 1994, The Ashanti gold mine at Obuasi, Ghana:
308 Mineralogical, geochemical, stable isotope and fluid inclusion studies on the
309 metallogensis of the deposit: *Geologisches Jahrbuch D*, v. 100, p. 31-129.

310 Prior, D. J., Boyle, A. P., Brenker, F., Cheadle, M. C., Day, A., Lopez, G., Peruzzo, L., Potts,
311 G. J., Reddy, S., and Spiess, R., 1999, The application of electron backscatter
312 diffraction and orientation contrast imaging in the SEM to textural problems in rocks:
313 *American Mineralogist*, v. 84, p. 1741-1759.

314 Putnis, A., 2009, Mineral Replacement Reactions: *Reviews in Mineralogy and Geochemistry*,
315 v. 70, no. 1, p. 87-124.

316 Ramsay, J., and Huber, M., 1983, *The Techniques of Modern Structural Geology, Vol. 1:*
317 *Strain Analysis* Academic Press, London.

318 Reddy, S. M., and Hough, R. M., 2013, Microstructural evolution and trace element mobility
319 in Witwatersrand pyrite: *Contributions to Mineralogy and Petrology*, v. 166, no. 5, p.
320 1269-1284.

321 Schwartz, M. O., Oberthür, T., Amanor, J., and Gyapong, W. A., 1992, Fluid inclusion re-
322 equilibration and P-T-X constraints on fluid evolution in the Ashanti gold deposit,
323 Ghana: *European Journal of Mineralogy*, v. 4, no. 5, p. 1017-1033.

324 Sung, Y. H., Brugger, J., Ciobanu, C. L., Pring, A., Skinner, W., and Nugus, M., 2009,
325 Invisible gold in arsenian pyrite and arsenopyrite from a multistage Archaean gold
326 deposit: Sunrise Dam, Eastern Goldfields Province, Western Australia: *Mineralium*
327 *Deposita*, v. 44, no. 7, p. 765-791.

328 Timms, N. E., Reddy, S. M., Gerald, J. D. F., Green, L., and Muhling, J. R., 2012, Inclusion-
329 localised crystal-plasticity, dynamic porosity, and fast-diffusion pathway generation
330 in zircon: *Journal of Structural Geology*, v. 35, p. 78-89.

331 Urai, J., Means, W., and Lister, G., 1986, Dynamic recrystallization of minerals: *Mineral and*
332 *Rock Deformation: Laboratory Studies: The Paterson Volume*, p. 161-199.

333 Vukmanovic, Z., Reddy, S. M., Godel, B., Barnes, S. J., Fiorentini, M. L., Barnes, S.-J., and
334 Kilburn, M. R., 2014, Relationship between microstructures and grain-scale trace
335 element distribution in komatiite-hosted magmatic sulphide ores: *Lithos*, v. 184, p.
336 42-61.

337

338 FIGURE CAPTIONS

339 Figure 1. A) Boudinaged mineralized quartz vein, Obuasi underground mine; B) Micrograph
340 of mineralized metasedimentary ore, comprising gold-bearing arsenopyrites with quartz strain
341 shadows deformed by a crenulation cleavage; C) Schematic of shear strain calculation
342 estimation and equations used in D (modified after Ramsay and Hubert 1983); D) Frequency
343 of shear strain (e_{stretch}) calculated from 100 strain shadows surrounding arsenopyrites.

344

345 Figure 2. Sample 215-20, grain 1 - the arsenopyrite boundaries, cores and rims are highlighted
346 in red. A) Backscattered electron (BSE) image; B) Cumulative orientation profile along line a-
347 a' (plotted relative to point a) parallel with the arsenopyrite long axis. Only minor
348 crystallographic misorientations are recorded in the grain despite the wall rock being subject
349 to high strain. C) Coloured EBSD maps showing crystallographic misorientations in the range
350 0-10°. Position of orientation profile a-a' is indicated; D) SIMS elemental gold distribution
351 map. The arsenopyrite is zoned with gold-poor core and gold-rich rim (A-rim).

352



353 Figure 3. Sample 215-20, grain 2 - the arsenopyrite boundaries, cores and rims are highlighted
354 in red; A-B) Backscattered electron (BSE) images; C-D) Coloured EBSD maps, showing
355 crystallographic misorientations in the range 0-10° for C and 0-90° for D. Grain 2 coincides
356 with a crenulation microfold and is weakly deformed by this later overprinting low strain event
357 ($D3_{\text{Ob}}$), especially in zone D. High-angle boundaries ($\geq 10^\circ$) are plotted in black and low-angle
358 boundaries ($\geq 2^\circ$) in red. Dynamic recrystallization has produced subgrains and new grains.
359 Position of orientation profiles a-a' and b-b' are indicated; E) SIMS elemental map of ^{197}Au ;
360 F) NanoSIMS composite image of ^{197}Au (yellow) and ^{34}S (blue) elemental distribution. B-rims
361 are gold depleted in comparison to A-rims, and are well developed around zone F; G) Pole

362 figures show the dispersion of orientation data; H) Cumulative orientation profiles (plotted
363 relative to first point) parallel with the long axis (a-a') and short axis (b-b') of grain 2. The
364 largest crystallographic misorientations are recorded along the crystal long axis (a-a'), parallel
365 with the D3 shortening direction.


366

367 Figure 4. Interpretation of the microstructural deformation sequence and replacement of
368 arsenopyrite grain 2.


369



C Stretching estimation: Strain shadow geometry





Shortening estimation: Fold geometry



370

371





373

374

