

1 **Lifetime physical activity and risk of breast cancer in pre-and post-**  
2 **menopausal women**

3 Si Si, PhD<sup>a</sup>, Terry Boyle, PhD<sup>b,c,d</sup>, Jane Heyworth, PhD<sup>e</sup>, Deborah C Glass PhD<sup>f</sup>, Christobel Saunders,  
4 MBBS<sup>g</sup>, Lin Fritschi, PhD<sup>a</sup>,

5

6 <sup>a</sup> School of Public Health, Curtin University, Kent Street, Bentley, Western Australia, Australia

7 <sup>b</sup> Cancer Control Research, British Columbia Cancer Agency, Vancouver, British Columbia, Canada

8 <sup>c</sup> School of Population and Public Health, The University of British Columbia, Vancouver, British Columbia,  
9 Canada

10 <sup>d</sup> Centre for Medical Research, The University of Western Australia, Perth, Western Australia, Australia

11 <sup>e</sup> School of Population Health, The University of Western Australia, 35 Stirling Hwy, Crawley, Western  
12 Australia, Australia

13 <sup>f</sup> Department of Epidemiology and Preventive Medicine Monash University, Melbourne, Victoria, Australia

14 <sup>g</sup> School of Surgery, The University of Western Australia, 35 Stirling Hwy, Crawley, Western Australia,  
15 Australia

16

17 Correspondence to:

18 Si Si

19 e-mail: [si.si@curtin.edu.au](mailto:si.si@curtin.edu.au)

20 Phone: 61-8-92669476

21

22 **Abstract**

23 **Purpose:** To investigate the association between different types of physical activity (PA) and breast  
24 cancer. **Methods:** A case-control study of breast cancer was conducted in Western Australia from  
25 2009-2011, in which 1202 women with breast cancer and 1785 frequency age-matched breast cancer-  
26 free control women were recruited. A self-administered questionnaire was used to collect information  
27 about lifetime and age-period recreational, household, occupational and transport physical activities.  
28 Detailed questions about demographic characteristics, and relevant reproductive, medical and lifestyle  
29 factors were also included. Logistic regression and restrictive cubic spline analyses were applied to  
30 investigate the association and dose-response relationship between PA and breast cancer risk. Sub-  
31 group analysis was performed regarding menopausal status. **Results:** We found non-linear dose-  
32 response associations between PA and risk of breast cancer. Overall, 95-130 MET-hours/week of total  
33 lifetime PA was associated with the lowest breast cancer risk. The effects were stronger among post-  
34 menopausal women. We also found medium amounts of recreational PA (up to 21 MET-hours/week)  
35 were associated with lower breast cancer risk among post-menopausal women. Further analysis on the  
36 intensity of recreational PA demonstrated different dose-response associations between moderate- and  
37 vigorous-intensity recreational PA and breast cancer risk. **Conclusions:** We found that physical  
38 activity was associated with a reduced risk of breast cancer among post-menopausal women, but not  
39 in a linear fashion. Recreational PA of different intensities may have different dose-response  
40 associations with risk of breast cancer.

41 **Key words:** Physical activity; Recreational physical activity; breast cancer; dose-response association

## 42 Introduction

43 Breast cancer is the most commonly diagnosed invasive cancer in Australian women. It was also the  
44 sixth leading cause of burden of disease for females in 2012.[1] The health expenditure on breast  
45 cancer was estimated to be \$331 million in 2004-2005, accounting for 24% of all cancer expenditure  
46 for Australian women that year.[1] Most identified risk factors for breast cancer are non-modifiable in  
47 nature including age, height, family history, hormonal factors and child-bearing histories.[1]  
48 Nevertheless, some modifiable lifestyle-related risk factors for breast cancer have been identified,  
49 including physical activity (PA).[1] A review identified over 73 epidemiologic studies (including  
50 cohort and case-control studies) investigating the associations or dose-response relationships of PA  
51 and the risk of breast cancer worldwide.[2] Although results of these studies were divergent, a slight  
52 majority (51% of all studies) concluded that increasing PA significantly reduced the risk of breast  
53 cancer. Case-control studies, with an average risk reduction of 30%, generally yielded stronger effects  
54 than cohort studies for which the average risk reduction was 20%.[2] Similarly, a meta-analysis of 31  
55 prospective cohort studies yielded an overall 23% risk reduction in breast cancer related to PA.[3]

56 To establish a causal relationship between PA and reduced risk of breast cancer, potential biological  
57 mechanisms have been investigated and verified. It has been argued that PA decreases lifetime  
58 exposure to oestrogen by delaying menarche, reducing the number of ovulatory cycles and ovarian  
59 oestrogen production. [4] Higher levels of other sex hormones including testosterone and  
60 androstenedione have also been associated with an increased risk of breast cancer, especially among  
61 post-menopausal women; and PA might lower testosterone levels. [5] Increasing epidemiological  
62 evidence indicates that adiposity and change of body composition (waist circumference/abdominal fat)  
63 is associated with risk of breast cancer, especially among post-menopausal women. [6-8] Other  
64 possible biological pathways related to PA and risk of breast cancer have also been investigated  
65 including insulin-related factors, adipokines and inflammatory cytokines. However, limited  
66 epidemiological evidence has been found to verify these pathways.[2]

67 Many studies have investigated the dose-response relationship between PA and risk of BC. While  
68 some studies yielded linear association, several studies suggest a ceiling effect of lifetime PA in  
69 reducing risk of breast cancer, in which study participants with the highest level of PA were not the  
70 group at the lowest risk of breast cancer.[9-14] Types of physical activity (e.g. aerobic activity, weight  
71 lifting) may vary with regard to different domains of physical activity (e.g. recreational, occupational  
72 or household PA), therefore physical activity in different domains may have distinctive effects on  
73 breast cancer risk.[9] A meta-analysis of prospective cohort studies yielded 10% to 16% reduction in  
74 breast cancer risks with different domains of PA including recreational, occupational and household  
75 PA.[3] However, the definition and content of domains of PA varied in these studies. [10-13, 15-21]

76 Besides domains of PA, the intensity of PA may also influence the association between physical  
77 activity and breast cancer risk. A review concluded that moderate and vigorous intensity PA are  
78 associated with breast cancer risk reductions in the order of 15% and 18% respectively,[2] while there  
79 is some evidence that light-intensity physical activity may be inversely associated with the risk of  
80 breast cancer.[22] Similarly, the meta-analysis of prospective cohort studies reported 5% and 15%  
81 risk reduction related to moderate and vigorous PA respectively.[3] Even though the evidence seems  
82 to suggest stronger effects of vigorous-intensity PA on breast cancer risk reduction, current  
83 recommendations for recreational physical activities tend to make the assumption that moderate- and  
84 vigorous-intensity PA are related in terms of energy expenditure, with double the amount of  
85 moderate-intensity equivalent to vigorous-intensity PA.[23, 24]

86 The aim of our study was to investigate the relationships between risk of invasive breast cancer and  
87 physical activity including recreational, occupational, transport and household PA and PA of different  
88 intensity. As some breast cancer risk factors may vary for pre- and post-menopausal women (such as  
89 obesity,[2] and since most evidence suggests stronger associations of physical activities and risk of  
90 breast cancer in post-menopausal women,[2] subgroup analyses were performed based on menopausal  
91 status. Additionally, we investigated the dose-response associations of moderate- and vigorous-  
92 intensity recreational PA with risk of breast cancer.

93 **Methods**

94 Exposure to PA was collected as part of the case-control study, Breast Cancer Employment and  
95 Environment Study (BCEES). The details of patients' eligibility and recruitment procedures for this  
96 case-control study are described elsewhere.[25] In brief, women aged between 18 and 80 years with  
97 primary invasive breast cancer diagnosed between May 2009 and January 2011 were identified from  
98 the Western Australia (WA) Cancer Registry. Frequency age-matched control participants, who had  
99 not been diagnosed with invasive breast cancer, were randomly selected from the WA electoral roll  
100 during the same time period.

101 Informed consent was obtained for all study participants. The study was approved by the Human  
102 Research Ethics Committees of The University of WA and the WA Department of Health.

103 ***Data collection***

104 All participants were sent an invitation letter, consent forms and a study questionnaire. Questions  
105 regarding demographic characteristics (age, education level, socio-economic status, remoteness of  
106 residence), reproductive history (pregnancy and breastfeeding history), family history of breast  
107 cancer, lifestyles (alcohol consumption and smoking status), Body Mass Index (BMI) and  
108 reproductive history (menopausal status, age at menarche, oral contraceptive use, Hormone  
109 Replacement Therapy use (HRT)) were included in the questionnaire.[25]

110 Participants were asked to provide information on any job or occupations that they had held for at  
111 least 6 months in their lifetime. Questions included: age started, duration in years, job title, main  
112 duties, employer, industry, country of employment, hours per week, weeks per year worked and their  
113 self-rated intensity of activity (sedentary occupation, standing occupation, manual work and heavy  
114 manual work). This occupational activity question has been shown to have acceptable reliability and  
115 validity when measuring current activity.[26] Additionally, self-rated occupational activity and job-  
116 title based occupational activity have been shown to have very high agreement ( $\kappa = 0.73$ ) in this  
117 study population.[27]

118 A modified version of the Chasan-Taber Physical Activity Questionnaire (CT-PAQ) was used to  
119 assess recreational (walking, swimming, dancing, tennis, aerobics, netball and squash, and up to three  
120 other activities) and household (gardening and household chores) PA, with new questions added to  
121 assess transport-related PA (cycling and walking to/from work).[28] Both the CT-PAQ and the new  
122 transport-related physical activity questions have been shown to have acceptable test-retest  
123 reliability.[28, 29] Recreational, household and transport-related PAs were reported in three different  
124 age periods: 15-24 years, 25-39 years and 40 years above. Questions about age when the PA started  
125 and the number of years, months per year and hours per week undertaking each activity were included  
126 to quantify each PA undertaken.

### 127 *Exposure assessment*

#### 128 *Physical activities*

129 All PAs were assigned a metabolic equivalent (MET) value, derived from the Compendium of  
130 Physical Activities.[30] One MET is defined as the ratio of the metabolic rate for a specific activity  
131 compared to the resting metabolic rate. [30] MET-values were assigned to all measured PA in  
132 recreational, household, occupational and transport domains. For example, walking for exercise was  
133 assigned a MET-value of 4.3; swimming of 6; gardening of 3.8 and cycling to and from work of 7.5.  
134 MET-value of 1.5, 2.3, 3.5 and 6 were assigned to the four categories of occupational activities of  
135 sedentary, standing, manual and heavy manual occupations respectively. The intensity of each PA  
136 was classified as light, moderate or vigorous based on the MET-value assigned to it. Any PA with  
137 assigned MET-value between 1.6 and 3 was labelled light PA; the ones with assigned MET-value  
138 between 3 and 6 were classified as moderate PA; any PA with assigned MET-value 6 and above was  
139 classified as vigorous PA. MET-hours/week of each activity was calculated by multiplying the MET-  
140 value by its frequency and duration. For each age-period and over the lifetime, and for domain-  
141 specific and all physical activity (i.e., the four domains combined), we then calculated mean MET-  
142 hours per week in light-intensity PA, moderate-intensity PA, vigorous-intensity PA and total PA (i.e.,  
143 light, moderate and vigorous PAs combined).

#### 144 Menopausal status

145 Participants were classified as post-menopausal if they self-reported being post-menopausal; or were  
146 aged over 51 years and above and had one of the following self-reported conditions: use of HRT with  
147 regular periods; do not have regular period because of history of hysterectomy or oophorectomy; do  
148 not have regular period because of cancer treatment; or irregular periods (due to stress, endometriosis  
149 or relevant treatment; polycystic ovary; tubal ligation; or other endocrine disorders, etc.) We  
150 conservatively assumed that if women were missing information on whether they had regular periods  
151 and were over age 51 that they were postmenopausal.

#### 152 Potential confounding factors

153 Potential confounding factors collected in the study questionnaire included: age, socio-economic  
154 status (derived from residential postcode and the index of relative socioeconomic advantage and  
155 disadvantage for area [31], index of remoteness of residence (ARIA)[32], education attainment,  
156 family history of breast cancer, age at menarche, age at first pregnancy, breastfeeding history, oral-  
157 contraceptive use in the past 5 years, self-reported hormone replacement therapy use, alcohol  
158 consumption, smoking status and current BMI.

159 Family history of breast cancer was assessed in line with the Australian clinical guidelines:[33] “High  
160 risk” was assigned if the participant reported a first-degree female relative diagnosed with breast  
161 cancer before the age of 50, or two or more first-degree or second-degree female relatives with breast  
162 cancer on the same side of the family; “some family history” was assigned to respondents who  
163 reported any first-degree or second-degree female relative diagnosed with breast cancer at any stage;  
164 and all others were assigned “no family history”.

#### 165 *Data analysis*

##### 166 Data management

167 Those jobs with missing occupational activity intensity were assigned either the level of a similar job  
168 the same participant had held (based on their self-reported job titles and main job duties), or a  
169 physical demands strength rating (based on job-title and duties) from the Dictionary of Occupational  
170 Titles if the participant had not reported the activity level of a similar job.[34] Missing information on

171 hours per week or weeks per year for a job was assigned the median values of the existing/remaining  
172 records (of the same variable) from each individual. The missing duration of a job was calculated by  
173 subtracting the age started at the current job from the age started at the following job.

174 Hours per week were truncated at 14 hours per week for each individual recreational and transport-  
175 related activity, and household chores were truncated at 40 hours per week. If a participant was  
176 missing data for months per year and/or hours per week for an activity, and they had performed the  
177 same activity in a previous or subsequent age-period, the value(s) from that age-period were used. If  
178 they had not performed the same activity in a previous or subsequent age-period, they were assigned  
179 the median value from the study population.

#### 180 Statistical analysis

181 For the total sample, and separately for premenopausal and postmenopausal women in subgroup  
182 analyses, PA variables were categorized into 0 and tertiles of non-zero values of MET-hours/week PA  
183 if there were adequate controls in the 0 MET-hours/week category. Otherwise, quartiles of mean  
184 MET-hours/week of PA based on the distributions of controls were applied to categorize PA types.  
185 Univariate logistic regressions were performed with the potential confounding factors, which  
186 included: demographic characteristics (age, socio-economic status, remoteness of residence and  
187 education attainment); reproductive history (age at menarche, age at first pregnancy, breastfeeding  
188 history and oral-contraceptive use in the past 5 years); medical history (family history of breast cancer  
189 and use of hormone replacement therapy), and lifestyle risk factors (alcohol consumption, smoking  
190 status and current BMI). Variables were later introduced into a multivariate regression model based  
191 upon a conservative p-value of  $<0.25$  in the univariate regression models. A backward stepwise  
192 variable elimination was applied. Independent variables with  $p>0.10$  were removed from the  
193 regression model one at a time. Then, effect modification by risk factors were investigated in the  
194 analysis. Interactions between PA and family history of breast cancer; parity; BMI and ER status were  
195 included in the total PA model. If results indicated significant interaction effects, subgroup analysis  
196 were undertaken.

197 Restricted Cubic Spline (RCS) [35] function (4 knots option) was used to account for the non-linearity  
198 in the investigation of risk of breast cancer and recreational physical activities. In these analyses each  
199 of the PA variables was entered into the model as a continuous MET-hours/week variable rather than  
200 as a categorical variable.

## 201 **Results**

202 In the BCEES study, 58% eligible cases (1202/2084) and 41% eligible controls (1785/4356)  
203 responded to the questionnaire. Overall, controls were slightly older than the cases; and larger  
204 proportion of controls were post-menopausal than cases (77% vs 70%). The characteristics of study  
205 participants were summarized in Table 1. Compared to controls, cases were less likely to be post-  
206 menopausal; have no children and a short breast feeding history; have clear family history of breast  
207 cancer; receive mixed-hormone HRT (Table 1). Furthermore, Over 70% diagnosed breast cancer  
208 tested ER positive in our study sample.

### 209 **Table 1: Characteristics of breast cancer cases and controls**

#### 210 *Description of PA components*

211 Levels of physical activities are summarized in different dimensions in Table 2. Domain-wise,  
212 household and occupational physical activities were the major contributors to total lifetime PA.  
213 Recreational PA accounted for around 20% of lifetime PA among all participants. Intensity-wise,  
214 light, moderate and vigorous PA accounted for 57%, 31% and 12% of lifetime PA respectively. No  
215 significant differences were observed between cases and controls.

### 216 **Table 2: Summary of physical activity measures by breast cancer case and control status**

#### 217 *PA and risk of breast cancer*

218 The results suggested a non-linear association between lifetime total PA and risk of breast cancer.  
219 Women who undertook 95 to 130 MET-hours/week/year PA were at lower risk of breast cancer  
220 compared with participants in the other categories (Table 3), although these differences were not  
221 statistically significant. Higher amounts of PA did not further reduce risk of breast cancer and in fact  
222 the point risk estimate increased in the highest group. A similar pattern was observed among post-  
223 menopausal women, while increasing levels of lifetime all PA seemed to be associated with slightly

224 higher risk of breast cancer in the pre-menopausal sub-group. However, the trend was not statistically  
225 significant.

226 No significant interactions for lifetime total PA and BMI, parity and family history were observed  
227 (Supplementary Table 1). Analysis by ER status was also performed (Supplementary Table 2). The  
228 associations were generally stronger among the ER positive breast cancer cases than the ER negative  
229 cases. However, no significant dose-response associations were observed in any analysis.

230 **Table 3: Adjusted logistic regression analyses for lifetime total physical activities (recreational, household, occupational**  
231 **and transport) and breast cancer**

232 When taking into account domains of lifetime PA, our analysis did not yield significant associations  
233 between risk of breast cancer and either domains or intensity of lifetime PA in the overall analyses.  
234 Sub-group analysis of post-menopausal women yielded a significant association between recreational  
235 PA and breast cancer risk (Table 4). Medium rather than high amounts of recreational PA were  
236 associated with lower risk of breast cancer. Compared to light- and moderate-intensity PA, vigorous-  
237 intensity PA was more relevant for reducing breast cancer risk, especially among post-menopausal  
238 women (4-13 METs-hour/week) (Table 4). No clear risk reduction was observed in the pre-  
239 menopausal subgroup for any domain or intensity and in fact occupational physical activity seemed to  
240 be associated with increased risk.

241 **Table 4: Multiple logistic regressions for different domains and intensities of lifetime physical activity and breast cancer**  
242 **for all participants and stratified by menopausal status**

243 We further examined the associations between recreational PA and breast cancer risk using Restricted  
244 Cubic Spline (RCS) analyses. The RCS demonstrated a complicated pattern between the amount of  
245 lifetime recreational PA and risk of breast cancer. Compared with doing no recreational PA, up to 21  
246 METs hour/week recreational PA was associated with reduced risk of breast cancer (Figure 1).  
247 Recreational PA up to 60 METs hour/week yielded no further risk reduction than to 21 METs  
248 hour/week, while the extrapolation beyond 60 METs hour/week suggested a continuous risk  
249 reduction. However, only a limited number of participants (less than 5%) had than 60 METs  
250 hour/week recreational PA. A similar pattern is demonstrated among post-menopausal women with

251 larger risk reduction effects. Significant risk reduction was observed at level up to 20 METS-  
252 hours/week among post-menopausal women (Figure 1b).

253 **Figure 1: The dose-response analysis of breast cancer risk and recreational physical activity using multivariate restricted cubic splines**  
254 **in all participants (a) and post-menopausal participants (b)**

255 Further analysis of intensity of recreational PA indicated different patterns of dose-response  
256 associations of moderate- and vigorous-recreational PA with risk of breast cancer. A medium amount  
257 of moderate or vigorous recreational PA appeared to be associated with lower risk of breast cancer  
258 (Table 5). RCS further suggested different dose-response associations between moderate and vigorous  
259 intensity recreational PA with risk of breast cancer (Figures 2, 3). Increasing moderate-intensity  
260 recreational PA up to 16 METs-hour/week seemed to be associated with lower risk of breast cancer,  
261 with stronger effects among post-menopausal women. Higher amounts beyond 16 METs-hour/week  
262 were not associated with decreased risk of breast cancer (Figure 2). On the other hand, an increasing  
263 amount of vigorous-intensity recreational PA tended to continuously lower breast cancer risk and  
264 significant risk reduction was observed when the amount is higher than 40 METs-hour/week (Figure  
265 3).

266 **Table 5: Multiple logistic regressions on the intensity of lifetime recreational physical activity and breast cancer risk**  
267 **stratified by menopausal status**

269 **Figure 2: The dose-response analysis of breast cancer risk and moderate-intensity recreational physical activity using multivariate**  
270 **restricted cubic splines in all participants (a) and post-menopausal participants (b)**

271 **Figure 3: The dose-response analysis of breast cancer risk and vigorous-intensity recreational physical activity using multivariate**  
272 **restricted cubic splines in all participants (a) and post-menopausal participants (b)**

## 273 **Discussion**

274 This study investigated the dose-response relationship of lifetime physical activity (in terms of  
275 domains and intensity) and risks of breast cancer in general and further in pre- and post-menopausal  
276 subgroups. Our analyses indicated a small beneficial effect of lifetime PA in reducing risk of breast  
277 cancer overall and in the post-menopausal subgroup, but in a non-linear fashion. We found borderline  
278 significant risk reduction for recreational PA in general, but not for other domains of PA (household,  
279 occupational and transport PA). The association between recreational PA and breast cancer risk was  
280 stronger in the post-menopausal subgroup. The analysis regarding intensity of PAs demonstrated that  
281 compared to light and moderate-intensity PA, increasing vigorous-intensity PA is more relevant to

282 reducing breast cancer risk, especially among post-menopausal women although this is also a non-  
283 linear association. Finally the dose-response analysis of moderate- and vigorous recreational PA  
284 demonstrated distinctive dose-response association patterns with breast cancer risk.

285 We found stronger associations between recreational PA and risk of breast cancer than other domains  
286 of PA. One possible explanation for this is exposure misclassification. Household and occupational  
287 PA, particularly among women, is generally light-intensity, which is recalled less reliably than  
288 moderate- and vigorous-intensity PA.[29, 36] Also, these results corroborate the findings of previous  
289 studies that have measured recreational PA and two or more other PA domains[21, 37-45]; these  
290 previous studies have generally found that physical activity in the recreational domain confers the  
291 largest risk reduction[40-44], although some studies have observed larger risk reductions in the  
292 household domain[37, 39, 45]. We did not find any significant associations between transport PA and  
293 breast cancer risk. However, approximately 40% of all study participants reported no transport PA  
294 and overall it composed less than 3% of lifetime PA. Therefore, the statistical power was low in our  
295 analysis. In terms of intensity of PA, our findings are generally consistent with previous research  
296 suggesting stronger effects of vigorous-intensity PA in reducing risk of breast cancer than other lower  
297 intensity PA.[2] Again, our results demonstrated non-linear associations, especially among post-  
298 menopausal women. No clear associations were observed in the pre-menopausal subgroup in our  
299 study.

300 Our analysis of the recreational PA suggested lower risk of breast cancer among women partaking in  
301 a medium amount (6-26 MET-hours/week) of recreational PA. The literature seems to suggest a linear  
302 correlation between amount of recreational PA and breast cancer. [15, 16, 18] However, there is  
303 significant heterogeneity in the measurement of recreational PA in different studies, with the median  
304 value varying from 9 to 52 MET-hours/week. [9, 13, 15, 17] Therefore, the results in these studies  
305 may not be comparable. We applied Restricted Cubic Spline analysis to further the investigation of  
306 the dose-response associations between recreational PA and breast cancer risk. The results confirmed  
307 a non-linear association. Significant risk reductions were observed among post-menopausal women  
308 undertaking up to 20 MET-hours/week recreational PA. The pattern was consistent with findings of

309 the systematic review of prospective cohort studies, in which a spline demonstrated a relatively linear  
310 reduction in breast cancer risk with increasing amount of recreational PA up to 12 METs-hours/week.  
311 [3] However, the spline did not extend above 12 METs-hours/week. [3] Also, the results were broadly  
312 consistent with the recommendations from World Health Organization (WHO) and the World Cancer  
313 Research Fund. According to the WHO, 150 minutes of moderate-intensity PA or 75 minutes of  
314 vigorous-intensity PA per week is recommended for healthy adults,[24] which is roughly equivalent  
315 to 9 MET-hours/week. Our findings are also consistent with the World Cancer Research Fund  
316 recommendation of 60 minutes moderate-intensity or 30 minutes vigorous-intensity PA on a daily  
317 basis for healthy adults to improve fitness level, which corresponds to approximately 20 MET-  
318 hours/week.[23]

319 Our analysis further suggested different dose-response associations between moderate- (3-6 METs)  
320 and vigorous-intensity ( $\geq 6$  METs) recreational PA with risk of breast cancer respectively.  
321 Consistent with other relevant studies, [9, 13, 15, 17] increasing amount of vigorous-intensity  
322 recreational PA was associated with a decreasing risk of breast cancer, but with diminishing marginal  
323 benefits. Again, the effects were stronger among post-menopausal women. However, the results of  
324 moderate recreational PA suggested decreased breast cancer risk was associated with up to 9 MET-  
325 hours/week. Increasing amounts of moderate-intensity recreational PA (above 17 METs-hour/week)  
326 seemed to be associated with increasing risk of breast cancer. It is possible that the results could be  
327 partially attributed to reporting bias, as research suggests that cancer patients may over-report their  
328 level of PA compared to controls.[46] Further, since only a small proportion (around 10%) of our  
329 study participants self-reported more than 26 MET-hours/week of moderate-intensity recreational PA,  
330 the spline projection at higher levels may not be reliable. Finally, a few studies in the literature have  
331 investigated the association between moderate-intensity recreational PA and risk of breast cancer.[47,  
332 48] Since different definitions and measures were used in these studies to define moderate-intensity  
333 recreational PA, the results were not comparable to ours. Further study is needed to confirm the dose-  
334 response correlations between moderate-intensity recreational PA and risk of breast cancer.

335 Although both the PA guidelines from WHO and the World Cancer Research Fund make the  
336 assumption that moderate- and vigorous-intensity PA are interchangeable in terms of energy  
337 expenditure, with the implication that double amount of moderate-intensity is equivalent to vigorous-  
338 intensity PA.[23, 24] Our analysis indicated that moderate and vigorous-intensity recreational PA may  
339 not be interchangeable in terms of reducing risk of breast cancer. In this study, we found generally  
340 stronger associations between PA and risk of breast cancer among post-menopausal than pre-  
341 menopausal women. The results are consistent with previous research both epidemiological and  
342 biomedical.[2]

### 343 *Strengths*

344 This study had several strengths. We had a large sample size, and had information about a wide range  
345 of potentially confounding variables. We were also able to investigate a number of interaction effects  
346 in this study, including family history of breast cancer, parity, BMI and ER status. However, none of  
347 these variables yielded significant interactions with physical activity. A further strength was having  
348 detailed information about physical activity in four domains and across the lifetime. Finally, our  
349 investigation of PA as a continuous variable and the use of restricted cubic spline analyses better  
350 inform the potential dose-response relationship between different intensity of recreational PA and  
351 breast cancer.

### 352 *Limitations*

353 Our study may be subject to selection bias considering the relatively low response rates. Differences  
354 in age and residential remoteness were found between respondents and non-respondents in cases, and  
355 there was an age difference in controls. However, their potential influence on the amount of PA was  
356 not clear and unlikely to be substantial. Other limitations are associated with the measurement of  
357 physical activities. Self-reported physical activity is subject to reporting bias, especially for PAs in the  
358 early age-periods.[36]. A further limitation of this study was the lack of information regarding dietary  
359 intake. There are no convincing or probable dietary risk factors for breast cancer however, so it  
360 unlikely that controlling for dietary factors would have had a meaningful effect on the observed  
361 associations.[49] Finally, we did not have information about progesterone-receptor status or stage of

362 breast cancer for cases, so we were not able to investigate if the association between PA and breast  
363 cancer risk varied by these clinical characteristics.

### 364 **Conclusion**

365 In this study, we found non-linear associations between physical activity and risk of breast cancer.  
366 The associations between physical activities and breast cancer risk were stronger in post-menopausal  
367 women than pre-menopausal women. Medium amounts of recreational PA among post-menopausal  
368 women were associated with lower risk of breast cancer as was vigorous intensity recreational PA.  
369 Overall, the results of our study supported PA amount recommendations from the WHO and the  
370 World Cancer Research Fund/ American Institute for Cancer Research guidelines for cancer  
371 prevention. Finally, our study informs different dose-response associations of moderate- /vigorous-  
372 intensity recreational PA with breast cancer risks.

### 373 **Acknowledgement**

374 We thank the many West Australian women who participated in the study, Troy Sadkowsky for his  
375 role as data scientist throughout the BCEES project and the members of the BCEES team (Allyson  
376 Thomson, Ann D’Orsogna, Terry Slevin, Jen Girschik and Pierra Rogers). The Breast Cancer  
377 Environment and Employment Study was funded by a National Health and Medical Research Council  
378 Australia (NHMRC) project grant # 572530 and by a grant from the Cancer Council Western  
379 Australia (CCWA). Lin Fritschi is supported by fellowships from the NHMRC and CCWA. Terry  
380 Boyle is supported by an Early Career Fellowship from the NHMRC (#1072266), a Fellowship from  
381 the Canadian Institutes of Health Research (#300068), a Postdoctoral Fellowship from the Michael  
382 Smith Foundation for Health Research (#5553), and an Honorary Killam Postdoctoral Research  
383 Fellowship from The University of British Columbia.

### 384 **Conflict of interest**

385 The authors declare that they have no conflict of interest.

386

388 *Reference*

- 389 1. Bech AG. Breast Cancer in Australia: An Overview: AIHW; 2012.
- 390 2. Lynch BM, Neilson HK, Friedenreich CM. Physical activity and breast cancer prevention.  
391 Physical Activity and Cancer: Springer; 2011. p. 13-42.
- 392 3. Wu Y, Zhang D, Kang S. Physical activity and risk of breast cancer: a meta-analysis of  
393 prospective studies. Breast cancer research and treatment. 2013;137(3):869-82.
- 394 4. Friedenreich CM, Orenstein MR. Physical activity and cancer prevention: etiologic evidence  
395 and biological mechanisms. The Journal of nutrition. 2002;132(11):3456S-64S.
- 396 5. Key T, Appleby P, Barnes I, Reeves G. Endogenous sex hormones and breast cancer in  
397 postmenopausal women: reanalysis of nine prospective studies. Journal of the National Cancer  
398 Institute. 2002;94(8):606-16.
- 399 6. Velthuis MJ, Schuit AJ, Peeters PH, Monninkhof EM. Exercise program affects body  
400 composition but not weight in postmenopausal women. Menopause. 2009;16(4):777-84.
- 401 7. Irwin ML, Yasui Y, Ulrich CM, Bowen D, Rudolph RE, Schwartz RS, et al. Effects of  
402 Exercise on Total and Intra-abdominal Body Fat in Postmenopausal Women. 2003.
- 403 8. Friedenreich C, Woolcott C, McTiernan A, Terry T, Brant R, Ballard-Barbash R, et al.  
404 Adiposity changes after a 1-year aerobic exercise intervention among postmenopausal women: a  
405 randomized controlled trial. International Journal of Obesity. 2011;35(3):427-35.
- 406 9. Holtermann A, Hansen J, Burr H, Sjøgaard K, Sjøgaard G. The health paradox of occupational  
407 and leisure-time physical activity. British Journal of Sports Medicine. 2012;46(4):291-5.
- 408 10. John EM, Horn-Ross PL, Koo J. Lifetime Physical Activity and Breast Cancer Risk in a  
409 Multiethnic Population The San Francisco Bay Area Breast Cancer Study. Cancer Epidemiology  
410 Biomarkers & Prevention. 2003;12(11):1143-52.
- 411 11. Shin A, Matthews CE, Shu X-O, Gao Y-T, Lu W, Gu K, et al. Joint effects of body size,  
412 energy intake, and physical activity on breast cancer risk. Breast cancer research and treatment.  
413 2009;113(1):153-61.
- 414 12. Sprague BL, Trentham-Dietz A, Newcomb PA, Titus-Ernstoff L, Hampton JM, Egan KM.  
415 Lifetime recreational and occupational physical activity and risk of in situ and invasive breast cancer.  
416 Cancer Epidemiology Biomarkers & Prevention. 2007;16(2):236-43.
- 417 13. Schmidt ME, Steindorf K, Mutschelknauss E, Slanger T, Kropp S, Obi N, et al. Physical  
418 activity and postmenopausal breast cancer: effect modification by breast cancer subtypes and effective  
419 periods in life. Cancer Epidemiology Biomarkers & Prevention. 2008;17(12):3402-10.
- 420 14. Leitzmann MF, Moore SC, Peters TM, Lacey Jr JV, Schatzkin A, Schairer C, et al.  
421 Prospective study of physical activity and risk of postmenopausal breast cancer. Breast Cancer Res.  
422 2008;10(5):R92.
- 423 15. Kobayashi LC. Physical Activity Across the Life Course and Risk of Pre-and Post-  
424 menopausal Breast Cancer. 2012.
- 425 16. Peplonska B, Lissowska J, Hartman TJ, Szeszenia-Dabrowska N, Blair A, Zatonski W, et al.  
426 Adulthood lifetime physical activity and breast cancer. Epidemiology. 2008;19(2):226-36.
- 427 17. Friedenreich C, Bryant H, Courneya K. Case-control study of lifetime physical activity and  
428 breast cancer risk. American journal of epidemiology. 2001;154(4):336-47.
- 429 18. Kruk J. Lifetime physical activity and the risk of breast cancer: A case-control study. Cancer  
430 detection and prevention. 2007;31(1):18-28.
- 431 19. Lahmann PH, Friedenreich C, Schuit AJ, Salvini S, Allen NE, Key TJ, et al. Physical activity  
432 and breast cancer risk: the European Prospective Investigation into Cancer and Nutrition. Cancer  
433 Epidemiology Biomarkers & Prevention. 2007;16(1):36-42.
- 434 20. Steindorf K, Ritte R, Eomois PP, Lukanova A, Tjønneland A, Johnsen NF, et al. Physical  
435 activity and risk of breast cancer overall and by hormone receptor status: The European prospective  
436 investigation into cancer and nutrition. International Journal of Cancer. 2013;132(7):1667-78.
- 437 21. Steindorf K, Schmidt M, Kropp S, Chang-Claude J. Case-control study of physical activity  
438 and breast cancer risk among premenopausal women in Germany. American journal of epidemiology.  
439 2003;157(2):121-30.

- 440 22. Kobayashi LC, Janssen I, Richardson H, Lai AS, Spinelli JJ, Aronson KJ. A case-control  
441 study of lifetime light intensity physical activity and breast cancer risk. *Cancer Causes & Control*.  
442 2014;25(1):133-40.
- 443 23. World Cancer Research Fund. Food, nutrition, and the prevention of cancer: a global  
444 perspective. Washington: AICR, 1997.
- 445 24. World Health Organization. Global recommendations on physical activity for health.  
446 Switzerland, Geneva: 2010.
- 447 25. Fritschi L, Erren T, Glass D, Girschik J, Thomson A, Saunders C, et al. The association  
448 between different night shiftwork factors and breast cancer: a case-control study. *British journal of*  
449 *cancer*. 2013;109(9):2472-80.
- 450 26. Cust AE, Smith BJ, Chau J, van der Ploeg HP, Friedenreich CM, Armstrong BK, et al.  
451 Validity and repeatability of the EPIC physical activity questionnaire: a validation study using  
452 accelerometers as an objective measure. *International Journal of Behavioral Nutrition and Physical*  
453 *Activity*. 2008;5(1):33.
- 454 27. Boyle T, Leong S. Comparing ratings of occupational physical activity. *Epidemiology*.  
455 2012;23(6):934-6.
- 456 28. Chasan-Taber L, Erickson JB, McBride JW, Nasca PC, Chasan-Taber S, Freedson PS.  
457 Reproducibility of a self-administered lifetime physical activity questionnaire among female college  
458 alumnae. *American journal of epidemiology*. 2002;155(3):282-91.
- 459 29. Boyle T, Heyworth J, Bull FC, Fritschi L. Test-Retest Reliability of Transport-Related  
460 Physical Activity Performed During the Lifetime. *Journal of Physical Activity and Health*.  
461 2013;10:626-31.
- 462 30. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR, Tudor-Locke C, et al.  
463 2011 compendium of physical activities: a second update of codes and MET values. *Medicine and*  
464 *science in sports and exercise*. 2011;43(8):1575-81.
- 465 31. Australian Bureau of Statistics. Census of Population and Housing: Socio-Economic Indexes  
466 for Areas (SEIFA), Australia 2008. Available from:  
467 [http://www.abs.gov.au/AUSSTATS/abs@.nsf/allprimarymainfeatures/356A4186CCDDC4D1CA257](http://www.abs.gov.au/AUSSTATS/abs@.nsf/allprimarymainfeatures/356A4186CCDDC4D1CA257B3B001AC22C?opendocument)  
468 [B3B001AC22C?opendocument](http://www.abs.gov.au/AUSSTATS/abs@.nsf/allprimarymainfeatures/356A4186CCDDC4D1CA257B3B001AC22C?opendocument).
- 469 32. Department of Health and Aged Care. ARIA (Accessibility/Remoteness Index of Australia).  
470 Available from: [http://www.adelaide.edu.au/apmrc/research/projects/category/about\\_aria.html](http://www.adelaide.edu.au/apmrc/research/projects/category/about_aria.html).
- 471 33. Ackermann E, Harris M, Alexander K, Bailey L, Bennett J, Del Mar C, et al. Guidelines for  
472 Preventive Activities in General Practice. 2013.
- 473 34. U.S. Department of Labor. Dictionary of Occupational Titles 1991.
- 474 35. Desquilbet L, Mariotti F. Dose - response analyses using restricted cubic spline functions in  
475 public health research. *Statistics in medicine*. 2010;29(9):1037-57.
- 476 36. Klesges RC, Eck LH, Mellon MW, Fulliton W, Somes GW, Hanson CL. The accuracy of  
477 self-reports of physical activity. *Medicine & Science in Sports & Exercise*. 1990.
- 478 37. Friedenreich CM, Bryant HE, Courneya KS. Case-control study of lifetime physical activity  
479 and breast cancer risk. *Am J Epidemiol*. 2001 Aug 15;154(4):336-47. PubMed PMID: 11495857.  
480 English.
- 481 38. John EM, Horn-Ross PL, Koo J. Lifetime Physical Activity and Breast Cancer Risk in a  
482 Multiethnic Population: The San Francisco Bay Area Breast Cancer Study. *Cancer Epidemiol*  
483 *Biomarkers Prev*. 2003 November 1, 2003;12(11):1143-52.
- 484 39. Kobayashi L, Janssen I, Richardson H, Lai A, Spinelli J, Aronson K. Moderate-to-vigorous  
485 intensity physical activity across the life course and risk of pre- and post-menopausal breast cancer.  
486 *Breast Cancer Res Treat*. 2013 2013/06/01;139(3):851-61. English.
- 487 40. Kruk J. Intensity of lifetime physical activity and breast cancer risk among Polish women. *J*  
488 *Sports Sci*. 2009 Mar;27(5):437-45. PubMed PMID: 19253081. English.
- 489 41. Kruk J, Kruk J. Lifetime physical activity and the risk of breast cancer: a case-control study.  
490 *Cancer Detect Prev*. 2007;31(1):18-28. PubMed PMID: 17296272. English.
- 491 42. Matthews CE, Shu XO, Jin F, Dai Q, Hebert JR, Ruan ZX, et al. Lifetime physical activity  
492 and breast cancer risk in the Shanghai Breast Cancer Study. *Br J Cancer*. 2001 Apr 6;84(7):994-1001.  
493 PubMed PMID: 11286483. English.

- 494 43. Peplonska B, Lissowska J, Hartman TJ, Szeszenia-Dabrowska N, Blair A, Zatonski W, et al.  
495 Adulthood lifetime physical activity and breast cancer. *Epidemiology*. 2008 Mar;19(2):226-36.  
496 PubMed PMID: 18277160. English.
- 497 44. Tehard B, Friedenreich CM, Oppert J-M, Clavel-Chapelon F. Effect of physical activity on  
498 women at increased risk of breast cancer: results from the E3N cohort study. *Cancer Epidemiol*  
499 *Biomarkers Prev*. 2006 Jan;15(1):57-64. PubMed PMID: 16434587. English.
- 500 45. Lahmann PH, Friedenreich C, Schuit AJ, Salvini S, Allen NE, Key TJ, et al. Physical activity  
501 and breast cancer risk: the European Prospective Investigation into Cancer and Nutrition. *Cancer*  
502 *Epidemiol Biomarkers Prev*. 2007 Jan;16(1):36-42. PubMed PMID: 17179488. English.
- 503 46. Maruti SS, Willett WC, Feskanich D, Levine B, Rosner B, Colditz GA. Physical activity and  
504 premenopausal breast cancer: an examination of recall and selection bias. *Cancer Causes & Control*.  
505 2009;20(5):549-58.
- 506 47. Tehard B, Friedenreich CM, Oppert J-M, Clavel-Chapelon F. Effect of physical activity on  
507 women at increased risk of breast cancer: results from the E3N cohort study. *Cancer Epidemiology*  
508 *Biomarkers & Prevention*. 2006;15(1):57-64.
- 509 48. Verloop J, Rookus MA, van der Kooy K, van Leeuwen FE. Physical activity and breast  
510 cancer risk in women aged 20-54 years. *Journal of the National Cancer Institute*. 2000;92(2):128-35.
- 511 49. World Cancer Research Fund. *Breast Cancer 2010 Report*. 2010.

512

513

514

**Table 1: Characteristics of breast cancer cases and controls**

<b>Variables</b>	<b>Cases (%) (n=1,205)</b>	<b>Controls (%) (n=1,789)</b>	<b>P</b>
<b>Age group (years)</b>			<b>&lt;0.01</b>
< 44	168 (13.9)	179 (10.0)	
45-49	175 (14.5)	203 (11.4)	
50-54	152 (12.6)	237 (13.2)	
55-59	184 (15.3)	306 (17.1)	
60-64	193 (16.0)	322 (8.0)	
65-69	148 (12.3)	272 (15.2)	
70-80	185 (15.4)	270 (15.1)	
<b>Socioeconomic score</b>			0.50
Advantaged 1	54 (4.5)	91 (5.1)	
2	153 (12.7)	245 (13.7)	
3	234 (19.4)	372 (20.8)	
4	248 (20.6)	361 (20.2)	
Disadvantaged 5	515 (42.7)	720 (40.2)	
Missing	1 (0.1)	0	
<b>Remoteness</b>			0.31
Highly accessible	1,033 (85.7)	1,513 (84.6)	
Less accessible/remote	171 (14.2)	276 (15.4)	
Missing	1 (0.1)	0	
<b>Country of birth</b>			0.20
Australia/New Zealand	768 (63.7)	1,191 (66.6)	
UK/Ireland	261 (21.7)	385 (21.5)	
Other Europe	66 (5.5)	86 (4.8)	
Asia	64 (5.3)	67 (3.7)	
Other	46 (3.8)	60 (3.4)	
<b>Education</b>			<b>&lt;0.01</b>
Junior school	436 (36.2)	646 (36.1)	
Senior school	248 (20.6)	403 (22.5)	
Trade/apprenticeship	259 (21.5)	436 (24.4)	
University	262 (21.7)	304 (17.0)	
<b>Age at menarche</b>			0.11
<13 years old	543 (45.1)	737 (41.2)	
≥13 years old	650 (53.9)	1,032 (57.7)	
Missing	12 (1.0)	20 (1.1)	
<b>Parity</b>			<b>0.03</b>
No Child	158 (13.1)	188 (10.5)	
At least 1 Child	1,047 (86.9)	1,601 (89.5)	
Missing	0	0	
<b>Breast feeding history</b>			<b>0.02</b>
No breastfeeding	285 (23.6)	343 (19.2)	
Breastfed≤12 months	394 (32.7)	637 (35.6)	
Breastfed>12 months	507 (42.1)	773 (43.2)	
Missing	19 (1.6)	36 (2.0)	
<b>Menopausal status</b>			<b>&lt;0.01</b>
Pre-menopausal	839 (69.6)	1,368 (76.5)	
Post-menopausal	366 (30.4)	421 (23.5)	
<b>Family History of breast cancer</b>			<b>&lt;0.01</b>
None	729 (60.5)	1,281 (71.6)	
Some family history	302 (25.1)	373 (20.8)	
Clear family history	170 (14.1)	131 (7.3)	
Unknown/missing	4 (0.3)	4 (0.2)	
<b>HRT type</b>			<b>&lt;0.01</b>
No HRT	748 (62.1)	1,035 (57.8)	
Single hormone HRT	157 (13.0)	358 (20.0)	
Mixed hormone HRT	185 (15.4)	224 (12.5)	
Non-hormone/unknown HRT	114 (9.5)	171 (9.6)	
Missing	1 (0.1)	1 (0.1)	
<b>Smoking status</b>			0.39
Never smoked	658 (54.6)	1,022 (57.1)	
Ever smoked	543 (45.1)	761 (42.5)	
Missing	4 (0.3)	6 (0.3)	
<b>Alcohol consumption/week</b>			0.50
None	202 (16.8)	285 (15.9)	

<1 standard drink	138 (11.4)	230 (12.9)	
1-3 standard drinks	278 (23.1)	419 (23.4)	
4-6 standard drinks	148 (12.3)	235 (13.1)	
7-9 standard drinks	120 (10.0)	143 (8.0)	
≥10 standard drinks	315 (26.1)	469 (26.2)	
Missing	4 (0.3)	8 (0.5)	
<b>Current BMI</b>			0.98
Underweight/normal	503 (41.7)	756 (42.3)	
Overweight	388 (32.2)	581 (32.5)	
Obese	286 (23.7)	409 (22.9)	
Missing	28 (2.3)	43 (2.4)	
<b>ER status</b>			
ER positive	858 (71.2)	N/A	
ER negative	170 (14.1)		
Others	177 (14.7)		

*Significant results (p<0.05) were highlighted*

**Table 2: Summary of physical activity measures by breast cancer case and control status**

Variable	Total			Cases			Controls		
	No.	Mean (SD) (MET- hours/week/year)	% (SD)*	No.	Mean (SD) (MET- hours/week/year)	%*	No.	Mean (SD) (MET- hours/week/year)	% (SD)*
Lifetime All PA	<b>2,947</b>	<b>104.7 (56.9)</b>	<b>100</b>	<b>1,183</b>	<b>108.0 (60.2)</b>	<b>100</b>	<b>1,764</b>	<b>102.6 (54.5)</b>	<b>100</b>
Lifetime Household PA	<b>2,951</b>	42.3 (27.5)	43.9 (23.3)	<b>1,183</b>	41.9 (28.2)	42.4 (23.4)	<b>1,768</b>	42.5 (27.0)	44.9 (23.1)
Lifetime Occupational PA	<b>2,983</b>	39.3 (40.1)	32.9 (25.0)	<b>1,200</b>	42.2 (43.3)	34.4 (25.3)	<b>1,783</b>	37.4 (37.6)	31.9 (24.7)
Lifetime Recreational PA	<b>2,971</b>	20.2 (21.3)	20.3 (16.6)	<b>1,193</b>	20.8 (21.3)	20.3(16.1)	<b>1,778</b>	19.8 (21.3)	20.3 (16.9)
Lifetime Transport PA	<b>2,949</b>	2.9 (6.2)	2.9 (5.3)	<b>1,183</b>	3.0 (6.5)	2.9 (5.2)	<b>1,766</b>	2.8 (6.0)	2.8 (5.4)
Lifetime light PA	<b>2,988</b>	55.6 (33.1)	56.8 (23.3)	<b>1,200</b>	55.7 (33.6)	55.7 (23.5)	<b>1,788</b>	55.5 (32.8)	57.5 (23.2)
Lifetime moderate PA	<b>2,988</b>	34.2 (36.1)	30.6 (20.8)	<b>1,200</b>	36.3 (39.8)	31.5 (21.5)	<b>1,788</b>	32.9 (33.3)	30.0 (20.2)
Lifetime vigorous PA	<b>2,988</b>	14.3 (24.5)	12.6 (14.7)	<b>1,200</b>	15.2 (26.6)	12.8 (15.0)	<b>1,788</b>	13.6 (23.0)	12.5 (14.5)

*\*proportion of contribution to the lifetime All PA*

**Table 3: Adjusted logistic regression analyses for lifetime total physical activities (recreational, household, occupational and transport) and breast cancer**

<b>PA (MET-hrs/w/year)</b>	<b>Cases</b>	<b>Controls</b>	<b>OR [95% CI]*</b>
<b>All samples</b>	<b>(n=1205)</b>	<b>(n=1789)</b>	
<b>0-64.6</b>	298 (24.7)	441 (24.6)	1.00
<b>64.7-95.1</b>	267 (22.2)	443 (24.8)	0.92 (0.74, 1.15)
<b>95.2-129.9</b>	259 (21.5)	439 (24.5)	0.90 (0.72, 1.12)
<b>&gt;=130.0</b>	359 (29.8)	441 (24.6)	1.22 (0.98, 1.50)
<b>Missing</b>	22 (1.83)	25 (1.4)	
<b>P<sub>trend</sub></b>			0.02
<b>Post-menopausal Women</b>	<b>(n=839)</b>	<b>(n=1368)</b>	
<b>0-68.5</b>	203 (24.2)	338 (24.7)	1.00
<b>68.6-97.2</b>	196 (23.4)	337 (24.6)	0.98 (0.76, 1.27)
<b>97.3-131.2</b>	171 (20.4)	336 (24.6)	0.84 (0.65, 1.09)
<b>&gt;=131.3</b>	249 (29.7)	335 (24.5)	1.23 (0.97, 1.58)
<b>Missing</b>	20 (2.4)	22 (1.6)	
<b>P<sub>trend</sub></b>			0.03
<b>Pre-menopausal women</b>	<b>(n=366)</b>	<b>(n=421)</b>	
<b>0-50.8</b>	72 (19.7)	105 (24.9)	1.00
<b>50.9-83.5</b>	89 (24.3)	104 (24.7)	1.19 (0.78, 1.81)
<b>83.6-127.0</b>	92 (25.1)	104 (24.7)	1.25 (0.82, 1.91)
<b>&gt;=127.1</b>	111 (30.3)	105 (24.9)	1.46 (0.97, 2.20)
<b>Missing</b>	2 (0.6)	3 (0.7)	
<b>P<sub>trend</sub></b>			0.34

\* adjusted for age, menopausal status, family history of breast cancer, education levels, type of HRT, age at menarche and age at first birth

**Table 4: Multiple logistic regressions for different domains and intensities of lifetime physical activity and breast cancer for all participants and stratified by menopausal status**

PA (MET-hrs/w)	All participants			PA (MET-hrs/w)	Post-menopausal Odds Ratio [95% CI]	PA (MET-hrs/w)	Pre-menopausal Odds Ratio [95% CI]
	Cases (1,205)	Controls (1,789)	Odds Ratio [95% CI]				
<b>Occupational Physical Activities</b>							
<0.1	173 (14.4)	300 (16.8)	1.00	<0.1	1.00	<0.1	1.00
0.1-22.7	328 (27.2)	494 (27.6)	1.15 (0.90, 1.46)	0.1-21.3	1.04 (0.79, 1.37)	0.1-25.1	1.26 (0.77, 2.06)
22.8-52.7	315 (26.1)	496 (27.7)	1.09 (0.85, 1.39)	21.4-49.0	0.98 (0.74, 1.29)	25.2-61.9	1.63 (1.01, 2.64)
>=52.8	384 (31.9)	493 (27.6)	1.30 (1.02, 1.66)	>49.1	1.18 (0.90, 1.56)	>62.0	1.57 (0.96, 2.57)
Missing	5 (0.4)	6 (0.3)					
P <sub>trend</sub>			0.15	P <sub>trend</sub>	0.42	P <sub>trend</sub>	0.16
<b>Household Physical Activities</b>							
0-21.3	326 (27.1)	446 (24.9)	1.00	0-25.7	1.00	0-11.9	1.00
21.4-38.2	304 (25.2)	443 (24.8)	0.97 (0.78, 1.21)	25.8-42.5	0.94 (0.73, 1.21)	12.0-23.9	1.25 (0.83, 1.89)
38.3-60.5	288 (23.9)	443 (24.8)	0.95 (0.76, 1.19)	42.6-65.4	1.01 (0.79, 1.30)	24.0-39.9	1.00 (0.66, 1.52)
>=60.6	267 (22.2)	441 (24.7)	0.91 (0.71, 1.16)	>=65.5	0.92 (0.70, 1.19)	>=40.0	1.03 (0.67, 1.58)
Missing	20 (1.7)	16 (0.9)					
P <sub>trend</sub>			0.89	P <sub>trend</sub>	0.85	P <sub>trend</sub>	0.64
<b>Recreational Physical activities</b>							
0-6.2	302 (25.1)	445 (24.9)	1.00	0-6.2	1.00	0-6.6	1.00
6.3-13.9	264 (21.9)	444 (24.8)	0.84 (0.67, 1.05)	6.3-13.5	0.74 (0.57, 0.96)	6.7-14.7	1.10 (0.72, 1.68)
14.0-25.6	276 (22.9)	446 (24.9)	0.86 (0.69, 1.07)	13.6-25.5	0.83 (0.64, 1.07)	14.8-25.8	1.16 (0.76, 1.76)
>=25.7	351 (29.1)	443 (24.8)	1.08 (0.87, 1.34)	>=25.6	1.05 (0.82, 1.35)	>=25.9	1.24 (0.81, 1.90)
Missing	12 (1.0)	11 (0.6)					
P <sub>trend</sub>			0.06	P <sub>trend</sub>	0.02	P <sub>trend</sub>	0.79
<b>Transport Physical Activities</b>							
<0.1	446 (37.0)	721 (40.3)	1.00	<0.1	1.00	<0.1	1.00
0.1-1.2	235 (19.5)	351 (19.6)	1.05 (0.85, 1.29)	0.1-1.1	1.03 (0.81, 1.32)	0.1-1.3	1.08 (0.72, 1.62)
1.3-4.1	253 (21.0)	350 (19.6)	1.14 (0.92, 1.40)	1.2-4.0	1.05 (0.83, 1.34)	1.4-4.1	1.34 (0.90, 1.99)
>=4.2	252 (20.9)	352 (19.7)	1.14 (0.92, 1.41)	>=4.1	1.03 (0.80, 1.33)	>=4.2	1.34 (0.89, 2.00)
Missing	19 (1.6)	18 (0.8)					
P <sub>trend</sub>			0.53	P <sub>trend</sub>	0.98	P <sub>trend</sub>	0.37
<b>Light-intensity All PA (METs (1.5, 3))</b>							
0-29.5	303 (25.1)	447 (25.0)	1.00	0-32.2	1.00	0-23.3	1.00
29.6-51.5	304 (25.2)	448 (25.0)	1.02 (0.82, 1.27)	32.3-54.6	0.99 (0.77, 1.27)	23.4-43.6	1.04 (0.69, 1.57)
51.6-76.3	280 (23.2)	447 (25.0)	0.95 (0.76, 1.18)	54.7-79.3	0.92 (0.71, 1.18)	43.7-68.8	1.03 (0.68, 1.55)
>=76.4	313 (26.0)	446 (24.9)	1.04 (0.83, 1.29)	>=79.4	1.01 (0.79, 1.30)	>=68.9	1.17 (0.78, 1.77)
Missing	5 (0.4)	1 (0.1)					
P <sub>trend</sub>			0.85	P <sub>trend</sub>	0.87	P <sub>trend</sub>	0.87
<b>Moderate-intensity ALL PA (METs: [3,6])</b>							
0-9.8	271 (22.5)	448 (25.0)	1.00	0-10.4	1.00	0-7.9	1.00
9.9-21.8	308 (25.6)	446 (24.9)	1.16 (0.93, 1.45)	10.5-22.9	1.23 (0.95, 1.59)	8.0-18.4	1.14 (0.75, 1.75)
21.9-44.3	291 (24.1)	447 (25.0)	1.13 (0.91, 1.41)	23.0-46.0	1.11 (0.86, 1.43)	18.5-38.7	1.20 (0.79, 1.85)
>=44.4	330 (27.4)	447 (25.0)	1.27 (1.01, 1.58)	>=46.1	1.24 (0.96, 1.61)	>=38.8	1.44 (0.95, 2.19)
Missing	5 (0.4)	1 (0.1)					
P <sub>trend</sub>			0.21	P <sub>trend</sub>	0.30	P <sub>trend</sub>	0.39
<b>Vigorous-intensity All PA (METs: &gt;=6)</b>							
<0.1	271 (22.5)	448 (25.0)	1.00	<0.1	1.00	<0.1	1.00
0.1-4.1	308 (25.6)	446 (24.9)	0.94 (0.72, 1.21)	0.1-4.2	0.85 (0.64, 1.14)	0.1-4.2	0.99 (0.60, 1.63)
4.2-13.6	291 (24.1)	447 (25.0)	0.85 (0.66, 1.10)	4.3-12.8	0.74 (0.55, 0.99)	4.3-15.1	1.10 (0.67, 1.82)
>=13.7	330 (27.4)	447 (25.0)	0.92 (1.72, 1.19)	>=12.9	0.84 (0.63, 1.13)	>=15.2	0.97 (0.59, 1.61)
Missing	5 (0.4)	1 (0.1)					
P <sub>trend</sub>			0.65	P <sub>trend</sub>	0.26	P <sub>trend</sub>	0.91

Notes: analysis adjusted for age, menopausal status, family history of breast cancer, education levels, type of HRT, age at menarche, age at first birth, breastfeeding history and other types of physical activity

**Table 5: Multiple logistic regressions on the intensity of lifetime recreational physical activity and breast cancer risk stratified by menopausal status**

Recreational PA (MET-hrs/w)	All participants			Post-menopausal participants		Pre-menopausal participants	
	Cases (1,205) No. (%)	Controls (1,789) No. (%)	OR [95% CI]	Recreational PA (MET-hrs/w)	OR [95% CI]	Recreational PA (MET-hrs/w)	OR [95% CI]
<b>Moderate-intensity Recreational PA (METs: [3, 6))</b>							
<0.1	177 (14.7)	225 (12.6)	1.00	<0.1	1.00	<0.1	1.00
<b>0.1-4.3</b>	294 (24.4)	518 (28.9)	0.73 (0.56, 0.94)	<b>0.1-4.3</b>	0.74 (0.56, 0.99)	<b>0.1-3.9</b>	0.63 (0.36, 1.11)
<b>4.4-12.3</b>	345 (28.6)	516 (28.8)	0.84 (0.65, 1.09)	<b>4.4-12.4</b>	0.76 (0.57, 1.02)	<b>4.0-11.1</b>	0.93 (0.54, 1.61)
<b>&gt;=12.4</b>	377 (31.3)	519 (29.0)	0.93 (0.72, 1.20)	<b>&gt;=12.5</b>	0.94 (0.71, 1.25)	<b>&gt;=11.2</b>	0.87 (0.50, 1.51)
Missing	12 (1.0)	11 (0.6)					
<b>P<sub>trend</sub></b>			0.04	<b>P<sub>trend</sub></b>	0.06	<b>P<sub>trend</sub></b>	0.17
<b>Vigorous-intensity Recreational PA (METs: [6, ))</b>							
<0.1	201 (16.7)	270 (15.1)	1.00	<0.1	1.00	<0.1	1.00
<b>0.1-3.4</b>	342 (28.4)	498 (27.8)	0.95 (0.75, 1.22)	<b>0.1-3.4</b>	0.93 (0.71, 1.23)	<b>0.1-3.7</b>	0.91 (0.57, 1.44)
<b>3.5-10.8</b>	300 (24.9)	505 (28.2)	0.79 (0.62, 1.02)	<b>3.5-10.3</b>	0.74 (0.55, 0.98)	<b>3.8-12.8</b>	0.94 (0.59, 1.48)
<b>&gt;=10.9</b>	350 (29.1)	505 (28.2)	0.85 (0.66, 1.08)	<b>&gt;=10.4</b>	0.77 (0.58, 1.02)	<b>&gt;12.8</b>	0.83 (0.52, 1.32)
Missing	12 (1.0)	11 (0.6)					
<b>P<sub>trend</sub></b>			0.18	<b>P<sub>trend</sub></b>	0.09	<b>P<sub>trend</sub></b>	0.87

Note: final model adjusted for: age, menopausal status, family history of breast cancer, education levels, type of HRT, age at menarche, age at first birth, breastfeeding history and other types of physical activity




Figure 1a: All participants

Recreational PA (MET-hours/week)	Median OR (95%CI)
0	1.00
1-21*	0.77 (0.58, 1.02)
22-34	0.95 (0.74, 1.22)
35-50	1.01 (0.78, 1.30)
51-60*	0.97 (0.74, 1.26)
61-80**	0.90 (0.66, 1.23)

Note: knots of spline\*; 5% of the study sample\*\*

Spline analysis adjusted for SES, remoteness, education level, family history of breast cancer, smoking status, age at menarche, parity status, uptake of contraceptive drugs in the last 5 years, breastfeeding history, HRT history, and other types of PA




Figure 1b: Post-menopausal participants

Recreational PA (MET-hours/week)	Median OR (95%CI)
0	1.00
1-9*	0.80 (0.68, 0.93)
10-20*	0.65 (0.47, 0.90)
21-40	0.89 (0.67, 1.19)
41-60*	0.93 (0.69, 1.26)
60-80**	0.76 (0.53, 1.10)

Note: knots of spline\*; 5% of the study sample\*\*

Spline analysis adjusted for SES, remoteness, education level, family history of breast cancer, smoking status, age at menarche, parity status, uptake of contraceptive drugs in the last 5 years, breastfeeding history, HRT history, and other types of PA

Figure 1: The dose-response analysis of breast cancer risk and recreational physical activity using multivariate restricted cubic splines in all participants (a) and post-menopausal participants (b)




Figure 2a: All participants

Recreational MPA (MET-hours/week)	Median OR (95%CI)
0	1.00
1-3*	0.92 (0.84, 1.02)
4-10*	0.83 (0.65, 1.06)
11-16	0.95 (0.76, 1.17)
17-35*	1.12 (0.90, 1.41)
36-50**	1.32 (0.96, 1.83)

Note: knots of spline\*; 5% of the study sample\*\*

Spline analysis adjusted for SES, remoteness, education level, family history of breast cancer, smoking status, age at menarche, parity status, uptake of contraceptive drugs in the last 5 years, breastfeeding history, HRT history, and other types of PA





Figure 2b: Post-menopausal participants

Recreational MPA (MET-hours/week)	Median OR (95%CI)
0	1.00
1-3*	0.89 (0.80, 1.00)
4-10*	0.76 (0.57, 1.01)
11-20	0.90 (0.70, 1.16)
21-35*	1.10 (0.85, 1.43)
36-50**	1.30 (0.91, 1.86)

Note: knots of spline\*; 5% of the study sample\*\*

Spline analysis adjusted for SES, remoteness, education level, family history of breast cancer, smoking status, age at menarche, parity status, uptake of contraceptive drugs in the last 5 years, breastfeeding history, HRT history, and other types of PA

Figure 2: The dose-response analysis of breast cancer risk and moderate-intensity recreational physical activity using multivariate restricted cubic splines in all participants (a) and post-menopausal participants (b)




**Figure 3a: All participants**

Recreational VPA (MET-hours/week)	Median OR (95%CI)
0	1.00
0-9*	0.82 (0.65, 1.03)
10-40*	0.86 (0.69, 1.08)
41-50**	0.76 (0.59, 0.99)

*Note: knots of spline\*; 5% of the study sample\*\**

*Spline analysis adjusted for SES, remoteness, education level, family history of breast cancer, smoking status, age at menarche, parity status, uptake of contraceptive drugs in the last 5 years, breastfeeding history, HRT history, and other types of PA*



**Figure 3b: Post-menopausal participants**

Recreational VPA (MET-hours/week)	Median OR (95%CI)
0	1.00
1-2*	0.86 (0.75, 0.99)
3-8*	0.74 (0.56, 0.97)
9-37*	0.83 (0.64, 1.09)
38-50**	0.68 (0.50, 0.93)

*Note: knots of spline\*; 5% of the study sample\*\**

*Spline analysis adjusted for SES, remoteness, education level, family history of breast cancer, smoking status, age at menarche, parity status, uptake of contraceptive drugs in the last 5 years, breastfeeding history, HRT history, and other types of PA*

**Figure 3: The dose-response analysis of breast cancer risk and vigorous-intensity recreational physical activity using multivariate restricted cubic splines in all participants (a) and post-menopausal participants (b)**

## Supplementary tables

**Table 1: Statistical tests for interaction effect of lifetime physical activity by relevant variables in multivariate logistic regression models**

<b>Variables</b>	<b>P value</b>
Family history of breast cancer	0.9827
BMI	0.2185
Parity	0.4471

*Model adjusted for age, SES, remoteness, education levels, smoking status, age at menarche, use of contraceptive drug in the last 5 years, parity history, breastfeeding history, HRT type, BMI, menopausal status and family history of breast cancer*

**Table 2: Multiple logistic regressions for different domains and intensities of lifetime physical activity and breast cancer for all participants and stratified by ER status**

PA (MET-hrs/w)	No. (%)			Odds Ratio (OR) [95% CI]	
	ER+ (858)	ER- (170)	Controls (1789)	ER+ cases*	ER- cases^
<b>All PA</b>					
0-64.6	201 (23.4)	48 (28.2)	441 (25.0)	1.00	1.00
64.7-95.1	190 (22.1)	36 (21.2)	443 (25.1)	0.97 (0.76, 1.24)	0.76 (0.48, 1.21)
95.2-129.9	188 (21.9)	34 (20.0)	439 (24.9)	0.95 (0.74, 1.22)	0.76 (0.47, 1.21)
>=130.0	261 (30.4)	50 (29.4)	441 (25.0)	1.32 (1.04, 1.67)	1.12 (0.73, 1.72)
Missing	18 (2.1)	2 (1.2)	25 (1.4)		
P	0.03	0.06		0.02	0.23
<b>Occupational Physical Activities</b>					
<0.1	117 (13.6)	19 (11.2)	300 (16.8)	1.00	1.00
0.1-22.7	227 (26.5)	51 (30.0)	494 (27.6)	1.15 (0.88, 1.52)	1.62 (0.93, 2.83)
22.8-52.7	231 (26.9)	43 (25.3)	496 (27.7)	1.16 (0.88, 1.53)	1.38 (0.78, 2.43)
>=52.8	280 (32.6)	56 (32.9)	493 (27.6)	1.37 (1.04, 1.79)	1.63 (0.93, 2.84)
Missing	3 (0.35)	1 (0.6)	6 (0.3)		
P	0.19	0.25		0.14	0.31
<b>Household Physical Activities</b>					
0-21.3	231 (26.9)	51 (30.0)	446 (24.9)	1.00	1.00
21.4-38.2	216 (25.2)	39 (22.9)	443 (24.8)	0.99 (0.78, 1.27)	0.87 (0.55, 1.37)
38.3-60.5	202 (23.5)	44 (25.9)	443 (24.8)	0.96 (0.74, 1.24)	1.00 (0.63, 1.58)
>=60.6	192 (22.4)	34 (20.0)	441 (24.7)	0.97 (0.74, 1.28)	0.89 (0.54, 1.47)
Missing	17 (2.0)	2 (1.18)	16 (0.9)		
P	0.28	0.33		0.99	0.94
<b>Recreational Physical Activities</b>					
0-6.2	211 (24.6)	40 (23.5)	445 (24.9)	1.00	1.00
6.3-13.9	182 (21.2)	48 (28.2)	444 (24.8)	0.85 (0.66, 1.10)	1.15 (0.73, 1.80)
14.0-25.6	200 (23.3)	38 (22.4)	446 (24.9)	0.90 (0.71, 1.16)	0.8 (0.54, 1.41)
>=25.7	256 (29.8)	42 (24.7)	443 (24.8)	1.15 (0.90, 1.47)	1.01 (0.63, 1.62)
Missing	9 (1.1)	2 (1.2)	11 (0.6)		
P <sub>t</sub>	0.15	0.05		0.08	0.72
<b>Transport Physical Activities</b>					
<0.1	323 (37.6)	56 (32.9)	721 (40.3)	1.00	1.00
0.1-1.2	163 (19.0)	30 (17.6)	351 (19.6)	0.98 (0.77, 1.24)	1.08 (0.68, 1.73)
1.3-4.1	173 (20.2)	50 (29.4)	350 (19.6)	1.05 (0.83, 1.33)	1.64 (1.08, 2.48)
>=4.2	184 (21.5)	32 (18.8)	352 (19.7)	1.12 (0.88, 1.42)	1.13 (0.71, 1.82)
Missing	15 (1.8)	2 (1.18)	18 (0.8)		
P	0.29	0.05		0.75	0.11
<b>Light-intensity All PA (METs (1.5, 3))</b>					
0-29.5	216 (25.2)	34 (20.0)	447 (25.0)	1.00	1.00
29.6-51.5	211 (24.6)	49 (28.8)	448 (25.0)	0.99 (0.78, 1.26)	1.53 (0.96, 2.43)
51.6-76.3	194 (22.6)	43 (25.3)	447 (25.0)	0.92 (0.72, 1.18)	1.32 (0.82, 2.13)
>=76.4	234 (27.3)	43 (25.3)	446 (24.9)	1.09 (0.85, 1.38)	1.45 (0.90, 2.35)
Missing	3 (0.4)	1 (0.6)	1 (0.1)		
P	0.31	0.29		0.61	0.31
<b>Moderate-intensity ALL PA (METs:[3,6])</b>					
0-9.8	185 (21.6)	47 (27.7)	448 (25.0)	1.00	1.00
9.9-21.8	225 (26.2)	40 (23.5)	446 (24.9)	1.26 (0.99, 1.61)	0.87 (0.55, 1.36)
21.9-44.3	205 (23.9)	50 (29.4)	447 (25.0)	1.18 (0.92, 1.51)	1.06 (0.69, 1.64)
>=44.4	240 (28.0)	32 (18.8)	447 (25.0)	1.41 (1.10, 1.81)	0.67 (0.41, 1.10)
Missing	3 (0.3)	1 (0.6)	1 (0.1)		
P	0.19	0.01		0.05	0.27
<b>^Vigorous-intensity All PA (METs: &gt;=6)*</b>					
<0.1	116 (13.5)	25 (14.7)	448 (25.0)	1.00	1.00
0.1-4.1	241 (28.1)	45 (26.5)	446 (24.9)	0.96 (0.72, 1.27)	0.84 (0.50, 1.42)
4.2-13.6	235 (27.4)	44 (25.9)	447 (25.0)	0.89 (0.68, 1.18)	0.83 (0.49, 1.41)
>=13.7	263 (30.6)	55 (32.4)	447 (25.0)	0.91 (0.68, 1.21)	0.97 (0.59, 1.62)
Missing	3 (0.4)	1 (0.6)	1 (0.1)		
P	0.17	0.30		0.85	0.82

Note: \*final model adjusted for: age, menopausal status, education level, family history, parity status, breastfeeding history, family history of breast cancer, types of HRT and other domains of PA

^ final model adjusted for: age, menopausal status, smoking status, family history of breast cancer, age at menarche, contraceptive drug in 5 years and PA of other intensity