

Current Issues in Plant Cryopreservation

Anja Kaczmarczyk^{1,2}, Bryn Funnekotter^{1,2}, Akshay Menon^{1,2},
Pui Ye Phang^{1,2}, Arwa Al-Hanbali^{1,2}, Eric Bunn^{2,3} and Ricardo L. Mancera¹

¹*Curtin Health Innovation Research Institute, Western Australian Biomedical Research
Institute, Curtin University, Perth*

²*Botanic Gardens and Parks Authority, Fraser Avenue, West Perth*

³*School of Plant Biology, Faculty of Natural and Agricultural Sciences,
University of Western Australia
Australia*

1. Introduction

Plant cryopreservation involves the storage of plant tissues (usually seed or shoot tips) in liquid nitrogen (LN) at -196°C or in the vapour phase of LN at -135°C in such a way that the viability of stored tissues is retained following re-warming (Day et al., 2008; Hamilton et al., 2009). Cryopreservation is usually applied to species with recalcitrant (i.e. dehydration sensitive) seeds that are not storable by any other means, or preservation of specific cultivars of vegetatively propagated crop plants like banana or potato, or for unique ornamental genotypes (Halmagyi et al., 2004; Kaczmarczyk et al., 2011a; Panis et al., 2005). Another reason to utilise cryostorage is to conserve endangered plant species, particularly where seeds may be extremely scarce or of doubtful quality and/or the species is threatened with imminent extinction (Decruse et al., 1999; Mallon et al., 2008; Mandal & Dixit-Sharma, 2007; Paunescu, 2009; Sen-Rong & Ming-Hua, 2009; Touchell et al., 2002).

The main advantage of cryopreservation is that once material has been successfully cooled to LN temperatures, it can be conserved in principle indefinitely, because at these ultra-low temperatures no metabolic processes occur. Replenishing a small volume of LN weekly in cryo-dewars is the only on-going maintenance operation usually required in cryostorage. There are further advantages to this approach: the low costs of storage, minimal space requirements and reduced labour maintenance compared to living collections and even when compared to maintenance of tissue cultures at room temperature. Once in storage, there is no risk of new contamination by fungus or bacteria, and cryogenically stored material has been reported to retain genetic stability (Harding, 2004). Depending on the species, small cryopreserved samples may take several weeks to re-establish shoot cultures, and several months to a year may be required to produce micropropagated plants capable of transfer to soil under greenhouse conditions and (following weaning) into the field.

Shoot tips (containing the apical meristem) are the most commonly used plant material for cryostorage. The apical meristem is composed of small unvacuolated cells served with a relatively small vascular system. The organised structure of apical meristems generally results in direct shoot formation after re-warming, thereby maintaining the genetic integrity

of the resulting propagated material. While callus tissue (unorganised wound tissue) can also be cryostored, the risk of occurrence of genetic deviations may be higher when utilising the indirect organogenesis pathway. Besides shoot tips, callus cultures, cell cultures, somatic embryos, pollen or plant buds as well as recalcitrant and orthodox seeds can be used as explants in plant cryopreservation (Reed, 2008).

Plant cryopreservation began with research on the freezing of mulberry twigs in LN (Sakai, 1965). Since then, a huge variety of plants and genotypes have been successfully cryostored for conservation of agriculture and horticultural genotypes, as well as for endangered and threatened plant species (Gonzalez-Arno et al., 2008; Hamilton et al., 2009; Kaczmarczyk et al., 2011b; Mycock et al., 1995; Reed, 2008; Sakai & Engelmann, 2007). This chapter reviews and gives examples of different plant cryopreservation protocols that have been successfully applied. It will focus on free radical damage and membrane structure, both important topics in the cryopreservation of biological tissues. The topic of genetic and epigenetic stability in plant cryopreservation is also discussed. Recent reviews of plant cryopreservation have been written by Benson (2008), Day & Stacey (2007), Hamilton et al. (2009) and Reed (2008).

2. Plant cryopreservation methods

In the last three decades a number of different cryopreservation protocols, such as classical slow-cooling, vitrification, droplet vitrification, encapsulation/dehydration and encapsulation/vitrification protocols have been developed and utilised for germplasm storage (Reed, 2008). The choice of cryopreservation method to attain the highest survival rates is largely dependent on the plant species and tissue type that is being cryostored.

2.1 Slow cooling or controlled rate cooling

This technique involves the simple dehydration of plant material before cryogenic storage in LN. This can be done by slow cooling of the plant tissue to a temperature of approximately -40°C (Reinhoud et al., 2000). This forces the formation of extracellular ice ahead of intracellular ice, thus causing an outflow of water from the cells due to the resulting osmotic imbalance and, consequently, dehydration. Dehydration can also be brought about by incubation of tissue material on media containing a relatively high concentration of an osmoregulant, commonly sucrose, although other compounds can also be used (Panis et al., 2002). Usually water concentrations must be decreased to between 10% and 20% of the fresh weight for optimal cryogenic survival (Engelmann, 2004). This has the aforementioned effects of reducing the extent of ice crystal formation due to the reduced water concentration and assisting in the achievement of the vitrified state of water as a result of the increased solute concentration. These techniques do not necessarily make use of cryoprotective agents (CPAs), however they can be used in conjunction with them to further improve dehydration (Reinhoud et al., 2000), though these agents can be toxic to plant cells at high concentrations (Arakawa et al., 1990). Rapid re-warming rates are used after cryogenic storage to prevent ice crystal formation during thawing (Reinhoud et al., 2000). This approach can result in extreme rates of dehydration, which can cause cell volume reductions that are potentially lethal (Day et al., 2008). It has been suggested that slow-cooling is only suitable for non-organised tissues, as sufficient dehydration is more difficult to achieve in tissues with complex structures due to the different rates of water movement between and within plant cells with different characteristics (Gonzalez-Arno et al., 2008).

2.2 Encapsulation-dehydration

This method, developed by Fabre and Dereuddre (1990), involves encapsulating shoot tips, somatic embryos or callus cells within alginate beads. This is followed by incubation in media with high sugar concentrations in order to raise intracellular solute concentrations and promote desiccation. Finally, silica gel or airflow is used to dehydrate the beads until the moisture content drops to 20-30%, before they are immersed in LN (Fabre & Dereuddre, 1990; Hamilton et al., 2009; Reinhoud et al., 2000). The encapsulating material is thought to promote a vitrified state in the tissue regardless of the cooling and re-warming rates, thus reducing damage from ice crystal formation (Scottetz et al., 1992). Mechanical stress is also reduced because the bead protects the explants from damage during handling. The benefits of this method include avoiding the use of high concentrations of (potentially toxic) CPAs (Reinhoud et al., 2000) and the presence of a nutritive bead, which may enhance post-regeneration survival or re-growth of the material following immersion in LN and re-warming (Panis & Lambardi, 2005).

2.3 Vitrification

Vitrification involves the treatment of tissues in a mixture of highly concentrated penetrating and non-penetrating CPAs applied at non-freezing temperatures, followed by rapid cooling in LN (Gonzalez-Arno et al., 2008; Panis & Lambardi, 2005). The combination of high intracellular solute concentrations (due to dehydration and some CPA penetration) and rapid cooling prevents the nucleation of water and the formation of ice crystals, both intracellularly and extracellularly, thus promoting the vitrification of water (Kreck et al., 2011; Mandumpal et al., 2011; Reinhoud et al., 2000).

For plants sensitive to direct exposure to vitrification solutions, due to dehydration intolerance and osmotic stresses, a loading step of 10-20 minutes can be incorporated prior to incubation within the CPA solution. This is done by incubation of the samples within a less toxic/concentrated CPA solution (a media containing 0.4 M sucrose and 2 M glycerol proved highly effective [Nishizawa et al., 1993; Sakai et al., 1990]), thereby improving dehydration tolerance. The CPAs used in vitrification usually contain high concentrations of glycerol, dimethyl sulfoxide (DMSO), ethylene glycol and various sugars (Day et al., 2008). The most commonly used mixture of CPAs for vitrification is plant vitrification solution 2 (PVS2), which consists of 30% (w/v) glycerol, 15% (w/v) ethylene glycol and 15% (w/v) DMSO in basal culture medium containing 0.4 M sucrose (Sakai et al., 1990, 2008).

Exposure time to cryoprotective solutions is a vital step in the cryostorage process. Volk and Walters (2006) demonstrated that the extent of penetration of PVS2 into mint and garlic shoot tips was directly proportional to exposure time. The water content of the shoot tips also significantly decreased with an increase in exposure time to PVS2 (Volk & Walters, 2006). Greater penetration of CPAs can be useful as it helps to increase the internal solute concentration and may contribute to maintaining cell volume, thus preventing damage to the cells (Meryman, 1974). However, overexposure to CPAs may cause damage to the cells owing to the toxic nature of the CPAs or excessive dehydration.

The vitrification protocol is a more widely applied cryopreservation method than slow cooling due to its ease of use, high reproducibility and the wide range of species with complex tissue

structure (such as shoot tips and embryos) that have been successfully cryopreserved with this procedure (Takagi et al., 1997; Touchell et al., 1992; Vidal et al., 2005).

2.4 Droplet-vitrification

The droplet-vitrification technique is a modification of the basic vitrification protocol that involves placing the sample within a droplet of 1-10 μl of cryoprotective solution on a piece of aluminium foil before immersion in LN (as opposed to 1-2 mL of cryoprotective solution in the original protocol), as shown in Fig. 1.




Fig. 1. Cryopreservation procedure for the droplet vitrification method.

This approach achieves higher cooling and re-warming rates, as the small volume of liquid allows a higher rate of heat transfer to and from the sample (Sakai & Engelmann, 2007). Cooling rates are increased to $>130^{\circ}\text{C}/\text{sec}$ (Panis & Lambardi, 2005), therefore facilitating the direct transition of intracellular water from a liquid state to a glassy state far more rapidly, thus minimising water crystallisation. The droplet-vitrification protocol has been successfully applied in the cryopreservation of garlic and chrysanthemum (Kim et al., 2009), yams (Leunufna & Keller, 2005), lily (Chen et al., 2011), potato (Yoon et al., 2006) and other plants (Sakai & Engelmann, 2007).

2.5 Encapsulation-vitrification

Another modification of the vitrification approach termed encapsulation/vitrification combines elements of the encapsulation/dehydration method with the vitrification method. As with the standard encapsulation method, shoot tips or calluses are first encapsulated in alginate beads and then the encapsulated material is incubated in a vitrification solution to promote sufficient dehydration and vitrification rather than dehydration under a constant airflow, which is very time consuming and relatively imprecise (Hirai & Sakai, 1999). Successful protocols have been established for potato (Hirai & Sakai, 1999), gentian (Tanaka et al., 2004), strawberry (Hirai et al., 1998) and pineapple (Gamez-Pastrana et al., 2004).

3. Free radical and oxidative damage in plant cryopreservation

While achieving an optimum cryopreservation protocol that successfully avoids ice damage is important, there are various other factors that can affect post-cryogenic survival (Fig. 2). During the cryopreservation procedure plant tissues are susceptible to a variety of stresses, including oxidative stresses (Benson & Bremner, 2004).




Fig. 2. Main causes of damage to plant tissues during cooling and cryopreservation (modified from [Turner, 2001]).

3.1 Reactive oxygen species (ROS)

The formation of reactive oxygen species (ROS) during cryopreservation can occur during the many steps involved in this process. For example, ROS formation has been detected in photo-oxidative stress during tissue culture, during excision of shoot apices, osmotic injury and desiccation following application of CPAs, as well as during the rapid changes in temperature when the samples are first cryostored in LN and then re-warmed (Benson & Bremner, 2004; Roach et al., 2008). ROS are highly reactive molecules that can cause a wide range of damage in cells. There is a large variety of molecules that are classified as ROS, some of which include oxygen-free radical species and reactive oxygen non-radical derivatives (Table 1).


Radicals	Non-Radicals
Superoxide ($O_2^{\bullet-}$)	Hydrogen peroxide (H_2O_2)
Hydroxyl (OH^{\bullet})	Peroxynitrite ($ONOO^{\bullet-}$)
Hydroperoxyl (OOH^{\bullet})	Peroxynitrous acid ($ONOOH$)
Peroxyl (ROO^{\bullet})	Hypochlorous acid ($HOCl$)
Alkoxy (RO^{\bullet})	Hypobromous acid ($HOBr$)
Carbonate ($CO_3^{\bullet-}$)	Ozone (O_3)
Carbon dioxide ($CO_2^{\bullet-}$)	Singlet oxygen ($^1\Delta g$)
Singlet oxygen ($^1\Sigma g^+$)	

Table 1. Common reactive oxygen species (ROS) (Halliwell & Gutteridge, 2007).

Many of the more active ROS are free radicals, which are molecules that contain an unpaired electron, thus being able to react non-specifically with neighbouring molecules by removing electrons and causing a self-propagating chain reaction of radical formation. The removal of electrons can lead to a loss of function and structural alterations in macromolecules like proteins, lipids and DNA (Benson, 1990; Halliwell & Gutteridge, 2007). ROS are frequently produced as by-products during cellular metabolism. The electron transport chain used in respiration and photosynthesis are the major producers of ROS, caused by the leakage of free electrons onto molecular oxygen (O_2), resulting in the formation of superoxide (Benson & Bremner, 2004; Benson, 1990; Halliwell & Gutteridge, 2007). The formation of ROS is controlled by a high concentration of antioxidants and proteins that can quench the ROS and fix the damage in these regions. Nevertheless, if there is a sudden increase in ROS formation then cellular repair processes can be overwhelmed and excessive damage can occur.

Temporarily reducing cryo-sample exposure to light immediately after cryopreservation has been shown to increase post-cryogenic survival due to the removal of photo-oxidative stress in the plants (Senula et al., 2007; Touchell et al., 2002). Photo-oxidative stress in plants can result in high levels of singlet oxygen ($^1\Sigma_g^+$) and superoxide ($O_2^{\cdot-}$) being produced, either from direct UV radiation on oxygen or the leakage of light energy onto oxygen from chlorophyll when the carotenoid pigments become saturated (Wise, 1995). Plants are highly susceptible to photo-oxidative stresses at low temperatures when exposed to strong light conditions. This can be demonstrated in alpine plants, many of which display adaptations (especially to leaves, i.e. the production of carotenoid pigments) that reduce photo-oxidation damage from enhanced UV-B radiation at high altitudes (Streb et al., 1998). When the ability of antioxidants to quench the formation of ROS and the recycling of antioxidants is reduced, greater damage occurs to the chloroplast through lipid peroxidation, inactivation of photosynthetic proteins and loss of pigments (i.e. bleaching) (Wise, 1995; Wise & Naylor, 1987). Damage to the chloroplast during chilling stress has been shown to severely impede growth rates (Partelli et al., 2009); therefore, reducing the damage that occurs to plant cells due to low temperature oxidative stress is vitally important for improving survival and recovery in cryopreservation.

The most reactive ROS commonly found in plants include superoxide ($O_2^{\cdot-}$), the hydroxyl radical ($OH\cdot$), hydroperoxyl ($OOH\cdot$) and singlet oxygen ($^1\Sigma_g^+$). Superoxide and singlet oxygen are often formed as by-products of the electron transport chain from both metabolism and photosynthesis, while hydroperoxyl and hydroxyl radicals are commonly formed in a process called Fenton's reaction, where hydrogen peroxide is converted into the hydroxyl or hydroperoxyl radical (1).



The formation of the hydroxyl radical is the major cause of lipid peroxidation in membranes, but can also cause a wide range of damage to all cellular components, including proteins and DNA (Halliwell & Gutteridge, 2007). The addition of specific chelating agents (such as desferrioxamine) has been shown to reduce the levels of iron in cryopreserved tissues, with subsequent decreased levels of the hydroxyl radical observed (Benson et al., 1995; Fleck et al., 2000; Obert et al., 2005).

Damage caused by ROS is difficult to quantify as these molecules are non-specific in their interactions, reacting freely with lipids, proteins and DNA. ROS are highly reactive (and

therefore short-lived) and thus, direct measurement of the ROS present in cells is difficult due to time constraints, and does not reflect the damage that may be done prior to the ROS being quenched by antioxidants. Consequently, it is easier to measure the formation of by-products of oxidative damage or the antioxidant status of the cells. The ratio of oxidised to reduced antioxidants is a good indication of ROS formation and the ability of cells to regulate oxidative stress. Identifying end products of ROS oxidation is an indication of the damage caused and is a sign that cells have been unable to satisfactorily quench ROS activity.

3.2 Lipid peroxidation

The cell membrane represents one of the major areas where cryo-injury can occur. Any damage to the cell membrane can alter the delicate balance between intra and extracellular solutes, leading to cell death (Anchordoguy et al., 1987; Dowgert & Steponkus, 1984; Gordon-Kamm & Steponkus, 1984; Lynch & Steponkus, 1987). Lipid peroxidation of fatty acids (FA) in phospholipids can cause extensive damage to the cell membrane if the chain reaction is not controlled, leading to large areas where the semi-permeability of the membrane is altered and thus can no longer function normally (Benson et al., 1992; Halliwell & Gutteridge, 2007). Lipid peroxidation is caused when specific ROS (hydroxyl radical, peroxy radical and singlet oxygen) interact with a FA. Polyunsaturated fatty acids (PUFA) are the most susceptible to peroxidation (Møller et al., 2007; Young & McEneny, 2001). Glutathione peroxidase can detoxify lipid peroxides by reducing them back to their lipid alcohol form and, in the process, oxidising glutathione (Benson, 1990). The formation of the toxic end-products of lipid peroxidation, such as malondialdehyde (MDA) and 4-hydroxynonenal (HNE), can also cause damage to cells (Halliwell & Gutteridge, 2007). The toxicity of MDA is debatable (Halliwell & Gutteridge, 2007); however, there is evidence that MDA can interact with proteins and DNA, causing loss of function in proteins and mutations in DNA (Halliwell & Gutteridge, 2007; Hipkiss et al., 1997; Marnett, 1999). The formation of HNE has shown greater toxicity to cells as it can damage mitochondria, inhibit synthesis of DNA and proteins, and interfere with the action of repair proteins such as chaperones (Halliwell & Gutteridge, 2007). Identifying the formation of MDA and HNE is commonly used in cryopreservation as an indicator of oxidative stress. High levels of MDA or HNE detected have correlated with decreased survival rates in rice cell suspensions, olive somatic embryos, flax and blackberry shoot tips (Benson et al., 1992; Obert et al., 2005; Uchendu et al., 2010; Lynch et al., 2011).

Volatile headspace sampling (VHS) measures the formation of volatile compounds released from a sample in a non-destructive and non-invasive assay method. This provides an important tool for measuring oxidative stress. Free radical damage can cause the formation of volatile compounds such as methane, ethane, ethylene and pentane. Quantification of these compounds can be indicative of oxidative damage. The detection of ethylene in plants is of particular importance as ethylene is a vital hormone. Fang et al. (2008) observed that decreased levels of ethylene production correlated with decreased survival and growth. The production of the other volatile compounds is indicative of lipid peroxidation, where increased levels correlate to excessive oxidative damage. Benson et al. (1987) observed a large increase in volatile compounds produced after thawing that has since been observed in other species.

The use of DMSO as a free radical scavenger and probe for the hydroxyl radical can be utilised with VHS. The formation of methane when DMSO interacts with the hydroxyl radical can be measured if the sample is placed in an airtight container, with the levels of methane detected correlating with the formation of the hydroxyl radical. This technique has been used as a measurement of oxidative stress in multiple different plant species, such as rice, cocoa, *Daucus carota* and flax (Benson & Withers 1987; Benson et al., 1995; Obert et al., 2005; Fang et al., 2008). The production of methane is particularly strong during the initial phase of recovery, where it is predicted that antioxidant activity and production is reduced, resulting in increased ROS (Fang et al., 2008). The use of chelating agents to reduce the formation of hydroxyl radicals from the Fenton reaction has shown significant benefits in plant cryopreservation. Desferrioxamine is the most common chelating agent used. It binds to and reduces the amount of free iron. The addition of desferrioxamine to rice cells during cryopreservation showed a decreased recovery period after cryopreservation (Benson et al., 1995). Detection of methane formation was delayed in flax tissue when exposed to desferrioxamine (Obert et al., 2005), indicating a delayed production of hydroxyl radicals.

3.3 Antioxidants

An antioxidant can be defined as any molecule that “delays, prevents or removes oxidative damage to a target molecule” (Halliwell & Gutteridge, 2007). Antioxidants can be classified into two major groups: enzymes that catalytically remove ROS, or sacrificial antioxidant molecules that are preferentially oxidised to protect more important molecules by quenching ROS (Halliwell & Gutteridge, 2007). The addition of exogenous antioxidants during cryopreservation has been shown to result in increased survival in some cases (Uchendu et al., 2010; Wang & Deng, 2004). It is possible that the addition of exogenous antioxidants may also aid in reducing oxidative stresses immediately following re-warming, when cellular metabolism has not been restored to its original state.

Glutathione (GSH) is one of the main water soluble sacrificial antioxidants in plants (Kranter et al., 2006). This low molecular weight thiol is converted to its oxidised form (GSSG) upon interaction with ROS. The ratio of reduced to oxidised GSH is a good indicator of oxidative stress, as an increased amount of GSSG indicates the inability of the plant to control oxidative damage, thereby triggering premature cell death (Kranter et al., 2006). GSH is recycled by the enzyme GSH reductase using NADPH as the electron donor (Halliwell & Gutteridge, 2007). Ascorbic acid is the most abundant water-soluble antioxidant in plant cells (Foyer & Noctor, 2009). This antioxidant is able to quench free radicals, forming the stable radical semidehydroascorbate, which can be further oxidised to dehydroascorbate by another free radical. *Arabidopsis* mutants, which were unable to express ascorbic acid activity, demonstrated the important role this antioxidant plays in reducing oxidative stress, as these plants were not viable following exposure to photo-oxidative stress (Dowdle et al., 2007). Tocopherol (Vitamin E) is a lipophilic antioxidant. This is the main antioxidant involved in membrane protection against lipid peroxidation, as it is preferentially oxidised instead of PUFAs (Halliwell & Gutteridge, 2007; Uchendu et al., 2010). This antioxidant is thought to be reduced to its original, functional state by ascorbic acid (Packer et al., 1979). Carotenoid pigments are the main defence in chloroplasts, where large amounts of singlet oxygen can be produced if the activated chlorophylls transfer their energy onto oxygen. The carotenoid pigments can absorb the energy directly from the

chlorophylls, thus suppressing the formation of singlet oxygen, and they can also quench the singlet oxygen directly (Halliwell & Gutteridge, 2007).

The enzyme superoxide dismutase (SOD) catalytically removes the ROS superoxide, producing oxygen and hydrogen peroxide. The removal of superoxide is more important than the formation of hydrogen peroxide, as superoxide is a more reactive species, causing wider damage in the cells. There is potential for SOD to cause formation of ROS if levels of hydrogen peroxide are not controlled. SOD contains a metal cofactor that can cause Fenton reactions and the formation of the hydroxyl radical. Catalase is the main enzyme involved in removing hydrogen peroxide, resulting in the decomposition of hydrogen peroxide to water and oxygen. This enzyme is vital for the removal of hydrogen peroxide before it can damage the cell or be converted to the highly reactive hydroxyl radical through Fenton's reaction.

4. Plant cryopreservation and membrane structure

Membrane systems within cells are usually the site of freezing injury in plants (Steponkus, 1984). Membrane stability is therefore important for reducing such injury. There are four types of injury: (i) expansion-induced lysis, where the cells overexpand as a result of increased extracellular osmotic pressure during warming/thawing; (ii) loss of osmotic responsiveness, where there is no osmotic change during warming due to a slow cooling rate (cells remain dehydrated); (iii) altered osmotic behaviour, where cells membranes turn "leaky", resulting in the release of water and solutes into the surroundings; and (iv) intracellular ice formation, where rapid cooling causes membrane disruption due to the formation of ice crystals (Steponkus, 1984).




Fig. 3. Typical cell membrane structure consisting of a phospholipid bilayer with embedded sterols. Phospholipid chains are shown in grey, choline groups in blue and phosphate groups in red, while sterol molecules are shown in yellow.

The cell membrane is a bilayer consisting of different lipids and associated proteins (Fig. 3), where the lipids define the cell membrane structure and fluidity, and have a function in

signal transduction (Foubert et al., 2007; Furt et al., 2011). The three main classes of lipids found in cell membranes are glycerolipids (mostly phospholipids), sterols and sphingolipids (Furt et al., 2011).

4.1 Phospholipids

Phospholipids are amphiphilic molecules that form the core structure of the cell membrane lipid bilayer. They consist of a polar head (containing a phosphate group and simple organic molecule) and a (mostly) non-polar fatty acid tail. The most common phospholipid components of membranes include phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylglycerol (PG), phosphatidylinositol (PI), phosphatidylserine (PS) and phosphatidic acid (PA). Phospholipids essentially maintain the structure, fluidity and permeability of the membrane, determined to some extent by the type of phospholipid present and the level of unsaturation in their fatty acid tails (Emmelot & Vanhoeven, 1975; van Meer et al., 2008).

4.2 Sterols

Sterols are steroid alcohols that are integral components of living cells. They are formed as intermediate molecules during the metabolic production of hormones and are an integral component of cell membranes (Hartmann, 1998; Hodzic et al., 2008). Sterols are non-polar molecules with a polar hydroxyl (-OH) side chain that allows them to interact with the polar and non-polar groups of the phospholipid bilayer. They restrict the motion of the fatty acid chains in phospholipids, thus controlling the fluidity of the cell membrane (Hartmann, 1998). The major plant sterols are cholesterol, campesterol, sitosterol and stigmasterol (Grunwald, 1975). Sitosterol is usually found in higher concentration in apex tissue than stigmasterol (Grunwald & Saunders, 1978). The composition and concentration of these sterols within the membrane modify its permeability and fluidity (Grunwald, 1975; Grunwald & Saunders, 1978; Nes, 1974). Cholesterol has the greatest stabilising effect on membranes due to its small side chain (Grunwald & Saunders, 1978).

The ability of sterols to alter stability and permeability of membranes can have a large effect on post-cryopreservation survival of plant tissue. The ratio of stigmasterol to sitosterol has been found to increase after sucrose preculture and was unfavourable to obtaining high percentages of shoot regeneration after cryopreservation of banana meristems (Zhu et al., 2006). Marsan et al. (1998) investigated the interactions of sitosterol and stigmasterol with phosphatidylcholine molecules in soybean and discovered, using neutron scattering experiments, that sitosterol has a greater effect than stigmasterol on the ordering of the fatty acyl chains of PC and increasing the hydrophobic thickness of PC bilayers. Cold acclimation of winter rye (*Secale cereale*) seedlings showed an increase in free sterol content, with β -sitosterol having the largest increase (Lynch & Steponkus, 1987). Uemura and Steponkus (1994) also found an increase of β -sitosterol with cold acclimation in winter rye; however, these results were not mimicked in spring oats (*Avena sativa* L. cv Ogle), where there was no significant change in β -sitosterol, but the stigmasterol proportion increased, whilst cholesterol decreased.

4.3 Soluble sugars

Intracellular soluble sugars and sugar alcohols, such as the ones used in preculture media, reduce damage sustained by cell membranes when the cells undergo desiccation and can

improve membrane stability (Crowe et al., 1988; Steponkus, 1984). Smaller sugar molecules help membranes to osmotically retain water and may enter the interlamellar space to maintain a degree of hydration and increase the separation between membranes, thus reducing compressive stresses and, consequently, reducing the chance of a fluid-gel phase transition (Wolfe & Bryant, 1999). Furthermore, the polar hydroxyl (-OH) groups on sugars and sugar alcohols have been thought to interact with the polar membrane phospholipid headgroups and stabilise the membranes by maintaining the separation of the phospholipid molecules (Steponkus, 1984; Turner et al., 2001; Wolfe & Bryant, 1999). Turner et al. (2001) tested several sugars and sugar polyalcohols and determined that the small size of molecules such as glycerol and erythritol allowed them to 'pack' around membranes and have better bonding abilities. Additionally, the stereochemical arrangement of the -OH groups, particularly their orientation along one side of the molecules, imparted more stable hydrogen bonds with the membrane phospholipids and, consequently, resulted in more stable membranes (Turner et al., 2001). Nonetheless, recent biophysical studies have established that specific interactions of sugar molecules to phospholipid headgroups are not necessary to explain the stabilising effect of sugars on membrane gel to fluid transition temperatures (Lenne et al. 2006, 2007, 2009) and that sugars do not in fact insert between phospholipid headgroups but instead a solvation layer of water molecules separates them from the phospholipid headgroups (Kent et al., 2010).

4.4 Preconditioning and cold acclimation

Cold acclimation is the process in which plants being exposed to low non-freezing temperatures increase their freezing tolerance (Thomashow, 1999; Sakai & Engelmann, 2007). Preconditioning of raspberry and blueberry plants at 22/-1°C alternating temperatures for four weeks was essential for optimal post-cryopreservation shoot-tip re-growth using encapsulation-dehydration and vitrification protocols (Gupta & Reed, 2006). Similarly, better recovery rates were seen in mint (Senula et al., 2007) and yams (Leunufna & Keller, 2005) when they were cold acclimated for several weeks before cryostorage.

Cold acclimation is thought to activate genes that improve plant survival at low temperatures by improving stability in membranes (Guy et al., 1985; Thomashow, 1999). Cell membrane phospholipids and sterols have been observed to increase in proportion upon cold acclimation in winter rye, with a particular increase in di-unsaturated PC and β -sitosterol (Uemura & Steponkus, 1994). Changes in phospholipid and sterol composition were found in *Arabidopsis thaliana* after cold acclimation at 2°C for one week, which increased its freezing tolerance from -2°C down to -10°C (Uemura et al., 1995). Cell membrane phospholipid changes were also observed in oat leaves (Uemura & Steponkus, 1994). These changes may be related to improved membrane stability in these plants.

Preconditioning of plants has also been shown to increase antioxidant levels in plants prior to cryopreservation (Baek & Skinner, 2003; Dai et al., 2009; Harding et al., 2009; Zhao & Blumwald, 1998). Baek and Skinner (2003) analysed the expression of antioxidant genes in wheat species after cold acclimation and found increased expression of antioxidant enzymes, such as SOD and catalase. Increased levels of antioxidants may allow plants to better tolerate oxidative stress. Lynch and Steponkus (1987) observed an increase in the di-unsaturated levels of PC and PE in winter rye seedlings. Sucrose pre-treatment of banana

meristems prior to cryopreservation increased survival after warming and was related in most cases to a decrease of the double bond index in phospholipids, free fatty acids, glycolipids and sphingolipids (Zhu et al., 2006).

5. Genetic and epigenetic stability

The aim of successful cryopreservation is to maintain genetically stable plant material. While cryopreservation is now recognised as the method of choice for the long term preservation of plant material, the usefulness of cryostorage only applies if it does not lead to genetic changes in the plant species of interest (Zarghami et al., 2008). It is thus recommended to avoid the use of tissue in a non-organised dedifferentiated state, such as callus, and to use organised tissue like shoot tips instead to reduce the likelihood of non-desirable genetic mutations (Benson et al., 1996; Harding, 2004) as well as due to their higher regrowth rates (Bunn et al., 2007). Cryopreservation can cause injury at cellular level, but it is not clear if this injury can change the genetic composition of plants. The genetic stability of cryopreserved plants has nonetheless been confirmed for most of the analysed samples at morphological, histological and molecular level (Harding, 2004). Where differences between control and cryopreserved genotypes were found, it was suggested that the genetic changes might not be associated with the cryopreservation process itself, but rather that they are caused by the overall tissue culture, cryoprotection and regeneration process (Harding, 2004).

Comparisons of morphological development and analyses of the characteristics of control and cryopreserved plants have shown no differences in many species, such as *Prunus*, sugarcane, onion, kiwi, *Eucalyptus*, coffee, *Dendrobium* and *Cosmos* (Harding, 2004), as well as in the hybrid aspen (*Populus tremula*), an economically important woody plant and widely used model forest plant (Jokipii et al., 2004). Alterations in phenotype related to flower colouring have been observed in *Chrysanthemum*, which might be related to the chimeric structure of the plant (Harding, 2004). Morphological and phenotypic studies in potato, where shoot tips were used for cryopreservation, showed stable genetic integrity (Kaczmarczyk et al., 2010). Biometric studies examining morphological characters, agronomic traits or vegetal development descriptors in *Dioscorea floribunda*, sugarcane and banana revealed no significant differences between cryopreserved-derived and control plants (Harding, 2004). No abnormalities in chromosome number or cell structure were observed in cryopreserved *Vanda pumila* (*Orchidaceae*), with the regrown shoot primordia being able to induce new meristematic tissues like those of the non-cryopreserved controls (Na & Kondo, 1996). Long term storage of cryopreserved plants, such as strawberry, pea, *Rubus* and potato, did not result in any overall changes in regeneration capability and phenotype when regenerated explants were compared at the time point of storage as well as 10, 12 and 28 years later (Castillo et al., 2010; Caswell & Kartha, 2009; Keller et al., 2006; Mix-Wagner et al., 2003).

Histological studies and cytological analysis using flow cytometry have confirmed the genetic stability of plant species such as pea, oil palm, silver birch, *Rubus*, *Solanum tuberosum* and rice (Harding, 2004). Biochemical analyses have compared products of gene expression such as the formation and concentration of secondary metabolites. Examples of compounds compared in cryopreserved and control plants have been diosgenin in *Dioscorea floribunda*, chlorophyll and pyrethrin in *Chrysanthemum*, hypericin production in *Hypericum perforatum*

L. (Skyba et al., 2010; Urbanova et al., 2006), betalain pigments in *Beta vulgaris* and nicotin alkaloids in *Nicotiana rustica*, which were all unchanged in cryopreserved plants and thus confirmed the integrity of metabolic functions after cryostorage (Harding, 2004). Similar stability was observed upon comparison of proteins and enzymes (Marin et al., 1993; Paulet et al., 1993; Wu et al., 2001).

A variety of different techniques and markers have been applied to compare genomic DNA patterns, such as restriction fragment length polymorphism (RFLP), randomly amplified polymorphic DNA (RAPDs) fragments, simple sequence repeat (SSR) analysis and amplified fragment length polymorphism (AFLP). Most studies have confirmed the presence of genetic stability (Castillo et al., 2010; Helliott et al., 2002) and where changes in the genome have been found, such as in sugarcane and potato with RFLP markers, the changes could not be related to the process of cryopreservation itself (Castillo et al., 2010; Harding, 2004).

In contrast to genetic variations manifested by DNA nucleotide sequence alterations, epigenetic changes do not change the original DNA sequence (Boyko & Kovalchuk, 2008) but can result in heritable changes of gene expression. Typical features of epigenetic characteristics are DNA methylation, histone modification and changes in chromatin structure (Boyko & Kovalchuk, 2008). Epigenetic gene regulatory mechanisms have a function in plant development and might be influenced or changed by environmental conditions and osmotic stress during tissue culture and cryopreservation. Some recent studies have analysed epigenetic characteristics like DNA methylation in tissue culture and cryopreserved plants. Modifications in DNA methylation have been found in almond (Channuntapipat et al., 2003), papaya (Kaity et al., 2008), chrysanthemum (Martín & González-Benito, 2006), *Ribes* (Johnston et al., 2009), strawberry (Hao et al., 2002a), citrus (Hao et al., 2002b) and potato (Kaczmarczyk et al., 2010). Changes in methylation might be caused by stressful *in vitro* conditions, osmotic dehydration and the application of cryoprotective agents (Harding, 2004).

6. Conclusion

Many plant species have been successfully cryopreserved through the development of various cryopreservation methods. As a standard protocol, vitrification and droplet vitrification are widely applied. Shoot tips are the preferred material for cryostorage as they contain the meristem and an organised structure, with direct shoot development avoiding unstructured phases, which could lead to mutations. Preconditioning of plants (cold acclimation or sugar preculture) can have positive effects on survival and regeneration after cryopreservation, which could be due to increased membrane stability. Cryopreserved plants have been found to be genetically stable in most cases, but epigenetic changes have been detected, although most of the molecular analyses have only compared fractions of the genome.

Success in cryopreservation cannot be guaranteed for all plants, as some species are recalcitrant to tissue culture or the cryopreservation process. Fundamental studies looking at membrane composition, membrane damage and repair are likely to help to elucidate why some species are cryosensitive and how cryopreservation protocols can be improved for those species.

7. Acknowledgment

The authors thank ALCOA of Australia Ltd and BHP Billiton Worsley Alumina Pty Ltd for their generous support. AK is supported through the Australian Research Council Linkage Project Grant LP0884027. BF is supported through an Australian Postgraduate Award and a Curtin Research Scholarship. Dr. Zak Hughes is gratefully acknowledged for producing Fig. 3.

8. References

- Anchordoguy, T.J., Rudolph, A.S., Carpenter, J.F. & Crowe, J.H. (1987). Modes of interaction of cryoprotectants with membrane phospholipids during freezing. *Cryobiology* 24:pp. 324-331.
- Arakawa, T., Carpenter, J.F., Kita, Y.A. & Crowe, J.H. (1990). The basis for toxicity of certain cryoprotectants - a hypothesis. *Cryobiology* 27:pp. 401-415.
- Baek, K-H. & Skinner, D.Z. (2003). Alteration of antioxidant enzyme gene expression during cold acclimation of near isogenic wheat lines. *Plant Science* 165:pp. 1221-1227.
- Benson, E.E. (1990). *Free Radical Damage in Stored Plant Germplasm*, International Board for Plant Genetic Resources, ISBN 92-9043-196-2, Rome, Italy.
- Benson, E.E. (2008). Cryopreservation of phytodiversity: A critical appraisal of theory & practice. *Critical Reviews in Plant Sciences* 27:pp. 141-219.
- Benson, E.E. & Bremner, D. (2004). Oxidative stress in the frozen plant: a free radical point of view. In: *Life in the Frozen State*, Fuller, B.J., Lane, N. & Benson, E.E. (Eds.), pp. 206-241, CRC Press, ISBN 0-415-24700-4, Boca Raton, Florida.
- Benson, E.E., Lynch, P.T. & Jones, J. (1992). The detection of lipid peroxidation products in cryoprotected and frozen rice cells: consequences for post-thaw survival. *Plant Science* 85:pp. 107-114.
- Benson, E.E., Lynch, P.T. & Jones, J. (1995). The use of the iron chelating agent desferrioxamine in rice cell cryopreservation: a novel approach for improving recovery. *Plant Science* 110:pp. 249-258.
- Benson, E.E., Reed, B.M., Brennan, R.M., Clacher, K.A. & Ross, D.A. (1996). Use of thermal analysis in the evaluation of cryopreservation protocols for *Ribes nigrum* L. germplasm. *CryoLetters* 17:pp. 347-362.
- Benson, E.E. & Whithers, L.A. (1987). Gas-chromatographic analysis of volatile hydrocarbon production by cryopreserved plant-tissue cultures - a nondestructive method for assessing stability. *CryoLetters* 8:pp. 35-46.
- Boyko, A. & Kovalchuk, I. (2008). Epigenetic control of plant stress response. *Environmental and Molecular Mutagenesis* 49:pp. 61-72.
- Bunn, E., Turner, S., Panaia, M. & Dixon, K.W. (2007). The contribution of in vitro technology and cryogenic storage to conservation of indigenous plants, *Australian Journal of Botany* 55:pp. 345-355.
- Castillo, N.R.F., Bassil, N.V., Wada, S. & Reed, B.M. (2010). Genetic stability of cryopreserved shoot tips of *Rubus* germplasm. *In Vitro Cellular & Developmental Biology-Plant* 46:pp. 246-256.
- Caswell, K.L. & Kartha, K.K. (2009). Recovery of plants from pea and strawberry meristems cryopreserved for 28 years. *CryoLetters* 30:pp. 41-46.

- Channuntapipat, C., Sedgley, M. & Collins, G. (2003). Changes in methylation and structure of DNA from almond tissues during in vitro culture and cryopreservation. *J. Amer. Soc. Hort. Sci.* 128:pp. 890-897.
- Chen, X.-L., Li, J.-H., Xin, X., Zhang, Z.-E., Xin, P.-P. & Lu, X.-X. (2011). Cryopreservation of in vitro-grown apical meristems of *Lilium* by droplet-vitrification. *South African Journal of Botany* 77:pp. 397-403
- Crowe, J.H., Crowe, L.M., Carpenter, J.F., Rudolph, A.S., Wistrom, C.A., Spargo, B.J. & Anchordoguy, T.J. (1988). Interactions of sugars with membranes. *Biochimica et Biophysica Acta* 947:pp. 367-84.
- Dai, F., Huang, Y., Zhou, M. & Zhang G. (2009). The influence of cold acclimation on antioxidant enzymes and antioxidants in sensitive and tolerant barley cultivars. *Biologia Plantarum* 153:pp. 257-262.
- Day, J.G., Harding, K.C., Nadarajan, J. & Benson, E.E. (2008). Cryopreservation, conservation of bioresources at ultra low temperatures. In: *Molecular Biomethods Handbook*. Walker J.M & Rapley R. (Eds.), pp. 917-947, Humana Press, ISBN 978-1-60327-374-9, Totowa, NJ.
- Day, J.G. & Stacey, G.N. (2007). *Cryopreservation and Freeze-Drying Protocols*. Humana Press Inc., ISBN 978-1-58829-377-0, Totowa, NJ.
- Decruse, S.W., Seeni, S. & Pushpangadan, P. (1999). Cryopreservation of alginate coated shoot tips of in vitro grown *Holostemma annulare* (Roxb.) K. Schum, an endangered medicinal plant: Influence of preculture and DMSO treatment on survival and regeneration. *CryoLetters* 20:pp. 243-250.
- Dowdle, J., Ishikawa, T., Gatzek, S., Rolinski, S. & Smirnov, N. (2007). Two genes in *Arabidopsis thaliana* encoding GDP I galactose phosphorylase are required for ascorbate biosynthesis and seedling viability. *The Plant Journal* 52:pp. 673-689.
- Dowgert, M.F. & Steponkus, P.L. (1984). Behavior of the plasma membrane of isolated protoplasts during a freeze-thaw cycle. *Plant Physiol* 75:pp. 1139-1151.
- Emmelot, P. & Vanhoeven, R.P. (1975). Phospholipid unsaturation and plasma membrane organization. *Chemistry and Physics of Lipids* 14:pp. 236-246.
- Engelmann, F. (2004). Plant cryopreservation: Progress and prospects. *In Vitro Cellular & Developmental Biology-Plant* 40:pp. 427-433.
- Fabre, J. & Dereuddre, J. (1990). Encapsulation-dehydration: A new approach to cryopreservation of *Solanum* shoot-tips. *CryoLetters* 11:pp. 423-426.
- Fang, J.Y., Wetten, A. & Johnston, J. (2008). Headspace volatile markers for sensitivity of cocoa (*Theobroma cacao* L.) somatic embryos to cryopreservation. *Plant Cell Reports* 27: pp. 453-461.
- Fleck, R.A., Benson, E.E., Bremner, D.H. & Day, J.G. (2000). Studies of free radical-mediated cryoinjury in the unicellular green alga *Euglena gracilis* using a non-destructive hydroxyl radical assay: a novel approach for developing protistan cryopreservation strategies. *Free Radic Res* 32:pp. 157-70.
- Foubert, I., Dewettinck, K., Van de Walle, D., Dijkstra, A.J. & Quinn, P.J. (2007). Physical properties: structural and physical characteristics, In: *The Lipid Handbook with CD-ROM*, Gunstone, F.D., Harwood J.L. & Dijkstra A.J. (Eds.), pp. 471-534, CRC Press, Taylor & Francis Group, ISBN 0-8493-9688-3, LLC, Boca Raton, FL.

- Foyer, C.H. & Noctor, G. (2009). Redox regulation in photosynthetic organisms: signaling, acclimation, and practical implications. *Antioxidants & Redox Signaling* 11:pp. 861-905.
- Furt, F., Simon-Plas, F. & Sebastien, M. (2011). Lipids of the plant plasma membrane, In: *The Plant Plasma Membrane*. Murphy, A.S., Peer, W. & Schulz, B. (Eds.), pp. 3-30, Springer-Verlag, ISBN 978-3-642-13430-2, Berlin Heidelberg.
- Gamez-Pastrana, R., Martinez-Ocampo, Y., Beristain, C.I. & Gonzalez-Arnao, M.T. (2004). An improved cryopreservation protocol for pineapple apices using encapsulation-vitrification. *CryoLetters* 25:pp. 405-414.
- Gonzalez-Arnao, M.T., Panta, A., Roca, W.M., Escobar, R.H. & Engelmann, F. (2008). Development and large scale application of cryopreservation techniques for shoot and somatic embryo cultures of tropical crops. *Plant Cell Tissue and Organ Culture* 92:pp. 1-13.
- Gordon-Kamm, W.J. & Steponkus, P.L. (1984). Lamellar-to-hexagonal II phase transitions in the plasma membrane of isolated protoplasts after freeze-induced dehydration. *Proc Natl Acad Sci USA* 81:pp. 6373-6377.
- Grunwald, C. (1975). Plant sterols. *Annual Review of Plant Physiology and Plant Molecular Biology* 26:pp. 209-236.
- Grunwald, C. & Saunders, P.F. (1978). Function of sterols [and discussion]. *Philosophical Transactions of the Royal Society of London B Biological Sciences* 284:pp. 541-558.
- Gupta, S. & Reed, B.M. (2006). Cryopreservation of shoot tips of blackberry and raspberry by encapsulation-dehydration and vitrification. *CryoLetters* 27:pp. 29-42.
- Guy, C.L., Niemi, K.J. & Brambl, R. (1985). Altered gene expression during cold acclimation of spinach. *Proc Natl Acad Sci USA* 82: pp. 3673-3677.
- Halliwell, B. & Gutteridge, J.M.C. (2007). *Free Radicals in Biology and Medicine* (Fourth edition), Oxford University Press Inc., ISBN 978-0-19-856868-1, Oxford.
- Halmagyi, A., Fischer-Klüver, G., Mix-Wagner, G. & Schumacher, H.M. (2004). Cryopreservation of *Chrysanthemum morifolium* (*Dendranthema grandiflora* Ramat.) using different approaches. *Plant Cell Rep* 22:pp. 371-375.
- Hamilton, K.N., Turner, S.R. & Ashmore, S.E. (2009). Cryopreservation, In: *Plant Germplasm Conservation in Australia: Strategies and Guidelines for Developing, Managing and Utilising Ex Situ Collections*, Offord, C.A & Meagher, P.F. (Eds.), pp. 129-128, Australian Network for Plant Conservation Inc., ISBN 978-0-9752191-1-9, Canberra.
- Hao, Y.J., You, C.X. & Deng, X.X. (2002a). Analysis of ploidy and the patterns of amplified fragment length polymorphism and methylation sensitive amplified polymorphism in strawberry plants recovered from cryopreservation. *CryoLetters* 23:pp. 37-46.
- Hao, Y.J., You, C.X. & Deng, X.X. (2002b). Effects of cryopreservation on developmental competency, cytological and molecular stability of citrus callus. *CryoLetters* 23:pp. 27-35.
- Harding, K. (2004). Genetic integrity of cryopreserved plant cells: a review. *CryoLetters* 25:pp. 3-22.
- Harding, K.N., Johnston, J.W. & Benson, E.E. (2009). Exploring the physiological basis of cryopreservation success and failure in clonally propagated in vitro crop plant germplasm. *Agricultural and Food Science* 18:pp. 103-116.

- Hartmann, M.A. (1998). Plant sterols and the membrane environment. *Trends in Plant Science* 3:pp. 170-175.
- Helliot, B., Madur, D., Dirlewanger, E. & De Boucaud, M.T. (2002). Evaluation of genetic stability in cryopreserved *Prunus*. *In Vitro Cellular & Developmental Biology-Plant* 38:pp. 493-500.
- Hipkiss, A.R., Preston, J.E., Himsworth, D., Worthington, V.C. & Abbot, N.J. (1997). Protective effects of carnosine against malondialdehyde-induced toxicity towards cultured rat brain endothelial cells. *Neuroscience Letters* 238:pp. 135-138.
- Hirai, D. & Sakai, A. (1999). Cryopreservation of in vitro-grown meristems of potato (*Solanum tuberosum* L.) by encapsulation-vitrification. *Potato Research* 42:pp. 153-160.
- Hirai, D., Shirai, K., Shirai, S. & Sakai, A. (1998). Cryopreservation of in vitro grown meristems of strawberry (*Fragaria x ananassa* Duch.) by encapsulation-vitrification. *Euphytica* 101:pp. 109-115.
- Hodzic, A., Rappolt, M., Amenitsch, H., Laggner, P. & Pabst, G. (2008). Differential modulation of membrane structure and fluctuations by plant sterols and cholesterol. *Biophysical Journal* 94:pp. 3935-3944.
- Johnston, J.W., Benson, E.E. & Harding, K. (2009). Cryopreservation induces temporal DNA methylation epigenetic changes and differential transcriptional activity in *Ribes* germplasm. *Plant Physiology and Biochemistry* 47:pp. 123-131.
- Jokipii, S., Rynänen, L., Kallio, P.T., Aronen, T. & Häggmana, H. (2004). A cryopreservation method maintaining the genetic fidelity of a model forest tree, *Plant Science*. 166:pp. 799-806.
- Kaczmarczyk, A., Houben, A., Keller, E.R.J. & Mette, M.F. (2010). Influence of cryopreservation on the cytosine methylation state of potato genomic DNA. *CryoLetters* 31:pp. 380-391.
- Kaczmarczyk, A., Rokka, V.-M. & Keller, E.R.J. (2011a). Potato shoot tip cryopreservation, a review. *Potato Research* 54:pp. 45-79.
- Kaczmarczyk, A., Turner, S.R., Bunn, E., Mancera, R.L. & Dixon, K.W. (2011b). Cryopreservation of threatened native Australian species - what have we learned and where to from here? *In Vitro Cellular & Developmental Biology-Plant* 47:pp. 17-25.
- Kaity, A., Ashmore, S.E., Drew, R.A. & Dulloo, M.E. (2008). Assessment of genetic and epigenetic changes following cryopreservation in papaya. *Plant Cell Reports* 27:pp. 1529-1539.
- Keller, E.R.J., Senula, A., Leunufna, S. & Grube, M. (2006). Slow growth storage and cryopreservation - tools to facilitate germplasm maintenance of vegetatively propagated crops in living plant collections. *International Journal of Refrigeration-Revue Internationale Du Froid* 29:pp. 411-417.
- Kent, B., Garvey, C.J., Lenne, T., Porcar, L., Garamus, V.M. & Bryant, G. (2010). Measurement of glucose exclusion from the fully hydrated DOPE inverse hexagonal phase. *Soft Matter* 6:pp. 1197-1202.
- Kim, H.H., Lee, Y.G., Shin, D.J., Ko, H.C., Gwag, J.G., Cho, E.G. & Engelmann, F. (2009). Development of alternative plant vitrification solutions in droplet-vitrification procedures. *CryoLetters* 30:pp. 320-34.
- Kranner, I., Birtic, S., Anderson, K.M. & Pritchard, H.W. (2006). Glutathione half-cell reduction potential: A universal stress marker and modulator of programmed cell death? *Free Radical Biology and Medicine* 40:pp. 2155-2165.

- Kreck, C.A., Mandumpal, J.B. & Mancera, R.L. (2011). Prediction of the glass transition in aqueous solutions of simple amides by molecular dynamics simulations. *Chemical Physics Letters* 501:pp. 273-277.
- Lenne, T., Bryant, G., Garvey, C.J., Kelderling, U. & Koster, K.L. (2006). Location of sugars in multilamellar membranes at low hydration. *Physica B-Condensed Matter* 385-86:pp. 862-864.
- Lenne, T., Bryant, G., Holcomb, R. & Koster, K.L. (2007). How much solute is needed to inhibit the fluid to gel membrane phase transition at low hydration? *Biochimica et Biophysica Acta-Biomembranes* 1768:pp. 1019-1022.
- Lenne, T., Garvey, C.J., Koster, K.L. & Bryant, G. (2009). Effects of sugars on lipid bilayers during dehydration - SAXS/WAXS measurements and quantitative model. *Journal of Physical Chemistry B*, 113:pp. 2486-2491.
- Leunufna, S. & Keller, E.R.J. (2005). Cryopreservation of yams using vitrification modified by including droplet method: effects of cold acclimation and sucrose. *CryoLetters* 26:pp. 93-102.
- Lynch, D.V. & Steponkus, P.L. (1987). Plasma membrane lipid alterations associated with cold acclimation of winter rye seedlings (*Secale cereale* L. cv Puma). *Plant Physiol* 83:pp. 761-767.
- Lynch, P.T., Siddika, A., Johnston, J.W., Trigwell, S.M., Mehra, A., Benelli, C., Lambardi, M. & Benson, E.E. (2011). Effects of osmotic pretreatments on oxidative stress, antioxidant profiles and cryopreservation of olive somatic embryos. *Plant Science* 181:pp. 47-56.
- Mallon, R., Bunn, E., Turner, S.R. & Gonzalez, M.L. (2008). Cryopreservation of *Centaurea ultriae* (Compositae) a critically endangered species from Galicia (Spain). *CryoLetters* 29:pp. 363-370.
- Mandal, B.B. & Dixit-Sharma, S. (2007). Cryopreservation of in vitro shoot tips of *Dioscorea deltoidea* Wall., an endangered medicinal plant: effect of cryogenic procedure and storage duration. *CryoLetters* 28:pp. 461-470.
- Mandumpal, J.B., Kreck, C.A. & Mancera, R.L. (2011). A molecular mechanism of solvent cryoprotection in aqueous DMSO solutions. *Physical Chemistry Chemical Physics* 13:pp. 3839-3842.
- Marin, M.L., Gogorcena, Y., Ortiz, J. & Duranvila, N. (1993). Recovery of whole plants of sweet orange from somatic embryos subjected to freezing-thawing treatments. *Plant Cell Tissue and Organ Culture* 34:pp. 27-33.
- Marnett, L.J. (1999). Chemistry and biology of DNA damage by malondialdehyde. *IARC scientific publications* 150:pp. 1-17.
- Marsan, M.P., Bellet-Amalric, E., Muller, I., Zaccari, G. & Milon, A. (1998). Plant sterols: a neutron diffraction study of sitosterol and stigmasterol in soybean phosphatidylcholine membranes. *Biophysical Chemistry* 75:pp. 45-55.
- Martin, M.C. & González-Benito, M. E. (2006) Sequence comparison in somaclonal variant of cryopreserved *Dendranthema grandiflora* shoot apices. *Cryobiology* 53:pp. 424.
- Meryman, H.T. (1974). Freezing injury and its prevention in living cells *Annual Review of Biophysics and Bioengineering* 3:pp. 341-363.
- Mix-Wagner, G., Schumacher, H.M. & Cross, R.J. (2003). Recovery of potato apices after several years of storage in liquid nitrogen. *CryoLetters* 24:pp. 33-41.

- Møller, I.M., Jensen, P.E. & Hansson, A. (2007). Oxidative modifications to cellular components in plants. *Annu. Rev. Plant Biol.* 58:pp. 459-481.
- Mycock, D.J., Wesley-Smith, J. & Berjak, P. (1995). Cryopreservation of somatic embryos of four species with and without cryoprotectant pre-treatment. *Annals of Botany* 75: pp. 331-336.
- Na, H.-Y. & Kondo, K. (1996). Cryopreservation of tissue-cultured shoot primordia from shoot apices of cultured protocorms in *Vanda pumila* following ABA preculture and desiccation. *Plant Science* 118:pp. 195-201.
- Nes, W.R. (1974). Role of sterols in membranes. *Lipids* 9:pp. 596-612.
- Nishizawa, S., Sakai, A., Amano, Y. & Matsuzawa, T. (1993). Cryopreservation of *Asparagus officinalis* L. embryogenic suspension cells and subsequent plant regeneration by vitrification. *Plant Science* 91:pp. 67-73.
- Obert, B., Benson, E.E., Millam, S., Pretová, A. & Bremner, D.H. (2005). Moderation of morphogenetic and oxidative stress responses in flax in vitro cultures by hydroxynonenal and desferrioxamine. *Journal of Plant Physiology* 162:pp. 537-547.
- Packer, J.E., Slater, T. & Willson, R. (1979). Direct observation of a free radical interaction between vitamin E and vitamin C. *Nature* 278:pp. 737-738.
- Panis, B., Strosse, H., Van den Hende, S. & Swennen, R. (2002). Sucrose preculture to simplify cryopreservation of banana meristem cultures. *CryoLetters* 23:pp. 375-384.
- Panis, B. & Lambardi, M. (2005). Status of cryopreservation technologies in plants (crops and forest trees). Proceedings of the International Workshop on 'The Role of Biotechnology', pp. 43-54, Villa Gualino, Turin, Italy, 5.-7. March 2005.
- Panis, B., Piette, B. & Swennen, R. (2005). Droplet vitrification of apical meristems: a cryopreservation protocol applicable to all Musaceae. *Plant Science* 168:pp. 45-55.
- Partelli, F.L., Vieira, H.D., Pais, I.P., Quartin, V.L., Campos, P.S., Fortunato, A., Eichler, P., Viana, A.P., Ribeiro, A. & Ramalho, J.C. (2009). Chloroplast membrane lipids from *Coffea* sp. under low positive temperatures. Proceedings of 22nd International Conference on Coffee Science, pp. 891-898, ASIC 2008, ISBN 2900212219, Campinas, SP, Brazil, September, 2008.
- Paulet, F., Engelmann, F. & Glaszmann, J.C. (1993). Cryopreservation of apices of in vitro plantlets of sugarcane (*Saccharum* sp hybrids) using encapsulation dehydration. *Plant Cell Reports* 12:pp. 525-529.
- Paunescu, A. (2009). Biotechnology for endangered plant conservation: a critical overview *Romanian Biotechnological Letters* 14:pp. 4095-4103.
- Reed, B.M. (2008). *Plant Cryopreservation: A Practical Guide*, Springer, ISBN 978-0-387-72275-7, New York.
- Reinhold, P.J., Versteeg, I., Kars, I., van Iren, F. & Kijne, J.W. (2000). Physiological and molecular changes in tobacco suspension cells during development of tolerance to cryopreservation by vitrification, In: *Cryopreservation of Tropical Plant Germplasm. Current Research Progress and Application*, Engelmann, F. & Takagi, H. (Eds.), pp. 57-66, International Plant Genetic Resources Institute, ISBN 92-9043-428-7, Rome.
- Roach, T., Ivanova, M., Beckett, R.P., Minibayeva, F.V., Green, I., Pritchard, H.W. & Kranner, I. (2008). An oxidative burst of superoxide in embryonic axes of recalcitrant sweet chestnut seeds as induced by excision and desiccation. *Physiologia Plantarum* 133:pp. 131-139.

- Sakai, A. (1965). Survival of plant tissue at super-low temperatures III. Relation between effective prefreezing temperatures and the degree of frost hardiness. *Plant Physiol.* 40:pp. 882-887.
- Sakai, A. & Engelmann, F. (2007). Vitrification, encapsulation-vitrification and droplet-vitrification: a review. *CryoLetters* 28:pp. 151-172.
- Sakai, A., Hirai, D. & Niino, T. (2008). Development of PVS-based vitrification and encapsulation-vitrification protocols, In: *Plant Cryopreservation: A Practical Guide*, Reed, B. (Ed.), pp. 33-57, Springer, ISBN 978-0-378-72275-7, New York.
- Sakai, A., Kobayashi, S. & Oiyama, I. (1990). Cryopreservation of nucellar cells of navel orange (*Citrus sinensis* Osb. var. *brasiliensis* Tanaka) by vitrification. *Plant Cell Reports* 9:pp. 30-33.
- Scottez, C., Chevreau, E., Godard, N., Arnaud, Y., Duron, M. & Dereuddre, J. (1992). Cryopreservation of cold acclimated shoot tips of pear in vitro cultures after encapsulation-dehydration. *Cryobiology* 29:pp. 691-700.
- Sen-Rong, H. & Ming-Hua, Y. (2009). High-efficiency vitrification protocols for cryopreservation of in vitro grown shoot tips of rare and endangered plant *Emmenopterys henryi* Oliv. *Plant Cell Tissue and Organ Culture* 99:pp. 217-226.
- Senula, A., Keller, E.R.J., Sanduijav, T. & Yohannes, T. (2007). Cryopreservation of cold-acclimated mint (*Mentha* spp.) shoot tips using a simple vitrification protocol. *CryoLetters* 28:pp. 1-12.
- Skyba, M., Urbanova, M., Kapchina-Toteva, V., Kosuth, J., Harding, K. & Cellarova, E. (2010). Physiological, biochemical and molecular characteristics of cryopreserved *Hypericum perforatum* L. shoot tips. *CryoLetters* 31:pp. 249-260.
- Steponkus, P.L. (1984). Role of the plasma membrane in freezing injury and cold acclimation. *Annual Review of Plant Physiology* 35:pp. 543-584.
- Streb, P., Shang, W., Feierabend, J. & Bigny, R. (1998). Divergent strategies of photoprotection in high-mountain plants. *Planta* 207:pp. 313-324.
- Takagi, H., Thinh, N.T., Islam, O.M., Senboku, T. & Sakai, A. (1997). Cryopreservation of in vitro-grown shoot tips of taro (*Colocasia esculenta* (L) Schott) by vitrification 1. Investigation of basic conditions of the vitrification procedure. *Plant Cell Reports* 16:pp. 594-599.
- Tanaka, D., Niino, T., Isuzugawa, K., Hikage, T. & Uemura, M. (2004). Cryopreservation of shoot apices of in-vitro grown gentian plants: Comparison of vitrification and encapsulation-vitrification protocols. *CryoLetters* 25:pp. 167-176.
- Thomashow, M.F. (1999). Plant cold acclimation: Freezing tolerance genes and regulatory mechanisms. *Annual Review of Plant Physiology and Plant Molecular Biology* 50:pp. 571-599.
- Touchell, D.H., Dixon, K.W. & Tan, B. (1992). Cryopreservation of shoot-tips of *Grevillea scapigera* (Proteaceae) - a rare and endangered plant from Western-Australia. *Australian Journal of Botany* 40:pp. 305-310.
- Touchell, D.H., Turner, S.R., Bunn, E. & Dixon, K.W. 2002. Cryostorage of somatic tissues of endangered Australian species, In: *Cryopreservation of Plant Germplasm*. Towill, L.E. & Bajaj, Y.P.S. (Eds.), pp. 357-372, Springer Verlag, ISBN 7055-8630, Berlin Heidelberg, Germany.

- Turner, S. 2001. *Cryopreservation of somatic germplasm of selected Australian monocotyledonous taxa (Haemodoraceae)*, PhD Thesis, Curtin University of Technology, Perth, Western Australia.
- Turner, S., Senaratna, T., Touchell, D., Bunn, E., Dixon, K. & Tan, B. (2001). Stereochemical arrangement of hydroxyl groups in sugar and polyalcohol molecules as an important factor in effective cryopreservation. *Plant Science* 160:pp. 489-497.
- Uchendu, E.E., Leonard, S.W., Traber, M.G. & Reed, B.M. (2010). Vitamins C and E improve regrowth and reduce lipid peroxidation of blackberry shoot tips following cryopreservation. *Plant Cell Reports* 29:pp. 25-35.
- Uemura, M. & Steponkus, P.L. (1994). A contrast of the plasma membrane lipid composition of oat and rye leaves in relation to freezing tolerance. *Plant Physiology* 104:pp. 479-496.
- Uemura, M., Joseph, R.A. & Steponkus, P.L. (1995). Cold-acclimation of *Arabidopsis thaliana* - effect on plasma membrane lipid composition and freeze-induced lesions. *Plant Physiology* 109:pp. 15-30.
- Urbanova, M., Kosuth, J. & Cellarova, E. (2006). Genetic and biochemical analysis of *Hypericum perforatum* L. plants regenerated after cryopreservation. *Plant Cell Reports* 25:pp. 140-147.
- van Meer, G., Voelker, D.R. & Feigenson, G.W. (2008). Membrane lipids: where they are and how they behave. *Nature Reviews Molecular Cell Biology* 9:pp. 112-124.
- Vidal, N., Sanchez, C., Jorquera, L., Ballester, A. & Vieitez, A.M. (2005). Cryopreservation of chestnut by vitrification of in vitro-grown shoot tips. *In Vitro Cellular & Developmental Biology-Plant* 41:pp. 63-68.
- Volk, G.M. & Walters, C. (2006). Plant vitrification solution 2 lowers water content and alters freezing behavior in shoot tips during cryoprotection. *Cryobiology* 52:pp. 48-61.
- Wang, Z.C. & Deng, X.X. (2004). Cryopreservation of shoot-tips of citrus using vitrification: effect of reduced form of glutathione. *CryoLetters* 25:pp. 43-50.
- Wise, R.R. (1995). Chilling-enhanced photooxidation: the production, action and study of reactive oxygen species produced during chilling in the light. *Photosynthesis Research* 45:pp. 79-97.
- Wise, R.R. & Naylor, A.W. (1987). Chilling-enhanced photooxidation: evidence for the role of singlet oxygen and superoxide in the breakdown of pigments and endogenous antioxidants. *Plant Physiology* 83:pp. 278.
- Wolfe, J. & Bryant, G. (1999). Freezing, drying, and/or vitrification of membrane-solute-water systems. *Cryobiology* 39:pp. 103-129.
- Wu, Y.J., Zhao, Y.H., Engelmann, R. & Zhou, M.D. (2001). Cryopreservation of kiwi shoot tips. *CryoLetters* 22:pp. 277-284.
- Yoon, J.-W., Kim, H.-H., Ko, H.-C., Hwang, H.-S., Hong, E.-S., Cho, E.-G. & Engelmann, F. (2006). Cryopreservation of cultivated and wild potato varieties by droplet vitrification: effect of subculture of mother-plants and of preculture of shoot tips. *CryoLetters* 27:pp. 211-222.
- Young, I. & McEneny, J. (2001). Lipoprotein oxidation and atherosclerosis. *Biochemical Society Transactions* 29:pp. 358-362.
- Zarghami, R., Pirseyedi, S.M., Hasrak, Sh. & Pakdaman, B.S. (2008). Evaluation of genetic stability in cryopreserved *Solanum tuberosum*. *African Journal of Biotechnology* 7:pp. 2798-2802.

- Zhao, S. & Blumwald, E. (1998). Changes in oxidation reduction state and antioxidant enzymes in the roots of jack pine seedlings during cold acclimation. *Physiologia Plantarum* 104: pp.134-142.
- Zhu, G.Y., Geuns, J.M.C., Dussert, S., Swennen, R. & Panis, B. (2006). Change in sugar, sterol and fatty acid composition in banana meristems caused by sucrose-induced acclimation and its effects on cryopreservation. *Physiologia Plantarum* 128:pp. 80-94.