

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368


Editors: W.T.A. Harrison, J. Simpson and
M. Weil

3-Pyridin-2-yl-1*H*-1,2,4-triazol-5-amine

Anton V. Dolzhenko, Geok Kheng Tan, Lip Lin Koh, Anna V. Dolzhenko
and Wai Keung Chui

Acta Cryst. (2009). **E65**, o125

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Pyridin-2-yl-1H-1,2,4-triazol-5-amine

Anton V. Dolzhenko,^{a*} Geok Kheng Tan,^b Lip Lin Koh,^b
Anna V. Dolzhenko^a and Wai Keung Chui^a^aDepartment of Pharmacy, Faculty of Science, National University of Singapore, 18 Science Drive 4, Singapore 117543, Singapore, and ^bDepartment of Chemistry, Faculty of Science, National University of Singapore, 3 Science Drive 3, Singapore 117543, Singapore

Correspondence e-mail: phada@nus.edu.sg


Received 13 November 2008; accepted 11 December 2008

Key indicators: single-crystal X-ray study; $T = 223$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.042; wR factor = 0.110; data-to-parameter ratio = 14.6.

In the title compound, $C_7H_7N_5$, the non-H atoms are almost coplanar (r.m.s. deviation = 0.050 Å), with the N atom of pyridine ring oriented to the N—N(H) side of the 1,2,4-triazole ring. The mean planes of the pyridine and 1,2,4-triazole rings form a dihedral angle of 5.58 (7)°. The N atom of the amino group adopts a pyramidal configuration. The molecules are linked into a two-dimensional network parallel to (10 $\bar{1}$) by N—H...N hydrogen bonds.

Related literature

For 1,2,4-triazol-5-amines as building blocks in the synthesis of fused heterocyclic systems, see: Dolzhenko *et al.* (2006, 2007*a,b*); Fischer, (2007). For a summary of structural data for 1,2,4-triazoles, see: Buzykin *et al.* (2006). For crystal structures of Cu^{II} complexes with 3-pyridin-2-yl-1,2,4-triazol-5-amine, see: Ferrer *et al.* (2004).



Experimental

Crystal data

$C_7H_7N_5$
 $M_r = 161.18$
Monoclinic, $P2_1/n$
 $a = 7.3863$ (6) Å

$b = 7.9096$ (6) Å
 $c = 13.2157$ (11) Å
 $\beta = 91.832$ (2)°
 $V = 771.70$ (11) Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹

$T = 223$ (2) K
 $0.36 \times 0.16 \times 0.12$ mm

Data collection

Bruker SMART APEX CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2001)
 $T_{min} = 0.967$, $T_{max} = 0.989$

5336 measured reflections
1772 independent reflections
1519 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.110$
 $S = 1.05$
1772 reflections
121 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{max} = 0.21$ e Å⁻³
 $\Delta\rho_{min} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N2-H2N\cdots N5^i$	0.90 (2)	2.01 (2)	2.9010 (16)	171 (1)
$N4-H4A\cdots N3^{ii}$	0.90 (2)	2.11 (2)	2.9971 (16)	172 (1)
$N4-H4B\cdots N1^i$	0.93 (2)	2.19 (2)	3.0264 (16)	151 (1)

Symmetry codes: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 1, -y, -z + 1$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINTE* (Bruker, 2001); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the National Medical Research Council, Singapore (grant Nos. NMRC/NIG/0019/2008 and NMRC/NIG/0020/2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2719).

References

- Bruker (2001). *SMART* and *SAINTE*. Bruker AXS GmbH, Karlsruhe, Germany.
- Buzykin, B. I., Mironova, E. V., Nabiullin, V. N., Gubaidullin, A. T. & Litvinov, I. A. (2006). *Russ. J. Gen. Chem.* **76**, 1471–1486.
- Dolzhenko, A. V., Dolzhenko, A. V. & Chui, W. K. (2006). *Heterocycles*, **68**, 1723–1759.
- Dolzhenko, A. V., Dolzhenko, A. V. & Chui, W. K. (2007*a*). *Heterocycles*, **71**, 429–436.
- Dolzhenko, A. V., Dolzhenko, A. V. & Chui, W. K. (2007*b*). *Tetrahedron*, **63**, 12888–12895.
- Ferrer, S., Ballesteros, R., Sambartolome, A., Gonzalez, M., Alzuet, G., Borrás, J. & Liu, M. (2004). *J. Inorg. Biochem.* **98**, 1436–1446.
- Fischer, G. (2007). *Adv. Heterocycl. Chem.* **95**, 143–219.
- Sheldrick, G. M. (2001). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o125 [doi:10.1107/S1600536808042177]

3-Pyridin-2-yl-1*H*-1,2,4-triazol-5-amine

A. V. Dolzhenko, G. K. Tan, L. L. Koh, A. V. Dolzhenko and W. K. Chui

Comment

1,2,4-Triazol-5-amines have been recognized as valuable synthons for the construction of fused heterocyclic systems, *e.g.* 1,2,4-triazolo[1,5-*a*]pyrimidines (Fischer, 2007) and 1,2,4-triazolo[1,5-*a*][1,3,5]triazines (Dolzhenko *et al.*, 2006). It also should be mentioned that 1,2,4-triazol-5-amines are widely used as ligands and crystallographic data on three different mono-nuclear complexes of 3-pyridin-2-yl-1,2,4-triazol-5-amine with Cu^{II} have been reported by Ferrer *et al.* (2004). However, no crystallographic study has been performed on the ligand.

In continuation of our investigations on using 1,2,4-triazol-5-amines in the synthesis of fused heterocyclic systems (Dolzhenko *et al.*, 2007*a,b*), we report herein the crystal structure of a synthetically important building block *viz.* 3-pyridin-2-yl-1,2,4-triazol-5-amine.

Due to annular tautomerism, 3-pyridin-2-yl-1,2,4-triazol-5-amine may theoretically exist in three tautomeric forms (**A**, **B** and **C**) and for each of them, rotameric structures **A'**, **B'** and **C'** are possible (Fig.1). As observed in reported Cu^{II} complexes (Ferrer *et al.*, 2004), 3-pyridin-2-yl-1,2,4-triazol-5-amine was the only tautomeric form found in the crystal (Fig. 2). However, the molecule exists in the crystal as rotamer **A** in contrast to rotamer **A'** found in Cu^{II} complexes.

Bond lengths and angles in the molecule of 3-pyridin-2-yl-1,2,4-triazol-5-amine are within normal ranges, and comparable with values summarized for 1,2,4-triazoles by Buzykin *et al.* (2006). 3-Pyridin-2-yl-1,2,4-triazol-5-amine has practically planar geometry with slight deviation of the pyridyl moiety, which makes a dihedral angle of 5.58 (7)° with mean plane of the 1,2,4-triazole ring. The nitrogen atom (N4) of the amino group adopts a pyramidal configuration with 0.26 (2) Å deviation of the nitrogen atom from the C2/H4A/H4B plane.

The molecules are linked into a two-dimensional network parallel to the (10 $\bar{1}$) by N—H \cdots N hydrogen bonds (Table 1 and Fig.3).

Experimental

3-Pyridin-2-yl-1,2,4-triazol-5-amine was prepared according to general method reported by Dolzhenko *et al.* (2007*a,b*). Single crystals suitable for crystallographic analysis were grown by recrystallization from ethanol.

Refinement

N-bound H-atoms were located in a difference map and refined freely [N—H = 0.90 (2)–0.92 (2) Å]. C-bound H atoms were positioned geometrically (C—H = 0.94 Å) and were constrained in a riding motion approximation with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Possible tautomers and rotamers of 3-pyridin-2-yl-1,2,4-triazol-5-amine.




Fig. 2. The molecular structure of the title compound, with the atomic numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 3. Molecular packing of the title compound, viewed along the *c* axis.

3-Pyridin-2-yl-1*H*-1,2,4-triazol-5-amine

Crystal data

$C_7H_7N_5$

$M_r = 161.18$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1/n$

$a = 7.3863$ (6) Å

$b = 7.9096$ (6) Å

$c = 13.2157$ (11) Å

$\beta = 91.832$ (2)°

$V = 771.70$ (11) Å³

$Z = 4$

$F_{000} = 336$

$D_x = 1.387$ Mg m⁻³

Melting point: 493 K

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 1841 reflections

$\theta = 3.0$ – 26.6 °

$\mu = 0.10$ mm⁻¹

$T = 223$ (2) K

Block, colourless

$0.36 \times 0.16 \times 0.12$ mm

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 223$ (2) K

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2001)

$T_{\min} = 0.967$, $T_{\max} = 0.989$

5336 measured reflections

1772 independent reflections

1519 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\text{max}} = 27.5$ °

$\theta_{\text{min}} = 3.0$ °

$h = -9 \rightarrow 9$

$k = -8 \rightarrow 10$

$l = -14 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.042$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.110$	$w = 1/[\sigma^2(F_o^2) + (0.0539P)^2 + 0.1487P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
1772 reflections	$(\Delta/\sigma)_{\max} = 0.001$
121 parameters	$\Delta\rho_{\max} = 0.21 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.72741 (16)	0.34999 (14)	0.68693 (8)	0.0383 (3)
N2	0.66107 (16)	0.19639 (15)	0.71630 (8)	0.0393 (3)
H2N	0.669 (2)	0.164 (2)	0.7817 (14)	0.052 (5)*
N3	0.61324 (14)	0.20352 (14)	0.55193 (8)	0.0345 (3)
N4	0.51489 (16)	-0.04181 (15)	0.64256 (9)	0.0398 (3)
H4A	0.488 (2)	-0.095 (2)	0.5842 (13)	0.049 (4)*
H4B	0.564 (2)	-0.106 (2)	0.6948 (13)	0.050 (4)*
N5	0.80901 (15)	0.62905 (15)	0.56803 (8)	0.0388 (3)
C1	0.69411 (16)	0.34759 (16)	0.58805 (9)	0.0331 (3)
C2	0.59494 (16)	0.11150 (17)	0.63532 (9)	0.0343 (3)
C3	0.74353 (16)	0.48955 (17)	0.52229 (9)	0.0343 (3)
C4	0.72299 (19)	0.47748 (19)	0.41755 (10)	0.0425 (3)
H4	0.6758	0.3786	0.3873	0.051*
C5	0.77288 (19)	0.6128 (2)	0.35870 (11)	0.0491 (4)
H5	0.7598	0.6076	0.2878	0.059*
C6	0.8419 (2)	0.7554 (2)	0.40516 (12)	0.0486 (4)
H6	0.8781	0.8489	0.3668	0.058*

supplementary materials

C7	0.85663 (19)	0.75796 (19)	0.50930 (12)	0.0455 (4)
H7	0.9029	0.8561	0.5408	0.055*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0476 (6)	0.0387 (6)	0.0282 (5)	-0.0007 (5)	-0.0076 (4)	0.0006 (4)
N2	0.0518 (7)	0.0392 (6)	0.0261 (6)	-0.0013 (5)	-0.0092 (5)	0.0000 (5)
N3	0.0362 (5)	0.0396 (6)	0.0273 (5)	0.0042 (4)	-0.0059 (4)	-0.0028 (4)
N4	0.0498 (7)	0.0397 (6)	0.0291 (6)	-0.0021 (5)	-0.0110 (5)	-0.0001 (5)
N5	0.0420 (6)	0.0409 (6)	0.0332 (6)	0.0030 (5)	-0.0043 (5)	0.0012 (5)
C1	0.0330 (6)	0.0385 (7)	0.0273 (6)	0.0057 (5)	-0.0056 (5)	-0.0035 (5)
C2	0.0357 (6)	0.0391 (7)	0.0274 (6)	0.0053 (5)	-0.0068 (5)	-0.0030 (5)
C3	0.0305 (6)	0.0419 (7)	0.0301 (6)	0.0076 (5)	-0.0028 (5)	-0.0010 (5)
C4	0.0442 (7)	0.0522 (8)	0.0310 (7)	0.0046 (6)	-0.0016 (5)	-0.0025 (6)
C5	0.0488 (8)	0.0683 (10)	0.0304 (7)	0.0080 (7)	0.0032 (6)	0.0071 (7)
C6	0.0444 (8)	0.0558 (9)	0.0457 (8)	0.0057 (7)	0.0046 (6)	0.0150 (7)
C7	0.0454 (8)	0.0447 (8)	0.0460 (8)	0.0013 (6)	-0.0035 (6)	0.0050 (6)

Geometric parameters (\AA , $^\circ$)

N1—C1	1.3221 (16)	N5—C3	1.3410 (17)
N1—N2	1.3708 (16)	C1—C3	1.4729 (18)
N2—C2	1.3422 (16)	C3—C4	1.3909 (18)
N2—H2N	0.902 (19)	C4—C5	1.380 (2)
N3—C2	1.3312 (17)	C4—H4	0.94
N3—C1	1.3656 (16)	C5—C6	1.374 (2)
N4—C2	1.3538 (18)	C5—H5	0.94
N4—H4A	0.896 (18)	C6—C7	1.377 (2)
N4—H4B	0.925 (17)	C6—H6	0.94
N5—C7	1.3354 (18)	C7—H7	0.94
C1—N1—N2	102.14 (10)	N5—C3—C4	122.10 (13)
C2—N2—N1	109.99 (11)	N5—C3—C1	116.99 (11)
C2—N2—H2N	129.2 (11)	C4—C3—C1	120.91 (12)
N1—N2—H2N	120.8 (11)	C5—C4—C3	119.01 (14)
C2—N3—C1	102.81 (10)	C5—C4—H4	120.5
C2—N4—H4A	116.4 (10)	C3—C4—H4	120.5
C2—N4—H4B	112.7 (10)	C6—C5—C4	119.12 (14)
H4A—N4—H4B	117.2 (14)	C6—C5—H5	120.4
C7—N5—C3	117.65 (12)	C4—C5—H5	120.4
N1—C1—N3	115.02 (12)	C5—C6—C7	118.36 (14)
N1—C1—C3	122.04 (12)	C5—C6—H6	120.8
N3—C1—C3	122.94 (11)	C7—C6—H6	120.8
N3—C2—N2	110.03 (12)	N5—C7—C6	123.76 (15)
N3—C2—N4	127.19 (11)	N5—C7—H7	118.1
N2—C2—N4	122.71 (12)	C6—C7—H7	118.1

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2N···N5 ⁱ	0.90 (2)	2.01 (2)	2.9010 (16)	171 (1)
N4—H4A···N3 ⁱⁱ	0.90 (2)	2.11 (2)	2.9971 (16)	172 (1)
N4—H4B···N1 ⁱ	0.93 (2)	2.19 (2)	3.0264 (16)	151 (1)

Symmetry codes: (i) $-x+3/2, y-1/2, -z+3/2$; (ii) $-x+1, -y, -z+1$.

Fig. 1




Fig. 2




Fig. 3

