

$C_{62}H_{89}N_3O_7$ requires C, 75.3; H, 9.08; N, 4.25%. Mass spectrum (FAB): m/z 988 [M^+], 1010 [$M^+ - H + Na$]. NMR ($CDCl_3$): 1H , δ 0.95 [s, 18H, $C(CH_3)_3$], 1.17 [s, 9H, $C(CH_3)_3$], 1.19 [s, 9H, $C(CH_3)_3$], 1.11–1.27 (br m, 18H, NCH_2CH_3), 3.19 (d, $J = 13$ Hz, 2H, CH_2), 3.24 (d, $J = 13$ Hz, 2H, CH_2), 3.27–3.47 (br m, 12H, NCH_2), 4.51 (d, $J = 13$ Hz, 2H, CH_2), 4.55 (d, $J = 14$ Hz, 2H, OCH_2), 5.05 (d, $J = 14$ Hz, 2H, OCH_2), 5.09 (d, $J = 13$ Hz, 2H, CH_2), 5.10 (s, 2H, OCH_2), 6.67 (d, $J = 2.3$ Hz, 2H, ArH), 6.70 (d, $J = 2.3$ Hz, 2H, ArH), 6.87 (s, 2H, ArH), 6.93 (s, 2H, ArH), 7.86 (s, 1H, OH). ^{13}C , δ 13.02 (br), 14.06, 14.25 (CH_2CH_3), 31.14, 31.46, 31.60 [$C(CH_3)_3$], 31.69, 32.09 (CH_2), 33.61, 33.73, 33.87 [$C(CH_3)_3$], 39.95, 40.22, 40.89, 41.44 (CH_2CH_3), 71.08, 72.59 (OCH_2), 124.69, 125.09, 125.31, 125.75 (aromatic CH), 127.67, 132.99, 133.20, 134.23, 140.25, 145.05, 145.58, 150.47, 152.71, 153.66 (aromatic C), 168.14, 169.74 (C=O).

5,11,17,23,29,35-Hexa-*tert*-butyl-36,39-dihydroxy-37,38,40,-41-tetrakis(diethylcarbamoylmethoxy)calix[6]arene (IV). A slurry of *p-tert*-butylcalix[6]arene (2.0 g, 2.1 mmol) and NaH (60% dispersion in oil, 1.1 g, ≈ 27 mmol) in dmf (10 cm^3) and thf (120 cm^3) was stirred at rt for 1 h, under an inert atmosphere. Potassium iodide (2.7 g, 16 mmol) and 2-chloro-*N,N*-diethylacetamide (2.46 g, 16 mmol) were then added, and the resulting mixture heated at reflux for 18 h. The solvents were then removed at reduced pressure, and 1 M HCl (100 cm^3) was added to the residue. The resulting off-white solid was collected by filtration, washed with water (3×50 cm^3), and dissolved in dichloromethane (50 cm^3). Hexane (approx. 50 cm^3) was added until a brown oil separated. The mother liquor was decanted from the oil, and upon further evaporation, white crystals of the product were precipitated. Recrystallisation from dichloromethane–acetonitrile gave the product as a white microcrystalline solid (0.8 g, yield 27%). (Found: C, 75.6; H, 8.74; N, 4.09. $C_{90}H_{128}N_4O_{10}$ requires C, 75.8; H, 9.05; N, 3.93%). Mass spectrum (FAB): m/z 1426 [M^+].

5,11,17,23,29,35-Hexa-*tert*-butyl-36,39-dimethoxy-37,38,40,-41-tetrakis(diethylcarbamoylmethoxy)calix[6]arene (V). A slurry of IV (0.10 g, 0.070 mmol) and K_2CO_3 (0.10 g, 0.72 mmol) in acetonitrile (50 cm^3) was heated at reflux for 1 h under an inert atmosphere. Methyl iodide (0.10 g, 0.69 mmol) was added, and the mixture heated at reflux for 12 h. The mixture was then filtered and the solvent removed at reduced pressure. The yellow residue was dissolved in dichloromethane (30 cm^3), and the solution was washed with water (3×20 cm^3). The organic layer was dried ($MgSO_4$), filtered, and ethanol added (20 cm^3). Upon evaporation the product was precipitated as off-white crystals (0.06 g, yield 60%). (Found: C, 76.0; H, 9.18; N, 3.84. $C_{92}H_{132}N_4O_{10}$ requires C, 76.0; H, 9.15; N, 3.85%). Mass spectrum (FAB): m/z 1454 [M^+]. NMR ($CDCl_3$): 1H , δ 1.05 [s, 36H, $C(CH_3)_3$], 1.37 [s, 18H, $C(CH_3)_3$], 1.15–1.25 (br m, 24H, NCH_2CH_3), 1.46 (s, 6H, OCH_3), 3.38 (d, $J = 15$ Hz, 4H, CH_2), 3.45–3.65 (br m, 16H, NCH_2), 3.99 (s, 4H, CH_2), 4.35 (d, $J = 15$ Hz, 4H, CH_2), 4.40 (d, $J = 12$ Hz, 4H, OCH_2), 4.69 (d, $J = 12$ Hz, 4H, OCH_2), 6.55 (d, $J = 2.1$ Hz, 4H, ArH), 7.21 (s, 4H, ArH), 7.61 (d, $J = 2.1$ Hz, 4H, ArH). ^{13}C , δ 12.87, 14.54 (CH_2CH_3), 31.29, 31.56 [$C(CH_3)_3$], 29.92, 32.36 (CH_2), 34.16, 34.21 [$C(CH_3)_3$], 40.11, 41.79 (CH_2CH_3), 58.80 (OCH_3), 72.18 (OCH_2), 123.30, 127.28, 127.88 (aromatic CH), 132.48, 132.72, 134.16, 145.79, 146.47, 152.30, 154.80 (aromatic C), 167.29 (C=O).

Complex syntheses

Complex 2a, [Fe(II – H)](ClO₄)₂. The calixarene II (0.15 g, 0.15 mmol) was dissolved in 1 : 1 dichloromethane–ethanol (2 ml). An approximately three fold excess of $Fe(ClO_4)_2 \cdot 6H_2O$ was added, resulting in a deep green solution which was filtered. Upon slow evaporation the product precipitated as small dark

green crystals. (Yield: 80%). (Found: C, 60.1; H, 7.21; N, 3.21. $C_{62}H_{88}Cl_2FeN_3O_{15}$ requires C, 60.0; H, 7.14; N, 3.38%). Mass spectrum (FAB): m/z 1043 [$\{M - 2(ClO_4)\}^+$], 1142 [$\{M - (ClO_4)\}^+$].

Complex 2b, [Fe(II – H)](FeCl₄)₂. The calixarene II (0.15 g, 0.15 mmol) was dissolved in 1 : 1 dichloromethane–ethanol (2 ml). An approximately three fold excess of $FeCl_3$ was added, resulting in a deep green solution which was filtered. Upon slow evaporation the product precipitated as dark green rhombic crystals suitable for X-ray diffraction studies. (Yield: 70%). (Found: C, 51.6; H, 6.12; N, 3.20. $C_{62}H_{88}Cl_8Fe_3N_3O_7$ requires C, 51.8; H, 6.17; N, 2.92%). Mass spectrum (FAB): m/z 1043 [$\{M - 2(FeCl_4)\}^+$].

Complex 4, [Fe(IV – H)](ClO₄)₂. The calixarene IV (0.14 g, 0.10 mol) was dissolved in 1 : 1 dichloromethane–ethanol (2 ml). An approximately three fold excess of $Fe(ClO_4)_2 \cdot 6H_2O$ was added, resulting in a deep green solution which was filtered. Upon slow evaporation the product precipitated as small dark green crystals, often contaminated with a red amorphous precipitate. (Yield: 40–50%). (Found: C, 61.6; H, 8.07; N, 3.01. $C_{90}H_{127}Cl_2FeN_4O_{18} \cdot 4H_2O$ requires C, 61.7; H, 7.77; N, 3.20%). Mass spectrum (FAB): m/z 1480 [$\{M - 2(ClO_4) - H\}^+$].

Electrochemistry

Following published procedures,^{12a} measurements were made using a MacLab Potentiostat, controlled by the Echem software package,^{12b} via a MacLab/2e. All reported voltammograms were recorded on the first sweep. Electrochemical solutions were prepared at 1.0×10^{-3} mol dm^{-3} in freshly distilled solvent. Bu_4NBF_4 was then added as supporting electrolyte (0.10 mol dm^{-3}). The electrochemical cell was an unsealed, one-compartment cell with a glassy carbon disc working electrode, an Ag/Ag^+ reference electrode (1.0×10^{-3} mol dm^{-3} $AgNO_3$, 0.10 mol dm^{-3} Bu_4NBF_4 in CH_3CN) and a platinum-wire counter electrode. Argon, saturated with the appropriate solvent, was used at all times for solution degassing. Ferrocene was used as an internal standard, except where peak overlap occurred; in these cases ferrocene was used as an external standard. The reference electrode was regularly checked against Fc/Fc^+ , and was found to be stable under the explored conditions. Potentials are reported relative to that of the Fc/Fc^+ couple as measured under the appropriate experimental conditions.

Structure determination

A full sphere of CCD area-detector data was measured at *ca.* 153 K [Bruker AXS instrument; SMART/SAINT/SADABS¹³ (*etc.*) proprietary software; monochromatic Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å; $2\theta_{max} = 58^\circ$] yielding 70920 reflections, merging after ‘empirical’ (multiscan) absorption correction to 18533 unique ($R_{int} = 0.028$), 13314 of these with $F > 4\sigma(F)$ being considered ‘observed’ and used in the full matrix least squares refinement. Anisotropic thermal parameter forms were refined for the non-hydrogen atoms, (x, y, z, U_{iso})_H also being refined. The chlorine atoms of one of the anions were modelled as disordered over two sets of sites, occupancies set at 0.5 after trial refinement and seemingly concerted with the disorder of a pair of nearby residues modelled as (half) water molecule oxygen atoms, no associated hydrogens being located; high ‘thermal motion’ in association with tail 2 of the ligand is presumably also a foil for unresolved disorder. Conventional residuals R, R_w on $|F|$ [weights $\{\sigma^2(F) + 4 \times 10^{-4}(F^2)\}^{-1}$] were 0.043, 0.048 at convergence. Pertinent results are given below and in the Tables and Figures. Neutral atom complex scattering factors were employed, computation using the Xtal 3.6 program system.¹³

Table 1 The iron atom environment in **2b**. *r* is the iron–oxygen atom distance (Å); other entries in the matrix are the angles (°) subtended at the metal by the relevant atoms at the head of the row and column^a

Atom	<i>r</i>	O(102)	O(21)	O(202)	O(31)	O(302)	O(41)
O(11)	2.300(4)	71.06(6)	74.79(5)	92.40(7)	138.91(6)	150.57(6)	84.09(6)
O(102)	2.064(2)		133.49(6)	82.82(7)	149.21(6)	79.51(7)	98.03(7)
O(21)	2.495(2)			67.93(7)	73.60(5)	129.34(6)	108.93(7)
O(202)	1.999(2)				99.33(7)	84.19(8)	175.88(8)
O(31)	2.365(2)					70.24(6)	82.02(6)
O(302)	2.045(2)						99.93(7)
O(41)	1.792(2)						

^a Fe–O(*n*1)–C(*n*01,*n*1) (*n* = 1,4) are: 114.7(1), 125.4(1); 113.0(1), 138.6(1); 114.9(1), 125.5(1); –, 164.2(2); Fe–O(*n*02)–C(*n*02) (*n* = 1–3) are: 122.8(2), 132.4(2), 127.4(2)°. O(*n*1)⋯O(*n* + 1 1) (*n* = 1, 4) are: 2.917(2), 2.913(2), 2.762(2), 2.766(2) Å. For the O(*n*1)₄ plane, deviations δO(*n*1) are 0.211(2), –0.201(2), 0.231(2), –0.245(2), δFe being 0.975(1) Å, dihedral angles of C₆(*n*) to it being 72.05(6), 67.71(7), 68.44(6), 28.47(7)°, *n* = 1/2, 2/3, 3/4, 4/1; 1/3, 2/4 being: 80.01(8), 81.28(8), 69.89(8), 77.62(8); 39.76(8), 83.87(8)°. δFe from the pendant O₂C₂N planes are: 0.622(4), 0.169(5), 0.168(4) Å; dihedral angles between the pendant planes and their parent C₆ plane are 84.74(9), 87.4(1), 85.1(1)°.

Crystal/refinement data for 2b. [Fe(II – H)](FeCl₄)₂·H₂O ≡ C₆₂H₉₀Cl₈Fe₃N₃O₈, *M* = 1456.6. Monoclinic, space group *P*2₁/*n* (*C*_{2h}, no. 14), *a* = 19.880(1), *b* = 14.765(1), *c* = 25.914(2) Å, β = 100.681(1)°, *V* = 7475 Å³. *D*_c (*Z* = 4) = 1.29₄ g cm^{–3}. μ_{Mo} = 9.1 cm^{–1}; specimen: 0.45 × 0.45 × 0.45 mm (rhomb); *T*_{min,max} = 0.67, 0.83. |Δρ_{max}| = 0.60(4) e Å^{–3}.

CCDC reference number 165283.

See <http://www.rsc.org/suppdata/dt/b1/b104963c/> for crystallographic data in CIF or other electronic format.

Results and discussion

Ligand syntheses

The required calix[4]arene ligands **I**, and **III**, were synthesised according to literature methods.^{9,10} The trisamide ligand **II** has been synthesised previously by adding the amide groups to a monoalkylated calix[4]arene derivative, followed by removal of the original substituent to give the trisamide.⁵ In this work, **II** was synthesised by adapting the published procedure for trialkylation of calix[4]arenes,¹¹ giving the required trisamide in modest yield. The synthesis of calix[6]arene derivative **IV** was serendipitous; under conditions expected to give the hexa-substituted calix[6]arene, a tetra-substituted product (by FAB MS) was obtained in significant and reproducible yield. The cause of this has not been clarified, but the synthesis provides a convenient route to this tetra-substituted product. The ¹H NMR of the product is extremely complicated, and is consistent with the molecule assuming a number of different rigid conformations in solution. Variable temperature NMR studies did not simplify the spectrum in the solvent used, and thus it was not possible to determine the substitution pattern of the calix[6]arene, although previous work with different alkylating agents indicated that a 1,2,4,5-substitution pattern was likely.^{1,14} Methylation of the remaining phenol groups of **IV** gave **V** which was readily characterised by ¹H and ¹³C NMR. The number of signals and integration of peaks in the ¹H NMR was found to be consistent with 1,2,4,5-substitution. From these studies, it was concluded that **IV** also has the 1,2,4,5-substitution pattern. The complex NMR spectra obtained for **IV** are presumably due to slow conformational change on the NMR timescale, caused by hydrogen bonds involving the unsubstituted phenol groups, although it is notable that the related bridged calix[6]arene tetraamide, where the remaining phenol groups are linked by a polyether bridge, is also conformationally restricted.¹⁵

Metal complexation

Iron complexation by the partly substituted calixarenes is clearly indicated by the rapid formation of an intensely coloured solution (purple for **I**, green for **II** and **IV**) when an iron salt is added to a dichloromethane–ethanol solution of the calixarene. As reported previously, **III** also forms an intense green solution initially, fading to a pale green solution over

24 h, from which the colourless iron(II) complex crystallises.⁷ Careful addition of an iron(III) perchlorate solution to dichloromethane–ethanol solutions of the calixarenes **I**, **II** and **IV** shows an increase in a broad absorbance in the visible region (**I**, 580 nm; **II**, **IV**, 800 nm), characteristic of a phenolate–iron(III) charge transfer band,¹⁶ up to a 1 : 1 metal : ligand ratio, after which no further changes are observed. This is consistent with the formation of a 1 : 1 complex in solution for all the calixarenes studied. The complexes are readily isolated by allowing the solutions to evaporate slowly, and microanalytical and FAB MS data were consistent with the formulation of the products as [Fe(**I** – 2H)](ClO₄)₂ (**1**), [Fe(**II** – H)](ClO₄)₂ (**2a**) and [Fe(**IV** – H)](ClO₄)₂ (**4**). Attempts to study the complexation process in more polar solvents such as dmf and dmsu were unsuccessful, and it appears that iron complexation does not occur under these conditions. Addition of a base is often required to induce metal complexation with ionisable calixarenes.^{1,17} In the present case, addition of triethylamine to the various reaction mixtures resulted in rapid precipitation of an amorphous red-brown solid which did not contain a significant proportion of the calixarene (presumably the solid is ‘iron hydroxide’ as no attempt was made to perform the reactions under strictly anhydrous conditions).

Reaction of trisamide **II** with iron(III) chloride rather than the perchlorate, resulted in the isolation of complex **2b**. This complex formed crystals appropriate for a single crystal X-ray study, the results of which are consistent with the formulation [Fe(**II** – H)](FeCl₄)₂·H₂O, one formula unit, devoid of crystallographic symmetry, comprising the asymmetric unit of the structure. Microanalytical data indicate that the solvent water molecule is lost upon drying of the product. The structure of the complex cation is shown in Fig. 1, and pertinent bond distances and angles are listed in Table 1. The calix[4]arene assumes a distorted cone conformation, although, given the dihedral angle between phenyl ring 4 and the phenol O₄ plane [28.47(7)°; see Table 1], it may be more appropriate to describe the conformation qualitatively as “up, up, up, out”.¹ The cavity defined by the phenyl rings is unoccupied.

The Fe atom lies in the coordination environment offered by the seven O atoms of the calixarene in an unsymmetrical manner, quasi-pentagonal bipyramidal, O(41), O(202) quasi-axial (Fig. 1). The shortest Fe–O distance involves the unsubstituted phenol atom [Fe–O(41), 1.792(2) Å], and the longest is found for the opposite phenol ether O atom [Fe–O(21), 2.495(2) Å]. The Fe–O distances increase in the order phenol, amide, phenol ether O atoms; this was also observed in the Fe(III) complex of diamide **I**,⁸ and with the exception of Fe–O(21), the magnitudes of the comparable Fe–O distances in the two structures differ by less than 0.07 Å.

Electrochemical studies

The metal complexes **1**, **2a**, **3** and **4** dissolve readily in acetonitrile and dichloromethane, and the electrochemical




Fig. 1 Projections of the cation of **2b** (a) down and (b) quasi-normal to the cone axis. 50% Displacement ellipsoids are shown for the non-hydrogen atoms.

behaviour of the complexes was studied in both solvents. Cyclic voltammograms for **3** and **4** were poorly defined and gave inconsistent results. The voltammograms of **1** and **2a** were more readily studied, and the results reported here will focus on these systems.

Representative cyclic voltammograms of **1** and **2a** in acetonitrile are shown in Fig. 2. The complex **2a** undergoes a reduction at $E_{1/2} = -0.15$ V (vs. Fc^+/Fc); varying the scan rate showed that the process is diffusion controlled, with changes in the oxidation and reduction potentials of less than 15 mV. The separation between the oxidation and reduction potentials at a scan rate of 200 mV s^{-1} is 90 mV, compared with 75 mV for ferrocene under the same conditions. A reduction process is also observed for **1** in acetonitrile (Fig. 2); in this case the $E_{1/2}$ is -0.73 V, and the process is somewhat less reversible, with a peak separation of 105 mV at a scan rate of 200 mV s^{-1} , increasing to 195 mV at 1000 mV s^{-1} . The behaviour in dichloro-




Fig. 2 Cyclic voltammograms of (a) **2a**, and (b) **1** in CH_3CN ($0.10 \text{ mol dm}^{-3} \text{ Bu}_4\text{NBF}_4$), reported vs. Fc^+/Fc . Scan rate 200 mV s^{-1} .

methane solvent is comparable to that observed in acetonitrile, albeit with poorer reversibility; for **1**, the $E_{1/2}$ is -0.77 V, and for **2a**, -0.25 V (vs. Fc^+/Fc).

To examine this process further, controlled potential bulk electrolysis of the solutions was performed. During this procedure the solutions became colourless, and UV/vis spectroscopy confirmed that the peak at 580 nm for **1**, and 800 nm for **2a** were no longer present. This observation is consistent with the reduction of iron(III) to iron(II).¹⁸ When performed in acetonitrile or dichloromethane solvents the reduction was irreversible, indicating that the complex may have dissociated or precipitated. It was found that the complexes could be dissolved (as judged by the UV/vis spectra) for bulk electrolysis experiments (although the solutions were colourless after 24 h indicating gradual dissociation of the complexes). In this solvent the bulk reduction process was reversible, and given that the complexes could not be synthesised in this solvent, it seems likely that the iron(III) complex is reduced without significant dissociation of the resulting iron(II) complex. These measurements show that the trisamide iron(III) complex **2a**, is more readily reduced than the bisamide complex **1**, consistent with increased stability of the higher metal oxidation state with increased number of phenolate O donor atoms.¹⁶

Cyclic voltammetry also shows that the two complexes undergo oxidation processes, although this aspect of the electrochemistry was not studied in detail. A thorough study of the anodic oxidation of **1** in dichloromethane solvent has recently been reported,¹⁹ showing that the calixarene undergoes two irreversible oxidations (with a minor, unassigned oxidation peak also observed in cyclic voltammograms) followed by chemical reactions to give electrochemically active species (an ECE-type mechanism); preparative coulometry produced the mono- and di-calixquinones resulting from oxidation of the unsubstituted phenol groups. In the present case, the complex **2a** was found to undergo an irreversible oxidation at 1.1 V (Fig. 3) in acetonitrile, after which a new reduction wave was observed, albeit overlapping with the process assigned to the iron(III) reduction, at approximately -0.3 V. This is consistent with a chemical reaction step following the initial oxidation, as observed for the uncomplexed calixarene **I**.

It is interesting to note that the cyclic voltammogram for **1** in acetonitrile exhibits an oxidation that is at least quasi-reversible at $E_{1/2} 0.62$ V (Fig. 4); scanning to higher potentials shows a second, irreversible, oxidation process at 1.1 V. The most probable assignment for these oxidation processes, is ligand-centred oxidation to give phenoxyl radicals, as is observed in the free




Fig. 3 Cyclic voltammograms of **2a** in CH_3CN ($0.10 \text{ mol dm}^{-3} \text{ Bu}_4\text{NBF}_4$), reported vs. Fc^+/Fc . Scan rate 200 mV s^{-1} .




Fig. 4 Cyclic voltammogram of **1** in CH_3CN ($0.10 \text{ mol dm}^{-3} \text{ Bu}_4\text{NBF}_4$), reported vs. Fc^+/Fc . Scan rate 500 mV s^{-1} .

ligand **1**.¹⁹ Presumably coordination to the metal ion stabilises the radical species sufficiently for the process to be quasi-reversible in some cases (Fig. 4), whereas in the free calixarene, this is not observed. Electrochemical generation of a range of coordinated phenoxyl radicals has been reported; for example, Wieghardt *et al.* have described detailed characterisation of such species generated from phenol-functionalised triaza-cyclononane derivatives.¹⁶ It should be noted, however, that electrochemical oxidation of aryl ether units has also been reported in studies of a *tert*-butylcalix[4]arene tetraester,²⁰ and thus definitive assignment of the anodic processes observed requires further work.

Conclusions

Iron(III) complexes of the calix[4]arene trisamide **II**, and the calix[6]arene tetraamide **IV**, have been synthesised, both forming 1 : 1 complexes under the conditions explored; the former was structurally characterised as the FeCl_4^- salt. The electrochemical behaviour of the trisamide calix[4]arene iron complex **2a**, and the previously described bisamide calix[4]arene iron complex **1**, was studied. Reduction of the complexes gave the related iron(II) species, with complex **2a** being more readily

reduced than **1**. Oxidation of the complexes was also observed, this process being assigned to oxidation of the phenoxy moieties.

Acknowledgements

This work was supported in part by the ARC Small Grant Scheme, and an Australian Postdoctoral Research Fellowship. M. I. O. is grateful to Assoc. Prof. Jack Harrowfield for providing laboratory facilities.

References

- 1 C. D. Gutsche, in *Calixarenes Revisited*, ed. J. F. Stoddart, The Royal Society of Chemistry, Cambridge, 1998.
- 2 F. Arnaud-Neu, M. A. McKerverey and M.-J. Schwing-Weill, in *Calixarenes 2001*, ed. Z. Asfari, V. Böhmer, J. Harrowfield and J. Vicens, Kluwer Academic Publishers, Dordrecht, Boston, London, 2001, p. 385.
- 3 M. A. McKerverey, M. J. Schwing-Weill and F. Arnaud-Neu, in *Comprehensive Supramolecular Chemistry*, ed. J. L. Atwood, J. E. D. Davies, D. D. McNicol and F. Vögtle, Pergamon, New York, 1996, p. 537; A. Casnati, L. Baldini, N. Pelizzi, K. Rissanen, F. Uguzzoli and R. Ungaro, *J. Chem. Soc., Dalton Trans.*, 2000, 3411; F. Arnaud-Neu, S. Barbosa, A. Casnati, A. Pinalli, M. J. Schwing-Weill and R. Ungaro, *New J. Chem.*, 2000, **24**, 967; M. F. Arnaud-Neu, S. Barbosa, F. Berny, A. Casnati, N. Muzet, A. Pinalli, R. Ungaro, M. J. Schwing-Weill and G. Wipff, *J. Chem. Soc., Perkin Trans. 2*, 1999, 1727; N. Muzet, G. Wipff, A. Casnati, L. Domiano, R. Ungaro and F. Uguzzoli, *J. Chem. Soc., Perkin Trans. 2*, 1996, 1065.
- 4 P. D. Beer, M. G. B. Drew, A. Grieve and M. I. Ogden, *J. Chem. Soc., Dalton Trans.*, 1995, 3455; P. D. Beer, M. G. B. Drew, M. Kan, P. B. Leeson, M. I. Ogden and G. Williams, *Inorg. Chem.*, 1996, **35**, 2202; P. D. Beer, M. G. B. Drew and M. I. Ogden, *J. Chem. Soc., Dalton Trans.*, 1997, 1489.
- 5 M. P. Oude Wolbers, F. C. J. M. Van Veggel, R. H. M. Heeringa, J. W. Hofstraat, F. A. J. Geurts, G. J. Van Hummel, S. Harkema and D. N. Reinhoudt, *Liebigs Ann. Chem.*, 1997, 2587.
- 6 F. Arnaud-Neu, G. Barrett, D. Corry, S. Cremin, G. Ferguson, J. F. Gallagher, S. J. Harris, M. A. McKerverey and M. J. Schwing-Weill, *J. Chem. Soc., Perkin Trans. 2*, 1997, 575; C. Wieser, D. Matt, J. Fischer and A. Harriman, *J. Chem. Soc., Dalton Trans.*, 1997, 2391.
- 7 P. D. Beer, M. G. B. Drew, P. B. Leeson and M. I. Ogden, *J. Chem. Soc., Dalton Trans.*, 1995, 1273.
- 8 P. D. Beer, M. G. B. Drew, P. B. Leeson and M. I. Ogden, *Inorg. Chim. Acta*, 1996, **246**, 133.
- 9 E. M. Collins, M. A. McKerverey, E. Madigan, M. B. Moran, M. Owens, G. Ferguson and S. J. Harris, *J. Chem. Soc., Perkin Trans. 1*, 1991, 3137.
- 10 A. Arduini, E. Ghidini, A. Pochini, R. Ungaro, G. D. Andreotti, G. Calestani and F. Uguzzoli, *J. Inclusion Phenom.*, 1988, **6**, 119.
- 11 C. D. Gutsche, B. Dhawan, J. A. Levine, K. H. No and L. J. Bauer, *Tetrahedron*, 1983, **39**, 409.
- 12 (a) P. D. Beer, Z. Chen and M. I. Ogden, *J. Chem. Soc., Faraday Trans.*, 1995, **91**, 295; (b) EChem software, ADInstruments, Sydney, Australia, 1995, version 1.3.1.
- 13 SMART/SAINT, Software for the CCD Detector System, Bruker Analytical X-Ray Systems, Inc., Madison, WI, 1998; G. M. Sheldrick, SADABS, Program for Siemens area detector absorption correction, Institut für Anorganische Chemie, Universität Göttingen, 1996; ed. S. R. Hall, D. J. du Boulay and R. Olthoff-Hazekamp, The Xtal 3.6 System, University of Western, Australia, 2000.
- 14 J. M. Harrowfield, M. Mocerino, B. W. Skelton and A. H. White, *Aust. J. Chem.*, 1998, **51**, 123.
- 15 A. Casnati, P. Jacopozzi, A. Pochini, F. Uguzzoli, R. Cacciapaglia, L. Mandolini and R. Ungaro, *Tetrahedron*, 1995, **51**, 591.
- 16 B. Adam, E. Bill, E. Bothe, B. Goerd, G. Haselhorst, K. Hildenbrand, A. Sokolowski, S. Steenken, T. Weyhermuller and K. Wieghardt, *Chem. Eur. J.*, 1997, **3**, 308.
- 17 P. Thuéry, M. Nierlich, J. Harrowfield and M. Ogden, in *Calixarenes 2001*, ed. Z. Asfari, V. Böhmer, J. Harrowfield and J. Vicens, Kluwer Academic Publishers, Dordrecht, Boston, London, 2001, p. 561.
- 18 D.-H. Jo, Y.-M. Chiou and L. Que, Jr., *Inorg. Chem.*, 2001, **40**, 3181.
- 19 R. Vataj, A. Louati, C. Jeunesse and D. Matt, *Electrochem. Commun.*, 2000, **2**, 769.
- 20 A. Pailleret and D. W. M. Arrigan, *Electrochem. Commun.*, 2001, **3**, 24.