

# Estimating eligibility for lung cancer screening in an Australian cohort, including the effect of spirometry

**L**ung cancer causes more deaths than any other cancer, and is a leading cause of disease-related burden in Australia.<sup>1</sup> The United States National Lung Screening Trial (NLST) showed that screening high risk current and former smokers for lung cancer with low dose chest computed tomography (LDCT) can significantly reduce lung cancer mortality.<sup>2</sup> LDCT lung cancer screening is recommended in the US for eligible individuals.<sup>3</sup>

Unlike other cancer screening programs that select participants according to the risk factors of age and sex, eligibility for lung cancer screening is more complex. The current US screening criteria, as defined by the US Preventive Service Task Force (USPSTF), are that current or former smokers are eligible if they are aged 55–80 years, have a smoking history of at least 30 pack-years, and, if a former smoker, have quit less than 15 years ago.<sup>3</sup> However, this approach is not necessarily the most efficient for identifying those who may benefit from screening.<sup>4</sup> Using probabilistic multivariate lung cancer risk prediction models to select individuals for screening may improve overall efficiency and cost-effectiveness, and are recommended by international organisations.<sup>5–7</sup>

The PLCO<sub>m2012</sub> lung cancer risk model was developed during the US Prostate, Lung, Colorectal, Ovarian Screening (PLCO) Trial, and has been validated in the NLST cohort.<sup>8</sup> This model uses data on age, smoking status, duration and intensity, family history, body mass index (BMI), and comorbidities (including self-reported chronic obstructive pulmonary disease [COPD], chronic bronchitis and emphysema) to estimate an individual's risk of developing lung cancer within 6 years.<sup>8</sup> A PLCO<sub>m2012</sub> lung cancer risk of greater than 1.5% has been proposed as an

## Abstract

**Objectives:** To estimate the proportion of ever-smokers who are eligible for lung cancer screening in an Australian cohort, and to evaluate the effect of spirometry in defining chronic obstructive pulmonary disease (COPD) when assessing screening eligibility.

**Design:** Cross-sectional study of 3586 individuals aged 50–68 years who live in the Busselton Shire of Western Australia.

**Outcomes:** Proportion of ever-smokers eligible for lung cancer screening based on United States Preventive Services Task Force (USPSTF) criteria, and PLCO<sub>m2012</sub> lung cancer risk >1.5%. The effect of using self-reported COPD, symptoms consistent with COPD, or spirometry to define COPD for screening eligibility according to the PLCO<sub>m2012</sub> criteria.

**Results:** Of ever-smokers aged 55–68 years, 254 (20.1%) would be eligible for screening according to USPSTF criteria; fewer would be eligible according to PLCO<sub>m2012</sub> criteria (225, 17.9%;  $P = 0.004$ ). This is equivalent to 8.9–10.0% of the total population aged 55–68 years, which suggests about 450 000 individuals in Australia may be eligible for lung cancer screening. The proportions of eligible participants were not significantly different whether spirometry results or symptoms consistent with COPD were used to determine PLCO<sub>m2012</sub> risk.

**Conclusions:** The proportion of ever-smokers in this population who were eligible for lung cancer screening was 17.9–20.1%. Using symptoms to define COPD is an appropriate surrogate measure for spirometry when determining the presence of COPD in this population. There are significant challenges for policy makers on how to identify and recruit these eligible individuals from the wider population.

alternative lung cancer screening eligibility criterion.<sup>9</sup>

COPD is an independent risk factor for the development of lung cancer, and self-reported, doctor-diagnosed COPD and chronic bronchitis or emphysema are included in the PLCO<sub>m2012</sub> risk prediction model (but not in the USPSTF eligibility criteria).<sup>8</sup> There are, however, significant differences in the proportions of individuals who report a diagnosis of COPD, who have symptoms consistent with COPD, or who have airflow obstruction detectable by spirometry.<sup>10</sup> Including airflow obstruction improves the accuracy of a lung cancer risk prediction that does not incorporate self-reported COPD.<sup>11</sup> Incorporating spirometry results into eligibility assessment for lung cancer screening has been suggested, but the impact of adding

spirometry-defined COPD to the existing eligibility criteria of self-reported COPD or symptoms consistent with COPD is not known.<sup>12</sup>

In Australia, about 2.2 million people aged 55–75 years are current or former smokers, but the proportion of those eligible for lung cancer screening, using either USPSTF or PLCO<sub>m2012</sub> criteria, is unknown.<sup>13</sup> Estimating the proportion of people eligible for lung cancer screening will both aid policy makers when planning future capacity and assessing infrastructure requirements, and help estimate screening participation rates.

The aims of this study were to estimate the proportion of an Australian cohort that is eligible for lung cancer screening, and to examine the utility of including spirometry-defined COPD as part of screening eligibility assessment.

David Manners  
MBBS(Hons)<sup>1</sup>

Jennie Hui  
PhD<sup>2,3</sup>

Michael Hunter  
PhD(Dist)<sup>2,3</sup>

Alan James  
MBBS, FRACP, PhD<sup>3</sup>

Matthew W Knuiman  
PhD<sup>2</sup>

Annette McWilliams  
FRACP, FRCP, MD<sup>4</sup>

Siobhain Mulrennan  
MRCP, MD, FRACP<sup>1,2</sup>

Arthur W (Bill) Musk  
MBBS, MD, FRACP<sup>2</sup>

Fraser JH Brims  
MD, MRCP, FRACP<sup>1,3</sup>

<sup>1</sup>Sir Charles Gairdner  
Hospital,  
Perth, WA.

<sup>2</sup>Busselton Population  
Medical Research Institute,  
Busselton, WA.

<sup>3</sup>University of Western  
Australia,  
Perth, WA.

<sup>4</sup>Fiona Stanley Hospital,  
Perth, WA.

fraser.brimbs@  
uwa.edu.au

doi:10.5694/mja16.00043

## Methods

The Busselton Healthy Ageing Study (BHAS) is a cross-sectional study of adults born between 1946 and 1964 who live in the Busselton Shire of Western Australia.<sup>14</sup> Participants were recruited sequentially at random between 2010 and 2013 from the electoral roll; the participation rate was greater than 80% of contacted individuals. Participants completed a self-administered questionnaire and physical assessments, including spirometry. The upper age limit of participants included in our study (68 years) was determined by the BHAS design.

Smoking status, duration and intensity were self-reported. Ever-smokers responded positively to “Have you ever smoked cigarettes?” Current smokers responded positively to “Do you currently smoke manufactured or hand-rolled cigarettes?”, and former smokers responded negatively. Pack-years of smoking were determined by multiplying the duration of smoking in years by the number of 20-cigarette packs smoked each day. Personal history of any cancer other than non-melanomatous skin cancer, family history of lung cancer, and other demographic information were self-reported.

BMI was calculated from the measured height and weight. Pre- and post-bronchodilator forced expiratory volume in one second (FEV<sub>1</sub>) and forced vital capacity (FVC) were measured using an EasyOne spirometer (NDD Medical Technologies) before and after administration of 200 µg of inhaled salbutamol with a metered dose inhaler and spacer. Participants were asked to withhold any bronchodilator medication, if possible, for 12 hours before testing.

Individuals were excluded from the analysis if they had a history of lung cancer, if BMI or demographic data were missing, or there was insufficient data on smoking duration or intensity to calculate the number of pack-years.

Lung cancer risk was calculated with the PLCO<sub>m2012</sub> 6-year lung cancer risk prediction model.<sup>8</sup> For the

purposes of this model, COPD was defined in three ways:

- self-reported, doctor-diagnosed COPD, defined as a positive response to “Has your doctor ever told you that you had COPD?”;
- symptoms consistent with COPD, defined as a positive response to “Do you cough on most days for as much as 3 months each year?” or “Do you bring up phlegm from your chest on most days for as much as 3 months each year?” or “Has your chest made a wheezing or whistling sound in the last 12 months?”; and
- spirometry-defined COPD, defined by the presence of post-bronchodilator airflow obstruction (FEV<sub>1</sub>/FVC ratio <0.7), excluding those with more than 12% improvement in FEV<sub>1</sub> or FVC after bronchodilator medication.

Eligibility for lung cancer screening was based on two criteria: the current USPSTF recommendation<sup>3</sup> and a PLCO<sub>m2012</sub> 6-year lung cancer risk of >1.5%, as recommended by Tammemägi and colleagues.<sup>8</sup> To estimate the proportion of the Australian population who are eligible for screening, an estimate of 2.2 million Australians aged 55–75 years who are current or former smokers<sup>13</sup> was multiplied by our estimate of the proportion of potentially eligible ever-smokers in the Busselton population according to USPSTF criteria.

Statistical analyses were performed in SPSS 22 (IBM); 95% confidence intervals for proportions were calculated using continuity adjustment. Comparisons of the proportions of independent and dependent samples were performed with  $\chi^2$  and McNemar tests respectively. Independent sample *t* tests compared the means of independent samples. Statistical significance was defined as  $P < 0.05$ .

## Ethics approval

The BHAS and the reported study received human research ethics

approval from the University of Western Australia Human Research Ethics Committee (references RA/4/A/2203, RA/4/1/7891).

## Results

Of all 3586 BHAS participants aged 50–68 years, 1903 (53.1%) were current or former smokers. Of these ever-smokers, 36 were excluded (16 had incomplete smoking information, 15 reported smoking less than one cigarette per day, one had prior lung cancer, and demographic or BMI data were missing for four), leaving 1867 participants for the analysis. In the 55–68-year-old age group, 1260 (49.9%) were ever-smokers (including 240 [6.7%] current smokers), 1241 (49.2%) were never-smokers, and 22 (0.9%) had incomplete smoking data.

The demographic details of all BHAS ever-smokers in two age groups, 50–68 years and 55–68 years, are summarised in [Box 1](#). The 55–68-year-old age group was selected as corresponding to the age cohort currently eligible for lung cancer screening in the US.

## Eligibility for lung cancer screening

Of ever-smokers aged 55–68 years, 254 (20.1%; 95% CI, 17.9–22.3%) would be eligible for screening according to USPSTF criteria. Using the PLCO<sub>m2012</sub> criteria with self-reported COPD, fewer people (225, 17.9%) would be eligible (95% CI, 15.8–20.0%;  $P = 0.004$ ).

The proportion of ever-smokers eligible for screening increased with age in the PLCO<sub>m2012</sub> model, but was stable when using USPSTF criteria ([Box 2](#)). Of the entire age-defined cohort, 2.6% would be eligible using PLCO<sub>m2012</sub> criteria but not according to USPSTF criteria, and 4.9% would be eligible using USPSTF criteria but not PLCO<sub>m2012</sub> criteria; 15.2% would be eligible according to both criteria sets ([Box 3](#)). This suggests that about 450 000 individuals in Australia may be eligible for lung cancer screening.

### 1 Demographic characteristics of ever-smokers in the Busselton Healthy Ageing Study

	50–68 years old	55–68 years old
Number	1867	1260
Age (years), mean ± SD	58.0 ± 4.8	60.7 ± 3.4
Sex (women)	927 (49.7%)	603 (47.9%)
Ethnic background		
European	1848 (99.0%)	1249 (99.1%)
Asian	12 (0.6%)	5 (0.4%)
Aboriginal and/or Torres Strait Islander	6 (0.3%)	5 (0.4%)
Māori or Pacific Islander	1 (0.1%)	1 (0.1%)
Body mass index (kg/m <sup>2</sup> ), mean ± SD	28.46 ± 4.90	28.70 ± 4.95
Smoking status		
Current	368 (19.7%)	240 (19.0%)
Former	1499 (80.3%)	1020 (81.0%)
Total pack-years, mean ± SD		
Ever-smokers	19.2 ± 19.6	20.5 ± 20.7
Current smokers	31.5 ± 20.0	32.1 ± 19.3
Former smokers	16.2 ± 18.4	17.8 ± 20.1
Highest education level obtained		
Did not complete high school	36 (1.9%)	27 (2.1%)
High school	1000 (53.6%)	702 (55.7%)
TAFE or other college	555 (29.7%)	352 (27.9%)
University	276 (14.8%)	179 (14.2%)
Previous cancer (excluding non-melanomatotic skin cancer)	155 (8.3%)	122 (9.7%)
Family history of lung cancer	146 (7.8%)	106 (8.4%)
Chronic obstructive pulmonary disease (COPD)		
Self-reported COPD	16 (0.9%)	13 (1.0%)
Symptoms consistent with COPD	617 (33.0%)	441 (35.0%)
Spirometry-defined COPD	179/1191 (15.0%)	108/782 (13.8%)

TAFE = Technical and Further Education. ♦

#### Effect of COPD definition on lung cancer risk and eligibility

Of the ever-smokers in the BHAS population, 1191 had concurrent spirometry, of whom 782 were aged 55–68 years. Using the PLCO<sub>m2012</sub> criteria, the proportion eligible for screening was 16.4% (using self-reported COPD), 18.8% (symptoms consistent with COPD), or 17.5% (airflow obstruction on spirometry) (Box 4). Fewer people were eligible for screening using self-reported than spirometry-defined COPD ( $P < 0.01$ ); there was no statistical difference between the proportions of eligible participants when spirometry-defined or symptom-defined COPD was used ( $P = 0.06$ ).

When spirometry-defined COPD was added to symptom-defined COPD to determine PLCO<sub>m2012</sub> risk, an additional seven participants (0.9%) would have been eligible for screening. If normal spirometry was used to exclude COPD in those who reported symptoms of COPD, 17 participants (2.1%) would be deemed ineligible for screening.

Of the 409 ever-smokers aged 50–55 years (and therefore ineligible for screening according to current criteria) with spirometry data, 18 (4.4%; 95% CI, 2.8–6.9%), 36 (8.8%; 95% CI, 6.4–11.9%) or 25 (6.1%; 95% CI, 4.2–8.9%) would have been eligible based on a PLCO<sub>m2012</sub> risk

greater than 1.5% when COPD was defined by doctor diagnosis, symptoms, or spirometry respectively.


#### Discussion

Our study suggests that 17.9% (PLCO<sub>m2012</sub>) to 20.1% (USPSTF criteria) of current or former smokers aged 55–68 years in Busselton would be eligible for lung cancer screening, corresponding to 8.9% (PLCO<sub>m2012</sub>) and 10% (USPSTF) of all those in the BHAS cohort aged 55–68 years. This is the first time that the proportion of ever-smokers eligible for lung cancer screening has been estimated in an Australian population. While small studies have shown the feasibility of lung cancer screening in Australia, significant hurdles remain before screening is endorsed locally.<sup>15–18</sup>

Our finding that the USPSTF criteria identified a higher proportion of eligible individuals than the PLCO<sub>m2012</sub> criteria is consistent with previous reports.<sup>9</sup> Our study could not validate the predictive performance of each criterion because mature lung cancer outcomes were not available. Retrospective studies suggest that the PLCO<sub>m2012</sub> model performs better than the USPSTF criteria in identifying high risk individuals, but prospective impact analyses in screening cohorts are required to determine which eligibility criteria should be preferred in the Australian context.


There have been four previous estimates of the proportion of the US population who would be eligible for screening (Box 5).<sup>9,19–21</sup> Three of these studies estimated that 18.8–24.4% of ever-smokers and 10.5–14.3% of all individuals in the respective age cohorts were eligible for screening according to USPSTF criteria.<sup>19–21</sup> These estimates are broadly comparable with our estimates of the proportion of eligible individuals in the BHAS. On the other hand, fewer BHAS ever-smokers would be eligible for screening than estimated for the PLCO intervention arm ever-smokers (17.9% *v* 34.6%;  $P < 0.001$ ).<sup>9</sup> This observation may partly be explained by differences in tobacco exposure between the cohorts; the

**2 Eligibility for lung cancer screening of 1260 55–68-year-old ever-smokers in Busselton, according to USPSTF and PLCO<sub>m2012</sub> criteria\***



PLCO = Prostate, Lung, Colorectal, Ovarian Screening Trial; USPSTF = US Preventive Service Task Force. \* Calculated using doctor-diagnosed chronic obstructive pulmonary disease, with 95% confidence intervals. ♦

**3 Schematic representation of the proportion of current and former smokers eligible for screening**



A = Ineligible current smokers (*n* = 103; 8.2%); B = Ineligible former smokers (*n* = 870; 69.0%); C = Individuals eligible on both USPSTF and PLCO<sub>m2012</sub> criteria (*n* = 192; 15.2%); D = Individuals eligible only on PLCO<sub>m2012</sub> criteria (*n* = 33; 2.6%); E = Individuals eligible only on USPSTF criteria (*n* = 62; 4.9%). The areas of the components of this figure are approximately proportional. ♦

**4 Proportions of 782 ever-smokers aged 55–68 years eligible for screening based on PLCO<sub>m2012</sub> criteria, using different definitions of chronic obstructive pulmonary disease (COPD)**

COPD definition for estimating PLCO <sub>m2012</sub> lung cancer risk	PLCO <sub>m2012</sub> risk > 1.5%	<i>P</i> *
Self-reported COPD	128 (16.4%; 95% CI, 13.8–19.0%)	< 0.01
Symptom-defined COPD	147 (18.8%; 95% CI, 16.1–24.5%)	0.06
Spirometry-defined COPD	137 (17.5%; 95% CI, 14.8–20.2%)	—

\* *v* spirometry-defined COPD (McNemar test). ♦

proportion of current and former smokers were similar, but BHAS participants reported lower smoking intensity (mean pack-years, BHAS *v* PLCO: current smokers, 32.1 *v* 40.8, *P* < 0.001; former smokers, 17.8 *v* 25.8; *P* < 0.001).<sup>22</sup>

There are two possible explanations for the lower smoking intensity in the BHAS cohort. The first relates to historical trends in smoking. Enrolment for the PLCO trial occurred between 1993 and 2001, whereas BHAS enrolment commenced in 2010.<sup>14,23</sup> Smoking consumption in Australia declined significantly between 2001 and 2010 among both men and women.<sup>24</sup> The second explanation is a potential healthy community effect after repeated cross-sectional health surveys in the Busselton community since 1966.<sup>14</sup>

The current study has highlighted the known challenges in defining and diagnosing COPD. Only 1% of 55–68-year-old BHAS ever-smokers reported being diagnosed with COPD, whereas 13.8% met the spirometric definition of COPD and 35.0% reported symptoms consistent with the disease. A similar discrepancy was previously described in a larger Australian study.<sup>10</sup> Airflow obstruction on spirometry is not only crucial to diagnosing COPD, but is an independent predictor of increased lung cancer incidence in screening studies, and may predict over-diagnosis.<sup>12</sup> Incorporating spirometry into screening eligibility assessment has been suggested, but this would require direct patient contact.<sup>12</sup> Our results suggest that performing spirometry is unlikely to meaningfully improve the identification of eligible individuals, with just one additional individual from every 100 ever-smokers assessed becoming eligible. We therefore contend that, in the absence of more epidemiological data comparing the relative lung cancer risks associated with airflow obstruction and with symptoms or self-reported COPD, defining COPD on the basis of symptoms is an appropriate surrogate measure for spirometry.

Policy makers in jurisdictions that recommend population-based cancer screening, such as Australia, are faced by significant implementation

## 5 Summary of studies reporting proportions of the population aged 55 years or more who are eligible for lung cancer screening

	BHAS	PLCO intervention arm <sup>9,22</sup>	US Health and Retirement Study <sup>19</sup>	US National Health Interview Survey <sup>20</sup>	US National Health and Nutrition Examination Survey <sup>18</sup>
Sample size	2523	77 445	16 901	7138	2562
Recruitment period	2010–2013	1993–2001	2012	2011	2007–2012
Age range, years	55–68	55–74	55–80	55–74	55–77
Smoking status					
Never	49.2%	45.1%	42.5%	NR	NR
Current	9.5%	10.4%	14.4%	NR	NR
Former	40.4%	42.0%	39.6%	NR	NR
Missing or incomplete	0.9%	2.5%	3.5%	NR	NR
Eligible individuals as proportion of entire cohort					
USPSTF criteria	10.0%	19.6%	10.5%	14.3%	NR
PLCO <sub>m2012</sub> > 1.5%	8.9%	17.9%	NR	NR	NR
Eligible individuals as proportion of cohort of all ever-smokers					
USPSTF criteria	20.1%	38.0%	18.8%	NR	24.4%
PLCO <sub>m2012</sub> > 1.5%	17.9%	34.6%	NR	NR	9–35%, depending on ethnicity*

BHAS = Busselton Healthy Ageing Study; NR = not reported; PLCO = Prostate, Lung, Colorectal and Ovarian Trial; USPSTF = United States Preventive Services Task Force. \*Extrapolated from figure 1 in reference 18. ♦

challenges.<sup>15,25</sup> No state or federal database contains the necessary data for directly identifying the estimated 10% of the age-defined population eligible for screening. Population-based recruitment strategies for lung cancer screening must therefore currently engage with and assess the eligibility of a broader population of ever-smokers, the majority of whom will be ineligible. The feasibility and unintentional impact of this recruitment approach is not known.

Smoking cessation in one lung cancer screening study reduced all-cause mortality by up to 43%.<sup>26</sup> Almost half of BHAS current smokers (42.9%; 103 of 240 current smokers) were ineligible for screening, regardless of the criteria used. Providing smoking cessation support without LDCT screening to ineligible current smokers who engage with a screening program may be a pragmatic way to combine primary prevention and early detection of lung cancer.

Of the BHAS ever-smokers aged 50–55 years, 4.4–8.8% had PLCO<sub>m2012</sub> lung cancer risk sufficient to potentially benefit from screening, depending on the definition of COPD used. This may represent an additional population at

risk who may benefit from lung cancer screening at a lower age threshold than currently recommended, although internationally there is little prospect of such a change.<sup>27</sup>

There are several potential limitations to this study. The upper age limit of our BHAS participants was 68 years, younger than the maximum recommended screening age of 80 years.<sup>3</sup> This study may therefore have underestimated the proportion of the population potentially eligible for screening, as PLCO<sub>m2012</sub> risk increases with age.<sup>21</sup> Ethnic background has an impact on lung cancer risk, and the BHAS participants were almost exclusively European in origin, which is not representative of the broader Australian population, although the Busselton population age distribution and smoking demographic can be broadly generalised to the wider Australian population. Other lung cancer risk prediction models are being used to select individuals for lung cancer screening studies, but it was not possible to assess them, as they include variables (such as asbestos exposure) not recorded by the BHAS.<sup>6</sup> The definition of COPD

in the PLCO<sub>m2012</sub> model is self-reported COPD, chronic bronchitis or emphysema, more inclusive than the definition used in the BHAS. We were unable to validate the PLCO<sub>m2012</sub> model, as 6-year lung cancer outcomes are not available.

In conclusion, we estimate that between 17.9% and 20.1% of all current or former smokers aged 55–68 years in the BHAS population would be eligible for lung cancer screening, depending on the criteria applied. Symptom-defined COPD may be an appropriate surrogate measure for spirometric testing for determining the presence of COPD as part of the assessment of lung cancer screening eligibility.

**Acknowledgements:** Funding was provided by the Western Australian Cancer and Palliative Care Network for a Clinical Cancer Fellowship for David Manners. We acknowledge the Western Australian Country Health Service – South West for core infrastructure support, Marianne Weber for assistance with the PLCO<sub>m2012</sub> model, and the residents of Busselton Shire for participating in the study. The baseline survey of the Busselton Healthy Ageing Study was funded by grants from the Office of Science and the Department of Health of the Government of Western Australia (grant numbers G05911, G06301), the City of Busselton, a bequest of the late Janet Elder, and private donors.

**Competing interests:** No relevant disclosures.

Received 12 Jan 2016, accepted 12 Apr 2016. ■

© 2016 AMPCo Pty Ltd. Produced with Elsevier B.V. All rights reserved.

- 1 Australian Institute of Health and Welfare, Cancer Australia. Lung cancer in Australia: an overview (Cat. No. CAN 58; Cancer Series No. 64). Canberra: AIHW, 2011. <http://www.aihw.gov.au/WorkArea/DownloadAsset.aspx?id=10737421095&libID=10737421094> (accessed Feb 2016).
- 2 Aberle DR, Adams AM, Berg CD, et al. Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med* 2011; 365: 395-409.
- 3 Moyer VA. Screening for lung cancer: US Preventive Services Task Force recommendation statement. *Ann Intern Med* 2014; 160: 330-338.
- 4 Kovalchik SA, Tammemägi M, Berg CD, et al. Targeting of low-dose CT screening according to the risk of lung-cancer death. *N Engl J Med* 2013; 369: 245-254.
- 5 Black WC, Gareen IF, Soneji SS, et al. Cost-effectiveness of CT screening in the national lung screening trial. *N Engl J Med* 2014; 371: 1793-1802.
- 6 Field JK, Duffy SW, Baldwin DR, et al. UK Lung Cancer RCT Pilot Screening Trial: baseline findings from the screening arm provide evidence for the potential implementation of lung cancer screening. *Thorax* 2015; 71: 161-170.
- 7 Kauczor HU, Bonomo L, Gaga M, et al. ESR/ERS white paper on lung cancer screening. *Eur Radiol* 2015; 25: 2519-2531.
- 8 Tammemägi MC, Katki HA, Hocking WG, et al. Selection criteria for lung-cancer screening. *N Engl J Med* 2013; 368: 728-736.
- 9 Tammemägi MC, Church TR, Hocking WG, et al. Evaluation of the lung cancer risks at which to screen ever- and never-smokers: screening rules applied to the PLCO and NLST cohorts. *PLoS Med* 2014; 11: e1001764.
- 10 Toelle BG, Xuan W, Bird TE, et al. Respiratory symptoms and illness in older Australians: the Burden of Obstructive Lung Disease (BOLD) study. *Med J Aust* 2013; 198: 144-148. <https://www.mja.com.au/journal/2013/198/3/respiratory-symptoms-and-illness-older-australians-burden-obstructive-lung>
- 11 Tammemägi MC, Lam SC, McWilliams AM, et al. Incremental value of pulmonary function and sputum DNA image cytometry in lung cancer risk prediction. *Cancer Prev Res (Phila)* 2011; 4: 552-561.
- 12 Young RP, Duan F, Chiles C, et al. Airflow limitation and histology shift in the National Lung Screening Trial. The NLST-ACRIN Cohort Substudy. *Am J Respir Crit Care Med* 2015; 192: 1060-1067.
- 13 Australian Bureau of Statistics. 4364.0. 55.001. Australian Health Survey: first results, 2011–12. Canberra: ABS, 2012. [http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/4364.0.55.001Main+Features12011-12?OpenDocument](http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/4364.0.55.001>Main+Features12011-12?OpenDocument) (accessed Feb 2016).
- 14 James A, Hunter M, Straker L, et al. Rationale, design and methods for a community-based study of clustering and cumulative effects of chronic disease processes and their effects on ageing: the Busselton healthy ageing study. *BMC Public Health* 2013; 13: 936.
- 15 Brims F, McWilliams A, Fong K. Lung cancer screening in Australia: progress or procrastination? *Med J Aust* 2016; 204: 4-5. <https://www.mja.com.au/journal/2016/204/1/lung-cancer-screening-australia-progress-or-procrastination>
- 16 Brims FJ, Murray CP, de Klerk N, et al. Ultra-low-dose chest computer tomography screening of an asbestos-exposed population in Western Australia. *Am J Respir Crit Care Med* 2015; 191: 113-116.
- 17 Marshall HM, Bowman RV, Ayres J, et al. Lung cancer screening feasibility in Australia. *Eur Respir J* 2015; 45: 1734-1737.
- 18 Department of Health, Standing Committee on Screening. Position statement. Lung cancer screening using low-dose computed tomography. 2015. <http://www.cancerscreening.gov.au/internet/screening/publishing.nsf/Content/lung-cancer-screening> (accessed Feb 2016).
- 19 Fiscella K, Winters P, Farah S, et al. Do lung cancer eligibility criteria align with risk among Blacks and Hispanics? *PLoS One* 2015; 10: e0143789.
- 20 Howard DH, Richards TB, Bach PB, et al. Comorbidities, smoking status, and life expectancy among individuals eligible for lung cancer screening. *Cancer* 2015; 121: 4341-4347.
- 21 Ma J, Ward EM, Smith R, et al. Annual number of lung cancer deaths potentially avertable by screening in the United States. *Cancer* 2013; 119: 1381-1385.
- 22 Tammemägi MC, Pinsky PF, Caporaso NE, et al. Lung cancer risk prediction: prostate, lung, colorectal and ovarian cancer screening trial models and validation. *J Natl Cancer Inst* 2011; 103: 1058-1068.
- 23 Oken MM, Hocking WG, Kvale PA, et al. Screening by chest radiograph and lung cancer mortality: the Prostate, Lung, Colorectal, and Ovarian (PLCO) Randomized Trial. *JAMA* 2011; 306: 1865-1873.
- 24 Scollo M, Winstanley MH. Tobacco in Australia: facts and issues. Fourth edition. Melbourne: Cancer Council Victoria, 2012. <http://www.tobaccoinaustralia.org.au> (accessed Feb 2016).
- 25 Australian Population Health Development Principal Committee, Screening Subcommittee. Population based screening framework. Canberra: Australian Health Ministers' Council, 2008. [http://www.cancerscreening.gov.au/internet/screening/publishing.nsf/Content/16AE0B0524753EE9CA257CEE000B5D7/\\$File/Population-based-screening-framework.PDF](http://www.cancerscreening.gov.au/internet/screening/publishing.nsf/Content/16AE0B0524753EE9CA257CEE000B5D7/$File/Population-based-screening-framework.PDF) (accessed Feb 2016).
- 26 Pastorino U, Boffi R, Machianò A, et al. Stopping smoking reduces mortality in low-dose computed tomography (LDCT) screening volunteers. *J Thorac Oncol* 2016; doi: <http://dx.doi.org/10.1016/j.jtho.2016.02.011> [Epub ahead of print].
- 27 Field JK, Devaraj A, Duffy SW, et al. CT screening for lung cancer: Is the evidence strong enough? *Lung Cancer* 2016; 91: 29-35. ■