

**Note:**

This is an accepted manuscript (pre-print version) of an article published in *Music Perception* in April 2018, available online at: <http://mp.ucpress.edu>

This paper is not the copy of record and may not exactly replicate the authoritative document published in the journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1525/MP/2018.35.4.454

You may download the published version directly from the journal (homepage: <http://mp.ucpress.edu>).

## Published citation:

Krause, A. E., Davidson, J. W., & North, A. C. (2018). Musical activity and well-being: A new quantitative measurement instrument. *Music Perception*, 35, 4, 454-474. doi:10.1525/MP/2018.35.4.454

Musical Activity and Well-being: A New Quantitative Measurement Instrument

Amanda E. Krause

Curtin University, Perth, Australia & The University of Melbourne, Parkville, Australia

Jane W. Davidson

The University of Melbourne, Parkville, Australia

Adrian C. North

Curtin University, Perth, Australia

### Abstract

A relationship between participation in musical activity and well-being has frequently been observed in recent research reports. Of these, some propose various well-being-related correlates of musical participation, but the varying samples and foci leave researchers without a reasoned appraisal of these correlates or a data-driven categorization of them. To address this lacuna, the current research reviewed existing literature, identifying 562 benefits of well-being perceived to be associated with musical participation. These items were used as the basis for developing a new quantitative measure to evaluate the perceived benefits of well-being arising from music participation. Principal axis factor analysis of data using this new, 36-item measure identified five discrete dimensions: mood and coping, esteem and worth, socialization, cognition, and self-actualization. The development of this well-being measure addresses a gap in the research and provides a tool for future research concerning musical participation.

*Received December 31, 2016, accepted November 14, 2017.*

**Key words:** well-being, music-making, qualitative meta-analysis, music, participation

Music has been associated with therapeutic, medical, and curative powers throughout history and across numerous disciplines (MacDonald, Kreutz, & Mitchell, 2012). Investigating the link between music, health, and well-being<sup>1</sup> has become a topical concern (MacDonald, 2013) as people search for a range of approaches to addressing quality of life from socio-emotional to medical perspectives. A recent surge in research attention shows there is a growing acceptance that participation in music, and the arts broadly, can benefit people's well-being and health (Clift, 2012a; Clift, Hancox, Staricoff, & Whitmore, 2008; MacDonald, 2013; MacDonald et al., 2012; Skingley et al., 2011). Thus, there has been a growth of musical activities, yet the espoused well-being and/or health benefits have not been categorized in any systematic or comprehensive manner. In a context where many consider the value of participating in the arts as self-evident and simply disregard the need for systematic investigation into the topic (Skingley et al. (2011), one particular challenge for researchers is to identify the specific benefits that diverse forms of musical engagement may afford. Anecdotal enthusiasm from participants regarding positive correlates of musical activity is evident in reports, but since much prior work is limited to single case-studies, the literature lacks objective evidence.

The goal of the current study was to provide an assessment of the existing literature, and, in turn, use this as foundational material to develop an explicit measure of the potential well-being correlates of musical participation. To address this aim, we used a three-stage research process. First, we summarize the results of a literature review concerning the possible well-being correlates of participation in music. Second, we present a scale to measure these, and third, an empirical study of this scale to test its viability.

### Literature Review

In line with the mini-review method (Griffiths, 2002), we took a systematic approach to review relevant studies (see similar approaches taken by, e.g., Schäfer, Sedlmeier, Städtler, & Huron, 2013; Skingley & Vella-Burrows, 2010). Internet keyword searches (e.g., combining “music,” “singing,” and “instruments” with “well-being,” “health,” and “quality of life”) identified pertinent literature for review. The following inclusion criteria were employed: full text availability in English; quantitative and qualitative empirical research studies or reviews; and work published since 1996, in order to focus on digitally accessible data. Additional publications, not originally identified in the searches, were included as a consequence of being referenced in the papers reviewed and otherwise meeting the inclusion criteria. This resulted in the identification of 202 published works (listed in the Appendix).

A previous systematic review of health and well-being correlates of singing (see Clift, Hancox, Staricoff, et al., 2008) had revealed that studies concerning music and well-being vary with regard to method, focus, and sample characteristics. This led the authors to conclude that a coherent quantitative synthesis was not possible (see Clift & Hancox, 2010, p. 80). In light of this finding, we propose that a critical assessment of identified works is possible when adopting a qualitative meta-analysis process (e.g., Finfgeld, 2003; Levine & Land, 2016; Timulak, 2009), because it allows an overview of studies with differing research questions and methods (Levine & Land, 2016; Timulak, 2009).

#### *Works Reporting Music and Well-Being Correlates*

The first author reviewed the 202 works, with the second author sampling 33% of the material to validate the selection process (agreement rate = 97%). Data concerning well-being correlates were interpreted broadly, such that while some of the studies listed statistical correlates (e.g., Pearson's  $r$  values, factor loadings) or experimental manipulations involving music and well-being, we also included other evidence concerning benefits, motivations, reasons, and/or perceived consequences/outcomes of music. Ninety-seven of the 202 works (48.0%) included data concerning a benefit, consequence, participation reason, and/or theme related to well-being. Most of these works that did report on music participation and well-being concerned making music (68 studies or 73.1%; see Table 1); and, when defining the type of music-making undertaken, more than half of the studies concerned singing (54.4%, see Table 1). Considering the studies that involved mixed types of music activity (i.e., singing alongside other music activities), it is clear that the present body of research is heavily skewed towards data concerning singing.

-Insert Table 1 about here-

### *Items for Scale Development*

Schäfer et al. (2013, p. 5) identified over 500 proposed functions of music, and aimed to create “the most comprehensive list yet of potential music-related functions,” via an “aggregate list of all the questions and statements encountered in the reviewed research.” In a process similar to that applied by Schäfer et al. (2013), we recorded any benefit, consequence, participation reason, and/or theme related to well-being that was found in the 202 studies. An inclusive approach was taken, such that each item mentioned was

treated as a distinct data point (Timulak, 2009). For example, questionnaire items were recorded individually, as were qualitative themes/quotes.

While the individual correlates identified in the different publications varied in number and phrasing, the works tended to cluster specific items into categories. Thus, in addition to a number of specific outcomes associated with music participation, several common groupings were also identified (with varying frequency). Similar to research concerning the uses and functions of music in everyday life (see, e.g., Schäfer et al., 2013), these dimensions included social, emotion, cognitive, and arousal-based benefits. Additionally, category labels pertaining to physical health, identity, self-improvement, quality of life, and music appreciation were also frequently used (Habron, Butterly, Gordon, & Roebuck, 2013).

This investigation revealed that many previous studies have highlighted positive relationships between music participation, health, and well-being by addressing social, emotional, cognitive, physical health, musicianship, spiritual, and life satisfaction benefits. Illustrative examples of social benefits include networking, socializing, and nurturing friendships (e.g., Adams, 2000; Bungay & Skingley, 2008; Cavitt, 2005; Coffman, 2008; Eley & Gorman, 2010; Hillman, 2002; Jutras, 2011; Lehmborg & Fung, 2010; Livesey, Morrison, Clift, & Camic, 2012; McQueen, Hallam, Creech, & Varvarigou, 2013; Pavlakou, 2009; Rohwer, 2012; Saarikallio, 2012). More broadly, the benefit of feeling connected to and involved in a community—which also includes producing a sense of community—has also been noted frequently (e.g., Adderley, Kennedy, & Berz, 2003; Bailey & Davidson, 2005; Cohen, 2007; Creech et al., 2013; Dingle, Brander, Ballantyne, & Baker, 2012; Gembris, 2012; Jutras, 2011; Kennedy, 2002; Lally, 2009; Livesey et al., 2012; Michalos, 2005; von Lob, Camic, & Clift, 2010).

Emotional benefits repeatedly highlighted in previous research include the use of music for mood regulation (e.g., Judd & Pooley, 2014; Laukka, 2007; Livesey et al., 2012; Schäfer et al., 2013) and producing positive emotions such as feeling uplifted (e.g., Bailey & Davidson, 2005; Bungay & Skingley, 2008; Coffman, 2008; Hallam, Creech, Varvarigou, & McQueen, 2012a; Hays & Minichiello, 2005; Jacob, Guptill, & Sumsion, 2009; Judd & Pooley, 2014; Livesey et al., 2012; Sixsmith & Gibson, 2007; Tonneijck, Kinébanian, & Josephsson, 2008). Stress release and relaxation have also been commonly reported (e.g., Cavitt, 2005; Clift, Hancox, Morrison, et al., 2008; Jutras, 2006, 2011; Lehmberg & Fung, 2010; Michalos, 2005; Sheldon, 1998). Moreover, individuals have included both processing and expressing emotions as a perceived benefit to their music participation (e.g. Bailey & Davidson, 2005; Hays, 2005; Schäfer & Sedlmeier, 2009; Schäfer, Tipandjan, & Sedlmeier, 2012).

Cognitive benefits have included thinking about self-esteem, self-worth, and identity (e.g., Bailey & Davidson, 2002, 2003; Bonde, 2014; Cavitt, 2005; Clift, Hancox, Morrison, et al., 2008; Coffman, 2008; Forssén, 2007; Hallam, Creech, Varvarigou, & McQueen, 2012b; Kennedy, 2002; Lally, 2009; Livesey et al., 2012; Saarikallio, 2012; Silber, 2005). Other benefits to cognitive processing make reference to memory, concentration, and intellectual stimulation (e.g., Coffman, 2008; Creech et al., 2013; Gick, 2011; Hays, 2005; Southcott, 2009). Outcomes concerning creativity and imagination have also been frequently expressed (e.g., Bruhn, 2002; Coffman, 2008; Jutras, 2006, 2011; Kokotsaki & Hallam, 2011; Lehmberg & Fung, 2010; Michalos, 2005).

Physical health benefits have included general perceived outcomes such as feeling healthier and avoiding illness (e.g., Coffman, 2008; Gembris, 2012; Hillman, 2002; McQueen et al., 2013; Michalos, 2005; Rohwer, 2013). Additionally, perceived improvements to specific types of functioning have been noted, such as motor coordination (e.g., Bungay &



Skingley, 2008; Jutras, 2011), the immune system (e.g., Gick, 2011; Livesey et al., 2012), muscle tone (e.g., Lehmborg & Fung, 2010), flexibility and posture (e.g., Coffman, 2008; Lally, 2009; Livesey et al., 2012) and breathing (e.g., Clift & Hancox, 2001; Engen, 2005; Hallam et al., 2012b; Lord et al., 2010; Tonneijck et al., 2008).

Music-specific observations have focused on musicianship: most notably regarding learning, maintaining, and improving musical skills (e.g., Coffman & Adamek, 1999; Dingle et al., 2012; Gridley, Astbury, Sharples, & Aguirre, 2011; McQueen et al., 2013). More obviously, outcomes concerning learning and being exposed to new music have also been noted (e.g., Hays, 2005; Shansky, 2010) along with the opportunity to make music and perform (e.g., Hills & Argyle, 1998; Pavlakou, 2009; Pitts, 2004; Rohwer & Rohwer, 2009).

Some work has also considered perceived spiritual benefits of music involvement. Music participation can facilitate spiritual experiences and transcendent feelings (e.g., Beck, Cesario, Yousefi, & Enamoto, 2000; Bonde, 2014; Clift & Hancox, 2001; Kokotsaki & Hallam, 2007; Tonneijck et al., 2008). These perceived benefits may be religious in nature (e.g., Adams, 2000; Livesey et al., 2012; Rohwer, 2010) or more broadly relate to experiencing something “deep and meaningful” (e.g., Beck et al., 2000), meditation (Bonde, 2013), a peak experience (Cohen, 2007), or an aesthetic experience of beauty (e.g., Laukka, 2007; Livesey et al., 2012; Sheldon, 1998).

Other reported benefits concern perceptions of improved life satisfaction and quality of life (e.g., Clift et al., 2010; Douglas, 2011; Gembris, 2012). Items referring to one’s quality of life, and well-being in particular, were not always labeled in the same way. In some cases, life satisfaction or quality of life was used (e.g., Clift, Hancox, Morrison, et al., 2008; Douglas, 2011; Gembris, 2008), while in other research phrasing included feelings of satisfaction or a satisfying experience (e.g., Kennedy, 2002; Lehmborg & Fung, 2010;

McQueen et al., 2013; Tonneijck et al., 2008). Moreover, well-being was sometimes referenced directly, albeit again in different ways, including mental well-being/health (e.g., Dabback, 2009; Hills & Argyle, 1998; Jutras, 2011), personal or overall well-being (e.g., Hays, 2005; Michalos, 2005) and general well-being (e.g., Lord et al., 2010), emotional well-being (e.g., Coffman, 2008; Michalos, 2005), and psychological well-being (e.g., Lehmborg & Fung, 2010; Tsugawa, 2009). In addition to these broader concepts, there is also evidence that musical involvement defines how one lives one's life (e.g., Hays, 2005; Murcia, Kreutz, Clift, & Bongard, 2010), provides an enriching experience (e.g., Tonneijck et al., 2008; Tsugawa, 2009), adds meaning/purpose to life (e.g., Coffman, 2008; Livesey et al., 2012; Pothoulaki, MacDonald, & Flowers, 2012; Sheldon, 1998; Southcott, 2009), and promotes feelings of vitality and rejuvenation (e.g., Creech et al., 2013; Forssén, 2007; Gembris, 2012; Pothoulaki et al., 2012; Varvarigou, Hallam, Creech, & McQueen, 2012). As Gembris (2012, p. 377) asserted, "benefits from music-making like vitality, happiness, and connectedness to other people are among the most important goals in human life ... [and] musical activities can make a significant contribution to the quality of life, regardless of musical genre or skills."

The recurrent appearance of a small number of items suggests that it may be possible to identify core concerns in the existing research and develop a means of measuring these all within a single instrument. However, it is important to remember that the research is limited in a number of important ways. For example, some of the relevant work was focused on one particular dimension or another (e.g., social or physical health), was conducted with a specific population (e.g., elderly, persons with dementia, professional musicians, etc.), and/or focused on a specific type of musical participation (e.g., singing, community band, etc.), such that the findings may not generalize to other possible correlates of other musical activities carried out by other people. Moreover, relevant work

has adopted a variety of methods, both quantitative and qualitative. As a result, the results are largely presented independently from other work (with Clift et al.'s, 2008, extensive systematic review of singing research is a notable exception), and often the studies make use of researcher-developed and defined categorizations. Additionally, it is important to note that potentially relevant music therapy literature may be missing as a result of how the targeted review was conducted.

Thus, while the musical correlates of well-being have received considerable recent research attention, there has been no empirical attempt to categorize these correlates. In fact, while some of the work made use of measures of various forms of possible positive outcomes (e.g., Ballantyne, Ballantyne, & Packer, 2014; Hallam et al., 2012b), these have tended to be ad hoc and extremely specific rather than comprehensive or empirically based.

Consequently, there is no comprehensive measure (or immediately obvious basis for one) that addresses the various perceived well-being benefits resulting from musical participation. Further, there has not been a large-scale study to consider the broad range of perceived benefits of music participation with singers and instrumentalists within the same research design. Therefore, the second part of the current research addressed this research gap. Specifically, the items identified in the existing research literature served as the starting point for development of a new comprehensive measure that addresses the various potential psychological correlates of music participation, thereby grounding the measure in that earlier work.

## Measure Creation

### *Item Identification and Reduction*

In accordance with Schäfer et al. (2013), the first step towards developing a measure involved assembling a list of the reported correlates of musical participation addressed in previous work. As a result of the literature review above, a total of 2,075 items were identified from the 97 works reviewed that provided correlate data. Via discussion with an independent impartial judge, the first author reviewed this master list in order to eliminate redundant and duplicate items. Likewise, similar and analogous items were combined into single items. After this elimination stage, the list contained a total of 562 distinct items.

With such a large number of items, another reduction stage was needed in order to create a manageable measure for pilot testing. Many of the items came from qualitative data, and because there is no single accepted approach to analyze large qualitative datasets (Namey, Guest, Thairu, & Johnson, 2008; Stanton, Sinar, Balzer, & Smith, 2002), a panel of experts was employed (as proposed by, Morey, 2003) to consider the face and content validity of the items (Dunn, Bouffard, & Rogers, 1999; Morey, 2003; Stanton et al., 2002). The panel consisted of four well-qualified judges who were familiar with the subject matter (Dunn et al., 1999; Morey, 2003): specifically, the panel included a practicing music therapist (with extensive observation experience), a psychology of music researcher (who has conducted extensive research in singing and well-being, publishing peer-reviewed papers in this field), an ethnomusicologist (with extensive participant observation and other ethnomusicology expertise), and a musicologist (possessing performance and critical music skills).

The judges were provided with clear and precise directions (Hardesty & Bearden, 2004) concerning their task. In particular, they were provided with information concerning the purpose of the new measure (Delgado-Rico, Carretero-Dios, & Ruch, 2012; Morey, 2003) as follows:

The purpose of this scale is to have participants report on the well-being benefits they perceive as resulting from their voluntary participation in musical activities. Importantly, these perceived benefits are considered to be applicable to any individual participant involved in a musical activity, regardless of: (a) musical training/experience level, (b) musical activity (i.e., a vocal /instrumental /therapy group, etc.), (c) demographic characteristics (age, gender, geographic location), and (d) frequency/length of participation.

The panel members were asked to “read each item and rate how well you think it addresses the definition of the measure.” They were prompted to answer the question, “how representative of the definition (i.e., how relevant) for the scale is this item?” using a five-point Likert scale where 1 represented *Not at all representative/relevant (poor match)* and 5 represented *Extremely representative/relevant (Excellent match)*. Each judge independently rated the list of 562 items. The four judges’ ratings were compiled, and the mean rating was computed for each item. In order to be retained in the measure, an item was required to produce a mean rating of at least 3.75, such that at least three of the four judges needed to rate the item as a 4 or 5 on the five-point scale. In turn, 40 distinct items were retained for the measure that was tested in part 3 of the present research (see Table 2).

### Questionnaire Development

The questionnaire phase of the research had two goals, namely: 1) to identify how people perceived the reduced list of items as grouping together, and 2) to determine which perceived correlates of music participation were experienced most commonly by the respondents, and whether this varied as a function of the type of music participation.

Establishing this information allows the questionnaire to be used by future research as an overarching measure of the perceived benefits of music participation.

With regard to the first goal, the review of the literature had demonstrated that although there was some commonality to the items considered, there was no comprehensive or established structure concerning these already in the literature. Thus, while commonalities were found, our research question was: what are the factors or dimensions of the perceived benefits of music participation for well-being?

Regarding the second goal, we were interested in whether people experience certain factors more or less frequently than others. Moreover, does the experience of these factors vary as a function of participant gender and/or the type of music participation? Previous research has indicated observable gender roles in musical behavior, preferences, and beliefs (see review by Dibben, 2002). For instance, research has identified gender stereotyping of musical instruments and singing (e.g., Conway, 2000; Harrison, 2007; O'Neill, 1997). Consequently, it is possible that individuals' subjective experiences of voluntary musical participation might differ by gender and type of participation.

### *Method*

The measure included the 40 items retained from the prior panel stage. All items were phrased so that they addressed the individual's personal experience (e.g., "It gives me..." and "I do not..."). As prior research has criticized use of only positively worded items (e.g., Clift & Hancox, 2010), half of the items were quasi-randomly selected and reworded, such that they required reverse coding. Instructions asked participants (selected as those who actively engaged in music participation) to "consider the benefits to your well-being

that result from your participation in music” and consequently to rate how strongly they agreed with each stated item on a seven-point Likert scale (1 = *not at all*, 7 = *fully agree*).

In total, 207 individuals completed the 40-item questionnaire online. Participants ranged from 18 to 72 years old ( $M = 32.95$ ,  $Mdn = 30$ ,  $SD = 13.40$ ) and 66.70% were female. We specifically sought to obtain a sample of people who participated in music as vocalists and instrumentalists. While people may be involved in multiple musical activities, 65.20% identified themselves as primarily instrumentalists, while 34.80% identified themselves as primarily singers. Participation in the study was voluntary: individuals were recruited through the researchers’ extensive networks of Australian musicians (including professional and amateur performers, music therapists, and music teachers).

### *Results*

To investigate the underlying structure of the measure, the 40 items were entered into a Principal Axis Factor analysis with promax rotation. The KMO value was .905, Bartlett’s Test was significant ( $p < .001$ ), all MSA values were above .74 (with most above .90), and all items demonstrated reasonable communality values. After visual inspection of the scree plot and evaluation of the eigenvalues and factor loadings, five meaningful factors were retained accounting for 50.90% of the total variance (see details in Table 2). Specifically, the first factor was labeled “mood and coping,” and includes 11 items concerning feeling uplifted, feeling better, and relieving stress/tension. The second factor, labeled “esteem and worth,” includes 11 items concerning feeling positive about life, a sense of belonging, and an increase in self-esteem. “Socializing,” factor 3, concerned perceived benefits such as making friends, bonding with other people, and connecting with a community (five items). Factor 4 concerned cognitive processes, and included five items

addressing thinking and processing, and so it was labeled “cognitive.” The fifth factor, “self-actualization” included four items reflecting self-improvement. Based on the results of the principal axis factor analysis, four of the forty items were removed, as they did not feature in the five-factor model. This results in a revised measure with 36 items that includes five subscales, labeled as per the factors.

To check the reliability of each factor, the items that loaded onto each were entered into separate reliability analyses. Each of the five factors demonstrated good reliability, based on resulting Cronbach alpha values between .75 and .90 (see Table 2). Considering the demonstrated reliability, each factor has the potential to be treated as an individual subscale measure, subject to further work to validate these factors and subscale measures (by, for example, establishing test-retest reliability and similar psychometric properties). Dimension scores can be interpreted as representing the degree to which individuals (or samples, when analyzed collectively) are experiencing the different categories of well-being benefit. Thus, the results of the separate dimensions have the potential to be applied in a variety of music participation contexts, including community-based, therapeutic, and music education activities.

**-Table 2 here –**

*Relative importance.* In order to consider the relative importance of the identified factors for people participating in music activities, mean scores were produced for the participants. The ratings for the set of items principally loading onto each factor were averaged (Schäfer et al., 2013). A one-way repeated measures ANOVA compared the subscale means. The ANOVA was significant,  $F(3, 715.86) = 5.37, p = .001, \eta_p^2 = .03$ , and pairwise comparisons indicated that the esteem and worth benefits ( $M = 5.91, SD = .07$ )



were experienced significantly more than those concerning socializing ( $M = 5.65, SD = .07$ ) or cognitive benefits ( $M = 5.71, SD = .07$ ). Additionally, the self-actualization factor ( $M = 5.89, SD = .08$ ) gave rise to a significantly higher rating than did both the socializing and cognitive factors.

*Perceived benefit responses.* A MANOVA was used to examine the effect of type of music participation and gender on the perceived benefit factor scores. Multivariate outliers were removed as the Mahalanobis distance exceeded the critical  $\chi^2$  value (for  $df = 5$  at  $\alpha = .001$ ), although the remaining assumptions were met. The interaction between participation type and gender was nonsignificant,  $F(5, 190) = 1.17, p = .33, \eta_p^2 = .03$ . Additionally, the main effect for gender was nonsignificant,  $F(5, 190) = 1.29, p = .27, \eta_p^2 = .03$ , and the main effect for participation type was also nonsignificant,  $F(5, 190) = 0.85, p = .52, \eta_p^2 = .02$ . These nonsignificant results indicate the absence of any meaningful difference in the extent of perceived benefits based on whether the participants were instrumentalists or singers or as a function of gender.

### General Discussion

The present research had two aims, namely to synthesize the current state of literature regarding music participation and well-being, and to produce a questionnaire measure of the subjective experience of the potential well-being benefits. First, we reviewed literature concerning the correlates of music participation that reflect various manifestations of well-being. From this literature survey, more than 500 potential benefits were identified. Following the reduction of the list via redundancy checks and evaluation by a panel of experts, a 40-item measure was established. Using questionnaire data concerning the frequency with which the potential benefits were experienced, a principal axis factor

analysis identified five dimensions of perceived well-being benefits pertaining to 36 items, encapsulated by five dimensions: mood and coping, esteem and worth, social, cognitive, and self-actualization.

Consequently, we have provided a comprehensive categorization of the perceived well-being benefits that may arise from music participation. Components inherent to the consideration of quality of life are present in more than one of the scale dimensions: the perceived benefits to one's well-being arising from music participation appear to be related to the notion of living a full, enriched life. In light of the current debate surrounding the imprecise definitions of health and well-being, these findings align with the view that well-being has an aspect of social capital embedded within it, for which a medical model of health does not account. The findings, then, are consistent with the WHO's broader, encompassing definition of health in that it is more than simply the narrowed perception of physical health. Moreover, this broader conceptualization of health/well-being is perhaps related to the literature matching the review criteria. For instance, clinical music therapy research conducted in hospitals may not have met the inclusion criteria for the present review, whereas community music therapy with a more social, community focus would have been more likely to meet the criteria (and thus represented). Additionally, future work could further interrogate the small amount of research regarding education/professional attempts to link music education to promoting long-term health and well-being.

Importantly, these dimensions (and this measure) specifically address *perceived* benefits rather than *actual* benefits. While this measure is effective in terms of measuring a person's subjective experience, it does not address, for example, actual physical health improvements. This explains why no distinct physical health-focused dimension is present. Moreover, as Gembris (2012, p. 375) stated, "usually, we do not pursue music for health

reasons as is the case with sports and gymnastics, but for its own sake.” So, while music participation can result in benefits to one’s physical health, this may not always be readily appreciated by participants, or necessarily valued highly as an outcome of musical activity. It also remains possible that any impact of music participation on physical health is associated more closely with certain types of activity or musicians than others: the present data cannot speak to this.

The experience of any of the five dimensions is not mutually exclusive. Any one individual may experience any or all of the five identified dimensions; and experiencing one dimension of well-being does not preclude experiencing another. It is likely that the experience of the perceived benefits overlaps, so that individuals participating in musical activity may not necessarily distinguish any particular correlate as different from a generalized sense of improved well-being.

These results have clear implications for the development of research that aims to assess well-being in the context of musical activity. The measure developed here can be used widely, as it is not specific to any particular type of music activity or participant. Moreover, it provides an overarching measure of perceived benefit, and also has the potential (pending appropriate validation work) to represent separate measures on each of five particular dimensions which cover the range of benefits as described in prior research. These findings and this tool, in turn, have implications for the facilitators of music activities, as well as to psychologists, music therapists, and medical professionals who make use of music in their practice and research. By defining the relationships between the types of benefits experienced via musical activity, the present findings help facilitators to make choices and, where necessary, changes to their practices to better promote desired outcomes. By understanding the different dimensions of well-being, practitioners may be

better able to elicit them, and could use the measure developed here to test the efficacy of this.

The present findings are not without limitations though. With the burgeoning interest in music and health, more work is being published that may address well-being correlates. It falls on future research to clearly delineate their methodologies and foci to better define their research questions and definitions of health and well-being. While we attempted to recruit those with a range of musical specialisms during the data collection phase of the present research, it should be recognized that music participation encompasses a large variety of sometimes less formal activities. Future research could involve additional (and more specific) types of music participation activities: in doing so, the perceived benefits of a wider range of different music activities could be compared. Such information, for instance, could be used to better tailor music opportunities to interested individuals. Moreover, cross-cultural work is needed to better understand any positive perceived correlates of music participation outside the countries represented in the present research, and the measure developed here could even be used to consider a broader range of arts and leisure activities, beyond music. Future research could make use of this tool to also understand how participants' motivations align with the benefits that they believe will arise from musical activity. For instance, such research may consider whether people's initial motivating reasons for participating map onto their perceived benefits, as well as whether both of these change via sustained involvement.

### Conclusion

In summary, there is a growing interest worldwide in the connection between music, health, and well-being (Clift, 2012b), which has been accompanied by a recent increase in

research attention (MacDonald, 2013; MacDonald et al., 2012; Skingley et al., 2011). By performing a review of the prior research concerning music and well-being, the present work identified more than five hundred items related to well-being that have been investigated for their possible correlation with musical activity. From this large pool of potential benefits, we created a 36-item measure that minimizes redundancy. Factor analysis of participants' responses to these indicates that we can categorize the perceived well-being correlates of musical activity in terms of mood and coping, esteem and worth, socialization, cognition, and self-actualization. Importantly, the development of this measure addresses a gap in the research, and we hope that the use of this new tool will vivify future research concerning musical participation.

## Footnote

<sup>1</sup> An issue that plagues this area of research is the debate concerning the definitions of health and well-being (Clift & Hancox, 2010; MacDonald et al., 2012; Saarikallio, 2012; Skingley, Bungay, & Clift, 2011). The World Health Organization (WHO) provides the often-used definition of health as “a state of complete physical, mental and social well-being and not merely the absence of disease and infirmity” (WHO, 1946). This definition of health makes it clear that health is a multidimensional phenomenon—one that includes “biochemical, psycho-motor, cognitive, emotional, social, cultural (moral, spiritual, aesthetic)” aspects (Ansdell & Denora, 2012). Moreover, “it is possible to enjoy ‘good health’ according to some of these measures, while being ‘ill’ according to others. In short, there is no ‘one form’ of health” (Ansdell & Denora, 2012, p. 106). Our use of “well-being” in this paper aligns with the WHO’s definition of health, as opposed to a more constricted medical model.

## Author Note

This work was carried out with the financial assistance of the Australian Research Council (DP140102679), which we acknowledge with gratitude.

*Correspondence concerning this article should be addressed to Amanda E. Krause, Melbourne Conservatorium of Music, The University of Melbourne, Melbourne, 3010 Australia, Australia. E-mail: [Amanda.Krause@unimelb.edu.au](mailto:Amanda.Krause@unimelb.edu.au)*

## References

- Adams, K. (2000). How can I keep from singing: A lifetime of sweet singing in the choir. In B. A. Roberts & A. Rose (Eds.), *Sharing the voices: The phenomenon of singing 2. Proceedings of the International Symposium*. St. John's, Newfoundland, Canada: Memorial University of Newfoundland.
- Adderley, C., Kennedy, M., & Berz, W. (2003). "A home away from home": The world of the music classroom. *Journal of Research in Music Education*, 51, 190–205.
- Ansdell, G., & Denora, T. (2012). Musical flourishing: Community music therapy, controversy, and the cultivation of wellbeing. In R. A. R. MacDonald, G. Kreutz, & L. A. Mitchell (Eds.), *Music, health, and wellbeing* (pp. 97–112). Oxford, UK: Oxford University Press.
- Bailey, B. A., & Davidson, J. W. (2002). Adaptive characteristics of group singing: perceptions from members of a choir for homeless men. *Musicae Scientiae*, 6(2), 221–256. DOI: 10.1177/102986490200600206
- Bailey, B. A., & Davidson, J. W. (2003). Amateur group singing as a therapeutic instrument. *Nordic Journal of Music Therapy*, 12(1), 18–32. DOI: 10.1080/08098130309478070
- Bailey, B. A., & Davidson, J. W. (2005). Effects of group singing and performance for marginalized and middle-class singers. *Psychology of Music*, 33(3), 269–303. DOI: 10.1177/0305735605053734
- Ballantyne, J., Ballantyne, R., & Packer, J. (2014). Designing and managing music festival experiences to enhance attendees' psychological and social benefits. *Musicae Scientiae*, 18(1), 65–83. DOI: 10.1177/1029864913511845



- Beck, R. J., Cesario, T. C., Yousefi, A., & Enamoto, H. (2000). Choral singing, performance perception, and immune system changes in salivary immunoglobulin A and cortisol. *Music Perception, 18*, 87–106.
- Bonde, L. O. (2013). The musical identities of Danish music therapy students: A study based on musical autobiographies. In L. O. Bonde, E. Ruud, M. S. Skånland, & G. Trondalen (Eds.), *Musical life stories. Narratives on health musicking* (pp. 307–328). Oslo, Norway: Centre for Music and Health, the Norwegian Academy of Music.
- Bonde, L. O. (2014). Music and health promotion - in the life of music therapy and music psychology researchers: A pilot study. *Voices: A World Forum for Music Therapy, 14*(1), n.p. DOI: 10.15845/voices.v14i1.740
- Bruhn, H. (2002). Musical development of elderly people. *Psychomusicology, 18*, 59–75.
- Bungay, H., & Skingley, A. (2008). *The silver song club project: Summary of a formative evaluation*. Canterbury, UK: Canterbury Christ Church University.
- Cavitt, M. E. (2005). Factors influencing participation in community bands. *Journal of Band Research, 41*(1), 42–57.
- Clift, S. (2012a). Singing, wellbeing, and health. In R. A. R. MacDonald, G. Kreutz, & L. A. Mitchell (Eds.), *Music, health, and wellbeing* (pp. 113-124). Oxford, UK: Oxford University Press.
- Clift, S. (2012b). Creative arts as a public health resource: moving from practice-based research to evidence-based practice. *Perspectives in Public Health, 132*(3), 120–127. DOI: 10.1177/1757913912442269
- Clift, S., & Hancox, G. (2001). The perceived benefits of singing: Findings from preliminary surveys of a university college choral society. *The Journal of the Royal Society for the Promotion of Health, 121*(4), 248–256.

- Clift, S., & Hancox, G. (2010). The significance of choral singing for sustaining psychological wellbeing: Findings from a survey of choristers in England, Australia and Germany. *Music Performance Research, 3*(1), 79–96.
- Clift, S., Hancox, G., Morrison, I., Hess, B., Kreutz, G., & Stewart, D. (2010). Choral singing and psychological wellbeing: Quantitative and qualitative findings from English choirs in a cross-national survey. *Journal of Applied Arts and Health, 1*(1), 19–34.  
DOI: 10.1386/jaah.1.1.19/1
- Clift, S., Hancox, G., Morrison, I., Hess, B., Stewart, D., & Kreutz, G. (2008). *Choral singing, wellbeing and health: Summary of findings from a cross-national survey*. Canterbury, UK: Canterbury Christ Church University.
- Clift, S., Hancox, G., Staricoff, R., & Whitmore, C. (2008). *Singing and health: Summary of a systematic mapping and review of non-clinical research*. Canterbury, UK: Canterbury Christ Church University.
- Coffman, D. D. (2008). Survey of New Horizons International Music Association musicians. *International Journal of Community Music, 1*(3), 375–390. DOI: 10.1386/ijcm.1.3.375/1
- Coffman, D. D., & Adamek, M. S. (1999). The contributions of wind band participation to quality of life of senior adults. *Music Therapy Perspectives, 17*, 27–31.
- Cohen, M. L. (2007). Explorations of inmate and volunteer choral experiences in a prison-based choir. *Australian Journal of Music Education, 1*, 61–72.
- Conway, C. (2000). Gender and musical instrument choice: A phenomenological investigation. *Bulletin of the Council for Research in Music Education, 146*, 1–17.

- Creech, A., Hallam, S., Gaunt, H., McQueen, H., Pincas, A., & Varvarigou, M. (2013). The power of music in the lives of older adults. *Research Studies in Music Education, 35*(1), 87–102. DOI: 10.1177/1321103X13478862
- Dabback, W. M. (2009). Identity formation through participation in the Rochester New Horizons band programme. *International Journal of Community Music, 1*(2), 267–286. DOI: 10.1386/ijcm.1.2.267/1
- Delgado-Rico, E., Carretero-Dios, H., & Ruch, W. (2012). Content validity evidences in test development: An applied perspective. *International Journal of Clinical and Health Psychology, 12*(3), 449–460.
- Dibben, N. (2002). Gender identity and music. In R. A. R. MacDonald, D. J. Hargreaves, & D. Miell (Eds.), *Musical identities* (pp. 117–133). Oxford, UK: Oxford University Press.
- Dingle, G. A., Brander, C., Ballantyne, J., & Baker, F. A. (2012). 'To be heard': The social and mental health benefits of choir singing for disadvantaged adults. *Psychology of Music, 41*(4), 405–421. DOI: 10.1177/0305735611430081
- Douglas, K. A. (2011). *A descriptive analysis of the psychological needs of adults participating in music ensembles: A survey of the New Horizons International Music Association ensemble participants* (Unpublished doctoral dissertation). Available from ProQuest Dissertations and Theses database. AAT3473456.
- Dunn, J. G. H., Bouffard, M., & Rogers, W. T. (1999). Assessing item content-relevance in sport psychology scale-construction research: Issues and recommendations. *Measurement in Physical Education and Exercise Science, 3*(1), 15–36. DOI: 10.1207/s15327841mpee0301\_2

- Eley, R., & Gorman, D. (2010). Didgeridoo playing and singing to support asthma management in Aboriginal Australians. *The Journal of Rural Health, 26*, 100–104. DOI: 10.1111/j.1748-0361.2009.00256.x
- Engen, R. L. (2005). The singer's breath: Implications for treatment of persons with emphysema. *Journal of Music Therapy, 42*(1), 20–48.
- Finfgeld, D. L. (2003). Metasynthesis: The state of the art—so far. *Qualitative Health Research, 13*(7), 893–904. DOI: 10.1177/1049732303253462
- Forssén, A. S. K. (2007). Humour, beauty, and culture as personal health resources: Experiences of elderly swedish women. *Scandinavian Journal of Public Health, 35*(3), 228-234. DOI: 10.1080/14034940601160680
- Gembris, H. (2008). Musical activities in the third age: An empirical study with amateur musicians. In A. Daubney, E. Longhi, A. Lamont, & D. J. Hargreaves (Eds.), *Second European Conference on Developmental Psychology of Music* (pp. 103–108). London, UK: Roehampton University.
- Gembris, H. (2012). Music-making as a lifelong development and resource for health. In R. A. R. MacDonald, G. Kreutz, & L. Mitchell (Eds.), *Music, health, and wellbeing* (pp. 367-382). Oxford, UK: Oxford University Press.
- Gick, M. L. (2011). Singing, health and well being: A health psychologist's review. *Psychomusicology: Music, Mind and Brain, 21*(1&2), 176–207. DOI: 10.1037/h0094011
- Gridley, H., Astbury, J., Sharples, J., & Aguirre, C. (2011). *Benefits of group singing for community mental health and wellbeing: Survey and literature review*. Carlton, VIC: Victorian Health Promotion Foundation.

- Griffiths, P. (2002). Evidence informing practice: Introducing the mini-review. *British Journal of Community Nursing*, 7(1), 38–39.
- Habron, J., Butterly, F., Gordon, I., & Roebuck, A. (2013). Being well, being musical: Music composition as a resource and occupation for older people. *British Journal of Occupational Therapy*, 76(7), 308–316. DOI: 10.4276/030802213X13729279114933
- Hallam, S., Creech, A., Varvarigou, M., & McQueen, H. (2012a). The characteristics of older people who engage in community music making, their reasons for participation and the barriers they face. *Journal of Adult and Continuing Education*, 18(2), 21–43. DOI: 10.7227/JACE.18.2.3
- Hallam, S., Creech, A., Varvarigou, M., & McQueen, H. (2012b). Perceived benefits of active engagement with making music in community settings. *International Journal of Community Music*, 5(2), 155–174.
- Hardesty, D. M., & Bearden, W. O. (2004). The use of expert judges in scale development: Implications for improving face validity of measures of unobservable constructs. *Journal of Business Research*, 57, 98–107. DOI: 10.1016/S0148-2963(01)00295-8
- Harrison, S. D. (2007). A perennial problem in gendered participation in music: What's happening to the boys? *British Journal of Music Education*, 24(3), 267–280. DOI: 10.1017/S0265051707007577
- Hays, T. (2005). Well-being in later life through music. *Australasian Journal on Ageing*, 24(1), 28–32.
- Hays, T., & Minichiello, V. (2005). The meaning of music in the lives of older people: A qualitative study. *Psychology of Music*, 33(4), 437–451. DOI: 10.1177/0305735605056160

- Hillman, S. (2002). Participatory singing for older people: A perception of benefit. *Health Education, 102*(4), 163–171. DOI: 10.1108/09654280210434237
- Hills, P., & Argyle, M. (1998). Musical and religious experiences and their relationship to happiness. *Personality and Individual Differences, 25*, 91–102.
- Jacob, C., Guptill, C., & Sumsion, T. (2009). Motivation for continuing involvement in a leisure-based choir: The lived experiences of university choir members. *Journal of Occupational Science, 16*(3), 187–193. DOI: 10.1080/14427591.2009.9686661
- Judd, M., & Pooley, J. A. (2014). The psychological benefits of participating in group singing for members of the general public. *Psychology of Music, 42*(2), 269–283. DOI: 10.1177/0305735612471237
- Jutras, P. J. (2006). The benefits of adult piano study as self-reported by selected adult piano students. *Journal of Research in Music Education, 54*(2), 97–110.
- Jutras, P. J. (2011). The benefits of new horizons band participation as self-reported by selected New Horizons band members. *Bulletin of the Council for Research in Music Education, 187*, 65–84.
- Kennedy, M. A. (2002). It's cool because we like to sing: Junior high school boys' experience of choral music as an elective. *Research Studies in Music Education, 18*, 24–35.
- Kokotsaki, D., & Hallam, S. (2007). Higher education music students' perceptions of the benefits of participative music making. *Music Education Research, 9*(1), 93–109. DOI: 10.1080/14613800601127577
- Kokotsaki, D., & Hallam, S. (2011). The perceived benefits of participative music making for non-music university students: a comparison with music students. *Music Education Research, 13*(2), 149–172. DOI: 10.1080/14613808.2011.577768

- Lally, E. (2009). 'The power to heal us with a smile and a song': Senior well-being, music-based participatory arts and the value of qualitative evidence. *Journal of Arts and Communities, 1*(1), 25–44. DOI: 10.1386/jaac.1.1.25/1
- Laukka, P. (2007). Uses of music and psychological well-being among the elderly. *Journal of Happiness Studies, 8*, 215–341. DOI: 10.1007/s10902-006-9024-3
- Lehmberg, L. J., & Fung, C. V. (2010). Benefits of music participation for senior citizens: A review of the literature. *Music Education Research International, 4*, 19–30.
- Levine, B., & Land, H. M. (2016). A meta-synthesis of qualitative findings about dance/movement therapy for individuals with trauma. *Qualitative Health Research, 26*(3), 330–344. DOI: 10.1177/1049732315589920
- Livesey, L., Morrison, I., Clift, S., & Camic, P. (2012). Benefits of choral singing for social and mental wellbeing: qualitative findings from a cross-national survey of choir members. *Journal Of Public Mental Health, 11*(1), 10–26. DOI: 10.1108/17465721211207275
- Lord, V. M., Cave, P., Hume, V. J., Flude, E. J., Evans, A., Kelly, J. L., et al. (2010). Singing teaching as a therapy for chronic respiratory disease - A randomised controlled trial and qualitative evaluation. *BMC Pulmonary Medicine, 10*, article 41.
- MacDonald, R. A. R. (2013). Music, health, and well-being: A review. *International Journal of Qualitative Studies on Health and Well-being, 8*, 20635. DOI: 10.3402/qhw.v8i0.20635
- MacDonald, R. A. R., Kreutz, G., & Mitchell, L. A. (2012). What is music, health, and wellbeing and why is it important? In R. A. R. MacDonald, G. Kreutz, & L. A. Mitchell (Eds.), *Music, health, and wellbeing* (3-11). Oxford, UK: Oxford University Press.

- McQueen, H., Hallam, S., Creech, A., & Varvarigou, M. (2013). A philosophical perspective on leading music activities for the over 50s. *International Journal of Lifelong Education, 32*(3), 353–377. DOI: 10.1080/02601370.2012.738432
- Michalos, A. C. (2005). Arts and the quality of life: An exploratory study. *Social Indicators Research, 71*, 11–59. DOI: 10.1007/s11205-004-8013-3
- Morey, L. C. (2003). Measuring personality and psychopathology. In I. B. Weiner (Ed.), *Handbook of psychology* (Vol. 3, pp. 377–405). Hoboken, NJ: John Wiley & Sons, Inc.
- Murcia, C. Q., Kreutz, G., Clift, S., & Bongard, S. (2010). Shall we dance? An exploration of the perceived benefits of dancing on well-being. *Arts and Health: An International Journal for Research, Policy and Practice, 2*(2), 149–163. DOI: 10.1080/17533010903488582
- Namey, E., Guest, G., Thairu, L., & Johnson, L. (2008). Data reduction techniques for large qualitative data sets. In G. Guest & K. M. MacQueen (Eds.), *Handbook for team-based qualitative research* (pp. 137–161). Lanham, MD: Altamira Press.
- O’Neill, S. A. (1997). Gender and music. In D. J. Hargreaves & A. C. North (Eds.), *The social psychology of music* (pp. 46–63). Oxford, UK: Oxford University Press.
- Pavlakou, M. (2009). Benefits of group singing for people with eating disorders: Preliminary findings from a non-clinical study. *Approaches: Music Therapy and Special Music Education, 1*(1), 30–48.
- Pitts, S. E. (2004). 'Everybody wants to be Pavarotti': The experience of music for performers and audience at a Gilbert and Sullivan festival. *Journal of the Royal Musical Association, 129*(1), 143–160.



- Pothoulaki, M., MacDonald, R. A. R., & Flowers, P. (2012). The use of music in chronic illness: Evidence and arguments. In R. A. R. MacDonald, G. Kreutz, & L. A. Mitchell (Eds.), *Music, health, and wellbeing* (pp. 239–256). Oxford, UK: Oxford University Press.
- Rohwer, D. (2010). Church musicians' participation perceptions: Applications to community music. *Research and Issues in Music Education*, 8(1), n.p. Retrieved from <http://files.eric.ed.gov/fulltext/EJ894767.pdf>.
- Rohwer, D. (2012). How participants envision community music in Welsh men's choirs. *Research & Issues in Music Education*, 10(1), n.p. Retrieved from <http://www.stthomas.edu/rimeonline/vol10/rohwer.htm>.
- Rohwer, D. (2013). Making music as an adult: What do the spouses think? *Texas Music Education Research*, 40-46.
- Rohwer, D., & Rohwer, M. (2009). A content analysis of choral students' participation perceptions: Implications for lifelong learning. *International Journal of Community Music*, 2, 255–262. DOI: 10.1386/ijcm.2.2&3.255/1
- Saarikallio, S. (2012). Cross-cultural approaches to music and health. In R. A. R. MacDonald, G. Kreutz, & L. Mitchell (Eds.), *Music, health, and wellbeing* (pp. 477–490). Oxford, UK: Oxford University Press.
- Schäfer, T., & Sedlmeier, P. (2009). From the functions of music to music preference. *Psychology of Music*, 37(3), 279–300. DOI: 10.1177/0305735608097247
- Schäfer, T., Sedlmeier, P., Städtler, C., & Huron, D. (2013). The psychological functions of music listening. *Frontiers in Psychology*, 4, article 311. DOI: 10.3389/fpsyg.2013.00511

- Schäfer, T., Tipandjan, A., & Sedlmeier, P. (2012). The functions of music and their relationship to music preference in India and Germany. *International Journal of Psychology, 47*(5), 370–380. DOI: 10.1080/00207594.2012.688133
- Shansky, C. (2010). Adult motivations in community orchestra participation: A pilot case study of the Bergen philharmonic orchestra (New Jersey). *Research and Issues in Music Education, 8*(1), n.p. Retrieved from <http://www.stthomas.edu/rimeonline/vol8/shansky.htm>.
- Sheldon, D. A. (1998). Participation in community and company bands in Japan. *Applications of Research in Music Education, 17*(1), 21–24. DOI: 10.1177/875512339801700105
- Silber, L. (2005). Bars behind bars: The impact of a women's prison choir on social harmony. *Music Education Research, 7*(2), 251–271. DOI: 10.1080/14613800500169811
- Sixsmith, A., & Gibson, G. (2007). Music and the wellbeing of people with dementia. *Ageing and Society, 27*, 127–145. DOI: 10.1017/S0144686X06005228
- Skingley, A., Bungay, H., & Clift, S. (2011). Researching participatory arts, well-being and health: some methodological issues. *Journal of Arts and Communities, 3*(1), 73–87. DOI: 10.1386/jaac.3.1.73\_1
- Skingley, A., & Vella-Burrows, T. (2010). Therapeutic effects of music and singing for older people. *Nursing Standard, 24*(19), 35–41.
- Southcott, J. E. (2009). 'And as I go, I love to sing': the Happy Wanderers, music and positive aging. *International Journal of Community Music, 2*(2&3), 143–156. DOI: 10.1386/ijcm.2.2&3.143/1
- Stanton, J. M., Sinar, E. F., Balzer, W. K., & Smith, P. C. (2002). Issues and strategies for reducing the length of self-report scales. *Personnel Psychology, 55*, 167–194.

- Timulak, L. (2009). Meta-analysis of qualitative studies: A tool for reviewing qualitative research findings in psychotherapy. *Psychotherapy Research, 19*(4-5), 591–600. DOI: 10.1080/10503300802477989
- Tonneijck, H. I. M., Kinébanian, A., & Josephsson, S. (2008). An exploration of choir singing: Achieving wholeness through challenge. *Journal of Occupational Science, 15*(3), 173–180. DOI: 10.1080/14427591.2008.9686627
- Tsugawa, S. (2009). *Senior adult music learning, motivation, and meaning construction in two New Horizons ensembles* (Unpublished doctoral dissertation).
- Varvarigou, M., Hallam, S., Creech, A., & McQueen, H. (2012). Benefits experienced by older people who participated in group music-making activities. *Journal of Applied Arts and Health, 3*(2), 183–198.
- von Lob, G., Camic, P., & Clift, S. (2010). The use of singing in a group as a response to adverse life events. *International Journal of Mental Health Promotion, 12*(3), 45–53. DOI: 10.1080/14623730.2010.9721818
- WHO. (1946). *Preamble to the Constitution of the World Health Organization as adopted by the International Health Conference*, New York. Signed on July 22, 1946 by the representatives of 61 States (Official Records of the World Health Organization, No. 2, p. 100) and entered into force on April 7, 1948.

## Appendix

Note this list includes the 202 works that were reviewed. The asterisks denote works that presented material (be it a discussed benefit, motivation, reason, and/or perceived consequence/outcome, etc.) that we included as a correlate of well-being.

- \* Adams, K. (2000). How can I keep from singing: A lifetime of sweet singing in the choir. In B. A. Roberts & A. Rose (Eds.), *Sharing the voices: The phenomenon of singing 2. Proceedings of the International Symposium*. St. John's, Newfoundland, Canada: Memorial University of Newfoundland.
- Adams, K. B., Leibbrandt, S., & Moon, H. (2011). A critical review of the literature on social and leisure activity and wellbeing in later life. *Ageing and Society*, 31, 683-712. DOI: 10.1017/S0144686X10001091
- \* Adderley, C., Kennedy, M., & Berz, W. (2003). "A home away from home": The world of the music classroom. *Journal of Research in Music Education*, 51, 190-205.
- Antoniadou, M., Michaelidis, V., & Tsara, V. (2012). Lung function in wind instrument players. *Pneumon*, 25(2), 180-183.
- Ashley, M. (2002). Singing, gender and health: Perspectives from boys singing in a church choir. *Health Education*, 102(4), 180-187. DOI: 10.1108/09654280210434255
- \* Bailey, B. A., & Davidson, J. W. (2002). Adaptive characteristics of group singing: perceptions from members of a choir for homeless men. *Musicae Scientiae*, 6(2), 221-256. DOI: 10.1177/102986490200600206
- \* Bailey, B. A., & Davidson, J. W. (2003). Amateur Group Singing as a Therapeutic Instrument. *Nordic Journal of Music Therapy*, 12(1), 18-32. DOI: 10.1080/08098130309478070
- \* Bailey, B. A., & Davidson, J. W. (2005). Effects of group singing and performance for marginalized and middle-class singers. *Psychology of Music*, 33(3), 269-303. DOI: 10.1177/0305735605053734
- \* Ballantyne, J., Ballantyne, R., & Packer, J. (2013). Designing and managing music festival experiences to enhance attendees' psychological and social benefits. *Musicae Scientiae*, 18(1), 65-83. DOI: 10.1177/1029864913511845
- Bamford, A., & Clift, S. (2006). *Making singing for health happen: Reflections on a 'singing for the brain' training course*. Canterbury, UK: Canterbury Christ Church University.
- Bamford, A., & Clift, S. (2006). *Music into Upton: Reflections on an initiative to bring live music into a hospital setting*. Canterbury, UK: Canterbury Christ Church University.
- Batt-Rawden, K. B., DeNora, T., & Ruud, E. (2005). Music listening and empowerment in health promotion: A study of the role and significance of music in everyday life of the long-term ill. *Nordic Journal of Music Therapy*, 14(2), 120-136. DOI: 10.1080/08098130509478134
- \* Beck, R. J., Cesario, T. C., Yousefi, A., & Enamoto, H. (2000). Choral singing, performance perception, and immune system changes in salivary immunoglobulin A and cortisol. *Music Perception*, 18, 87-106.

- Beck, R. J., Gottfried, T. L., Hall, D. J., Cisler, C. A., & Bozeman, K. W. (2006). Supporting the health of college solo singers: The relationship of positive emotions and stress to changes in salivary IgA and cortisol during singing. *Journal of Learning through the Arts: A Research Journal on Arts Integration in Schools and Communities*, 2(1), article 19.
- Biley, F. C. (2000). The effects on patient well-being of music listening as a nursing intervention: A review of the literature. *Journal of Clinical Nursing*, 9, 668-677.
- \* Boer, D., & Fischer, R. (2011). The functions of music-listening across cultures: The development of a scale measuring personal, social and cultural functions of music. In F. Deutsch, M. Boehnke, U. Kuehnen, & K. Boehnke (Eds.), *Crossing borders: Cross-cultural and cultural psychology as an interdisciplinary, multi-method endeavour* (pp. 201-215). Bremen: Jacobs-University.
- \* Boer, D., & Fischer, R. (2012). Towards a holistic model of functions of music listening across cultures: A culturally decentred qualitative approach. *Psychology of Music*, 40(2), 179-200. DOI: 10.1177/0305735610381885
- \* Bonde, L. O. (2013). The musical identities of Danish music therapy students: A study based on musical autobiographies. In L. O. Bonde, E. Ruud, M. S. Skånland, & G. Trondalen (Eds.), *Musical life stories. Narratives on health musicking* (pp. 307-328). Oslo: Centre for Music and Health, the Norwegian Academy of Music.
- \* Bonde, L. O. (2014). Music and health promotion - In the life of music therapy and music psychology researchers: A pilot study. *Voices: A World Forum for Music Therapy*, 14(1), n.p. DOI: 10.15845/voices.v14i1.740
- Bonilha, A. G., Onofre, F., Vieira, M. L., Prado, M. Y. A., & Martinez, J. A. B. (2009). Effects of singing classes on pulmonary function and quality of life of COPD patients. *International Journal of COPD*, 4, 1-8.
- \* Bruhn, H. (2002). Musical development of elderly people. *Psychomusicology*, 18, 59-75.
- Bugos, J. A., Perlstein, W. M., McCrae, C. S., Brophy, T. S., & Bedenbaugh, P. H. (2007). Individualized piano instruction enhances executive functioning and working memory in older adults. *Aging and Mental Health*, 11(4), 464-471. DOI: 10.1080/13607860601086504
- \* Bungay, H., & Skingley, A. (2008). *The silver song club project: Summary of a formative evaluation*. Canterbury, UK: Canterbury Christ Church University.
- Bygren, L. O., Konlaan, B. B., & Johansson, S.-E. (1996). Attendance at cultural events, reading books or periodicals, and making music or singing in a choir as determinants for survival: Swedish interview survey of living conditions. *British Medical Journal*, 313, 1577-1580.
- \* Camic, P. M., Williams, C. M., & Meeten, F. (2011). Does a 'Singing Together Group' improve the quality of life of people with a dementia and their carers? A pilot evaluation study. *Dementia*, 12(2), 157-176. DOI: 10.1177/1471301211422761
- \* Cavitt, M. E. (2005). Factors influencing participation in community bands. *Journal of Band Research*, 41(1), 42-57.

- \* Chamorro-Premuzic, T., & Furnham, A. (2007). Personality and music: Can traits explain how people use music in everyday life? *British Journal of Psychology*, *98*, 175-185. DOI: 10.1348/000712606X111177
- \* Chamorro-Premuzic, T., Swami, V., & Cermakova, B. (2012). Individual differences in music consumption are predicted by uses of music and age rather than emotional intelligence, neuroticism, extraversion or openness. *Psychology of Music*, *40*(3), 285-300. DOI: 10.1177/0305735610381591
- Chin, T., & Rickard, N. S. (2013). Emotion regulation strategy mediates both positive and negative relationships between music uses and well-being. *Psychology of Music*, *42*(5), 692-713. DOI: 10.1177/0305735613489916
- Clair, A. A., Mathews, M., & Kosloski, K. (2005). Assessment of active music participation as an indication of subsequent music making engagement for persons with midstage dementia. *American Journal of Alzheimer's Disease and Other Dementias*, *20*(1), 37-40.
- Clayton, M. (2009). The social and personal function of music in cross-cultural perspective. In S. Hallam, I. Cross, & M. Thaut (Eds.), *The Oxford handbook of music psychology* (pp. 35-44). Oxford, UK: Oxford University Press.
- Clift, S. (2012). Creative arts as a public health resource: moving from practice-based research to evidence-based practice. *Perspectives in Public Health*, *132*(3), 120-127. DOI: 10.1177/1757913912442269
- \* Clift, S. M., & Hancox, G. (2001). The perceived benefits of singing: Findings from preliminary surveys of a university college choral society. *The Journal of the Royal Society for the Promotion of Health*, *121*(4), 248-256.
- \* Clift, S., & Hancox, G. (2010). The significance of choral singing for sustaining psychological wellbeing: Findings from a survey of choristers in England, Australia and Germany. *Music Performance Research*, *3*(1), 79-96.
- \* Clift, S., Hancox, G., Morrison, I., Hess, B., Kreutz, G., & Stewart, D. (2009). *What do singers say about the effects of choral singing on physical health? Findings from a survey of choristers in Australia, England and Germany*. Paper presented at the Proceedings of the 7th Triennial Conference of European Society for the Cognitive Sciences of Music, Jyväskylä, Finland.
- \* Clift, S., Hancox, G., Morrison, I., Hess, B., Kreutz, G., & Stewart, D. (2010). Choral singing and psychological wellbeing: Quantitative and qualitative findings from English choirs in a cross-national survey. *Journal of Applied Arts and Health*, *1*(1), 19-34. DOI: 10.1386/jaah.1.1.19/1
- \* Clift, S., Hancox, G., Morrison, I., Hess, B., Stewart, D., & Kreutz, G. (2008). *Choral singing, wellbeing and health: Summary of findings from a cross-national survey*. Canterbury, UK: Canterbury Christ Church University.
- Clift, S., Hancox, G., Staricoff, R., & Whitmore, C. (2008). *Singing and health: Summary of a systematic mapping and review of non-clinical research*. Canterbury, UK: Canterbury Christ Church University.
- \* Clift, S., Mackenzie, K., & Bushell, F. (2006). *The MusicStart project: Evaluation of an Isle of Wight early years music initiative*. Canterbury, UK: Canterbury Christ Church University.
- Coffman, D. D. (2002). Music and quality of life in older adults. *Psychomusicology*, *18*(1-2), 76-88.

- Coffman, D. D. (2006). Voices of experience: Interviews of adult community band members in Launceston, Tasmania, Australia. *International Journal of Community Music*, n.p. Retrieved from <http://www.intljcm.com/articles/Volume%204/Coffman/Coffman.pdf>
- Coffman, D. D. (2007). An exploration of personality traits in older adult amateur musicians. *Research and Issues in Music Education*, 5(1), n.p.
- \* Coffman, D. D. (2008). Survey of New Horizons International Music Association musicians. *International Journal of Community Music*, 1(3), 375-390. DOI: 10.1386/ijcm.1.3.375/1
- Coffman, D. D. (2009). Learning from our elders: survey of New Horizons International Music Association band and orchestra directors. *International Journal of Community Music*, 2(2&3), 227-240. DOI: 10.1386/ijcm.2.2&3.227/1
- Coffman, D. D., & Adamek, M. S. (1999). The contributions of wind band participation to quality of life of senior adults. *Music Therapy Perspectives*, 17, 27-31.
- Cohen, A. (2002). Psychology of music and aging comes of age: Psychogeromusicology. *Psychomusicology*, 18, 3-9.
- Cohen, A., Bailey, B., & Nilsson, T. (2002). The importance of music to seniors. *Psychomusicology*, 18, 89-102.
- Cohen, G. D., Perlstein, S., Chapline, J., Kelly, J., Firth, K. M., & Simmens, S. (2006). The impact of professionally conducted cultural programs on the physical health, mental health, and social functioning of older adults. *The Gerontologist*, 46(6), 726-734.
- Cohen, G. D., Perlstein, S., Chapline, J., Kelly, J., Firth, K. M., & Simmens, S. (2007). The impact of professionally conducted cultural programs on the physical health, mental health, and social functioning of older adults—2-year results. *Journal of Aging, Humanities and the Arts*, 1, 5-22. DOI: 10.1080/19325610701410791
- \* Cohen, M. L. (2007). Explorations of inmate and volunteer choral experiences in a prison-based choir. *Australian Journal of Music Education*, 1, 61-72.
- Cohen, M. L. (2009). Choral singing and prison inmates: Influences of performing in a prison choir. *Journal of Correctional Education*, 60(1), 52-65.
- Constantinescu, M. (2013). Educating young people for quality of life improvement. *Procedia - Social and Behavioral Sciences*, 93, 395-399. DOI: 10.1016/j.sbspro.2013.09.210
- \* Conway, C., & Hodgman, T. M. (2008). College and community choir member experiences in a collaborative, intergenerational performance project. *Journal of Research in Music Education*, 56(3), 220-237. DOI: 10.1177/0022429408327297
- Cooke, M., Moyle, W., Shum, D., Harrison, S., & Murfield, J. (2010). A Randomized Controlled Trial exploring the effect of music on quality of life and depression in older people with dementia. *Journal of Health Psychology*, 15(5), 765-776. DOI: 10.1177/1359105310368188
- \* Cooper, T. L. (2001). Adults' perceptions of piano study: Achievements and experiences. *Journal of Research in Music Education*, 49(2), 156-168.
- \* Creech, A., Hallam, S., Gaunt, H., McQueen, H., Pincas, A., & Varvarigou, M. (2013). The power of music in the lives of older adults. *Research Studies in Music Education*, 35(1), 87-102. DOI: 10.1177/1321103X13478862

- Creech, A., Hallam, S., Varvarigou, M., McQueen, H., & Gaunt, H. (2013). Active music making: A route to enhanced subjective well-being among older people. *Perspectives in Public Health*, 133(1), 36-43. DOI: 10.1177/1757913912466950
- Croom, A. M. (2015). Music practice and participation for psychological well-being: A review of how music influences positive emotion, engagement, relationships, meaning, and accomplishment. *Musicae Scientiae*, 19(1), 44-64. DOI: 10.1177/1029864914561709
- \* Dabback, W. M. (2009). Identity formation through participation in the Rochester New Horizons band programme. *International Journal of Community Music*, 1(2), 267-286. DOI: 10.1386/ijcm.1.2.267/1
- Davidson, J. W., & Fedele, J. (2011). Investigating group singing activity with people with dementia and their caregivers: Problems and positive prospects. *Musicae Scientiae*, 15(3), 402-422. DOI: 10.1177/1029864911410954
- Di Benedetto, P., Cavazzon, M., Mondolo, F., Rugiu, G., Peratoner, A., & Biasutti, E. (2009). Voice and choral singing treatment: A new approach for speech and voice disorders in Parkinson's disease. *European Journal of Physical and Rehabilitation Medicine*, 45(1), 13-19.
- \* Dingle, G. A., Brander, C., Ballantyne, J., & Baker, F. A. (2012). 'To be heard': The social and mental health benefits of choir singing for disadvantaged adults. *Psychology of Music*, 41(4), 405-421. DOI: 10.1177/0305735611430081
- \* Douglas, K. A. (2011). *A descriptive analysis of the psychological needs of adults participating in music ensembles: A survey of the New Horizons International Music Association ensemble participants* (Doctoral dissertation). Available from ProQuest Dissertations and Theses database. AAT3473456.
- Driscoll, J. (2009). 'If I play my sax my parents are nice to me': opportunity and motivation in musical instrument and singing tuition. *Music Education Research*, 11(1), 37-55. DOI: 10.1080/14613800802547722
- Eley, R., & Gorman, D. (2008). Music therapy to manage asthma. *Aboriginal and Islander Health Worker Journal*, 32(1), 9-10.
- \* Eley, R., & Gorman, D. (2010). Didgeridoo playing and singing to support asthma management in Aboriginal Australians. *The Journal of Rural Health*, 26, 100-104. DOI: 10.1111/j.1748-0361.2009.00256.x
- \* Engen, R. L. (2005). The singer's breath: Implications for treatment of persons with emphysema. *Journal of Music Therapy*, 42(1), 20-48.
- \* Forssén, A. S. K. (2007). Humour, beauty, and culture as personal health resources: Experiences of elderly swedish women. *Scandinavian Journal of Public Health*, 35(3), 228. DOI: 10.1080/14034940601160680
- Galati, D., Costa, T., Rognoni, E., & Pisterzi, A. (2006, August). *The effect of singing on subjective emotional state and psychophysiological response in singers*. Paper presented at the 9th International Conference on Music Perception and Cognition, Alma Mater Studiorum, University of Bolgna.
- \* Gembris, H. (2008). Musical activities in the third age: An empirical study with amateur musicians. In A. Daubney, E. Longhi, A. Lamont, & D. J. Hargreaves (Eds.), *Second European Conference on Developmental Psychology of Music* (pp. 103-108). Roehampton University, England.



- \* Gembris, H. (2012). Music-making as a lifelong development and resource for health. In R. A. R. MacDonald, G. Kreutz, & L. Mitchell (Eds.), *Music, health, and wellbeing* (pp. 367-382). Oxford, UK: Oxford University Press.
- Gembris, H., & Heye, A. (2014). Growing older in a symphony orchestra: The development of the age-related self-concept and the self-estimated performance of professional musicians in a lifespan perspective. *Musicae Scientiae*, 18(4), 371-391. DOI: 10.1177/1029864914548912
- Getz, L. M., Chamorro-Premuzic, T., Roy, M. M., & Devroop, K. (2010). The relationship between affect, uses of music, and music preferences in a sample of South African adolescents. *Psychology of Music*, 40(2), 164-178. DOI: 10.1177/0305735610381818
- Getz, L. M., Marks, S., & Roy, M. (2014). The influence of stress, optimism, and music training on music uses and preferences. *Psychology of Music*, 42(1), 71-85. DOI: 10.1177/0305735612456727
- \* Gick, M. L. (2011). Singing, health and well being: A health psychologist's review. *Psychomusicology: Music, Mind & Brain*, 21(1&2), 176-207. DOI: 10.1037/h0094011
- Gold, A., & Clare, A. (2013). An exploration of music listening in chronic pain. *Psychology of Music*, 41(5), 545-564. DOI: 10.1177/0305735612440613
- Grape, C., Sandgren, M., Hansson, L.-O., Ericson, M., & Theorell, T. (2003). Does singing promote well-being? An empirical study of professional and amateur singers during a singing lesson. *Integrative Physiological and Behavioral Science*, 38(1), 65-74.
- \* Greaves, C. J., & Farbus, L. (2006). Effects of creative and social activity on the health and well-being of socially isolated older people: Outcomes from a multi-method observational study. *Journal of the Royal Society for the Promotion of Health*, 126(3), 134-142. DOI: 10.1177/1466424006064303
- \* Gridley, H., Astbury, J., Sharples, J., & Aguirre, C. (2011). *Benefits of group singing for community mental health and wellbeing: Survey and literature review*. Carlton, VIC: Victorian Health Promotion Foundation.
- \* Haake, A. B. (2011). Individual music listening in workplace settings: An exploratory survey of offices in the UK. *Musicae Scientiae*, 15(1), 107-129. DOI: 10.1177/1029864911398065
- \* Habron, J., Butterly, F., Gordon, I., & Roebuck, A. (2013). Being well, being musical: Music composition as a resource and occupation for older people. *British Journal of Occupational Therapy*, 76(7), 308-316. DOI: 10.4276/030802213X13729279114933
- Hacking, S., Secker, J., Spandler, H., Kent, L., & Shenton, J. (2008). Evaluating the impact of participatory art projects for people with mental health needs. *Health and Social Care in the Community*, 16(6), 638-648. DOI: 10.1111/j.1365-2524.2008.00789.x
- \* Hallam, S., & MacDonald, R. A. R. (2009). The effects of music in community and educational settings. In S. Hallam, I. Cross, & M. Thaut (Eds.), *The Oxford handbook of music psychology* (pp. 471-480). Oxford, UK: Oxford University Press.

- Hallam, S., Creech, A., McQueen, H., & Varvarigou, M. (2011). Perceptions of effective leadership in music facilitators working with older people. *Journal of Arts and Communities*, 3(3), 229-248.
- \* Hallam, S., Creech, A., McQueen, H., Varvarigou, M., & Gaunt, H. (2016). The facilitator of community music-making with older learners: Characteristics, Motivations and challenges. *International Journal of Music Education*, 34(1), 19-31. DOI: 10.1177/0255761415617039
- \* Hallam, S., Creech, A., Varvarigou, M., & McQueen, H. (2012). Perceived benefits of active engagement with making music in community settings. *International Journal of Community Music*, 5(2), 155-174.
- \* Hallam, S., Creech, A., Varvarigou, M., & McQueen, H. (2012). The characteristics of older people who engage in community music making, their reasons for participation and the barriers they face. *Journal of Adult and Continuing Education*, 18(2), 21-43. DOI: 10.7227/JACE.18.2.3
- Hallam, S., Creech, A., Varvarigou, M., McQueen, H., & Gaunt, H. (2014). Does active engagement in community music support the well-being of older people? *Arts & Health: An International Journal for Research, Policy and Practice*, 6(2), 101-116. DOI: 10.1080/17533015.2013.809369
- \* Hays, T. (2005). Well-being in later life through music. *Australasian Journal on Ageing*, 24(1), 28-32.
- \* Hays, T., & Minichiello, V. (2005). The meaning of music in the lives of older people: A qualitative study. *Psychology of Music*, 33(4), 437-451. DOI: 10.1177/0305735605056160
- \* Hillman, S. (2002). Participatory singing for older people: A perception of benefit. *Health Education*, 102(4), 163-171. DOI: 10.1108/09654280210434237
- \* Hills, P., & Argyle, M. (1998). Musical and religious experiences and their relationship to happiness. *Personality and Individual Differences*, 25, 91-102.
- Horvath, K. A. (2008). Adopting a healthy approach to instrumental music making. *Music Educators Journal*, 94(3), 30-34.
- Houston, D. M., McKee, K. J., Carroll, L., & Marsh, H. (1998). Using humour to promote psychological wellbeing in residential homes for older people. *Ageing and Mental Health*, 2(4), 328-332. DOI: 10.1080/13607869856588
- \* Hui, E., Chui, B. T.-k., & Woo, J. (2009). Effects of dance on physical and psychological well-being in older persons. *Archives of Gerontology and Geriatrics*, 49, e45-e50. DOI: 10.1016/j.archger.2008.08.006
- Hulme, C., Wright, J., Crocker, T., Oluboyede, Y., & House, A. (2010). Non-pharmacological approaches for dementia that informal carers might try or access: a systematic review. *International Journal of Geriatric Psychiatry*, 25, 756-763.
- Hyppä, M. T., & Mäki, J. (2001). Individual-level relationships between social capital and self-rated health in a bilingual community. *Preventive Medicine*, 32. DOI: 10.1006/pmed.2000.0782
- Hyppä, M. T., & Maki, J. (2003). Social participation and health in a community rich in stock of social capital. *Health Education Research*, 18(6), 770-779. DOI: 10.1093/her/cyf044

- \* Jacob, C., Guptill, C., & Sumsion, T. (2009). Motivation for continuing involvement in a leisure-based choir: The lived experiences of university choir members. *Journal of Occupational Science, 16*(3), 187-193. DOI: 10.1080/14427591.2009.9686661
- Johansson, L., & Theorell, T. (2003). Satisfaction with work task quality correlates with employee health. *Medical Problems of Performing Artists, 18*(4), 141-149.
- Johansson, S. E., Konlaan, B. B., & Bygren, L. O. (2001). Sustaining habits of attending cultural events and maintenance of health: A longitudinal study. *Health Promotion International, 16*(3), 229-234.
- \* Joseph, D. (2009). Sharing music and culture through singing in Australia. *International Journal of Community Music, 2*(2&3), 169-181. DOI: 10.1386/ijcm.2.2&3.169/1
- Joyce, Moon (1996). Turn off the radio and sing for your lives! Women, singing, and experiential education. In K. Warren (Ed.), *Women and experiential education* (253-266). Dubuque: Kendall/Hunt.
- \* Judd, M., & Pooley, J. A. (2014). The psychological benefits of participating in group singing for members of the general public. *Psychology of Music, 42*(2), 269-283. DOI: 10.1177/0305735612471237
- \* Juslin, P. N., & Laukka, P. (2004). Expression, perception, and induction of musical emotions: A review and a questionnaire study of everyday listening. *Journal of New Music Research, 33*(3), 217-238. DOI: 10.1080/0929821042000317813
- \* Jutras, P. J. (2006). The benefits of adult piano study as self-reported by selected adult piano students. *Journal of Research in Music Education, 54*(2), 97-110.
- \* Jutras, P. J. (2011). The benefits of new horizons band participation as self-reported by selected New Horizons band members. *Bulletin of the Council for Research in Music Education, 187*, 65-84.
- \* Kennedy, M. A. (2002). It's cool because we like to sing: Junior high school boys' experience of choral music as an elective. *Research Studies in Music Education, 18*, 24-35.
- Kenny, D. T., & Faunce, G. (2004). The impact of group singing on mood, coping, and perceived pain in chronic pain patients attending a multidisciplinary pain clinic. *Journal of Music Therapy, 41*(3), 241-258.
- Kenny, D., Driscoll, T., & Ackermann, B. (2014). Psychological well-being in professional orchestral musicians in Australia: A descriptive population study. *Psychology of Music, 42*(2), 201-131. DOI: 10.1177/0305735612463950
- Kleinerman, K. (2008). Women sing, women lead: The transformation of identity and emergence of leadership in women through voice. *GEMS- Gender, Education, Music, and Society, 4*. Accessed online at [www.queensu.ca/music/links/gems/kleinerman5.pdf](http://www.queensu.ca/music/links/gems/kleinerman5.pdf)
- \* Kokotsaki, D., & Hallam, S. (2007). Higher education music students' perceptions of the benefits of participative music making. *Music Education Research, 9*(1), 93-109.
- \* Kokotsaki, D., & Hallam, S. (2011). The perceived benefits of participative music making for non-music university students: a comparison with music students. *Music Education Research, 13*(2), 149-172. DOI: 10.1080/14613808.2011.577768
- Konieczna-Nowak, L. (2015). Music, text, music-and-text and psychophysiological

- responses: A randomized controlled trial. *The Arts in Psychotherapy*, 42, 57-62. DOI: 10.1016/j.aip.2014.12.004
- Konlaan, B. B., Björby, N., Bygren, L. O., Weissglas, G., Karlsson, L. G., & Widmark, M. (2000). Attendance at cultural events and physical exercise and health: A randomized controlled study. *Public Health*, 114, 316-319.
- Konlaan, B. B., Bygren, L. O., & Johansson, S.-E. (2000). Visiting the cinema, concerts, museums or art exhibitions as determinant of survival: A Swedish fourteen-year cohort follow-up. *Scandinavian Journal of Public Health*, 28(3), 174-178.
- Kreutz, G. (2014). Does singing facilitate social bonding? *Music and Medicine*, 6(2), 51-60.
- Kreutz, G., Bongard, S., Rohrmann, S., Hodapp, V., & Grebe, D. (2004). Effects of choir singing or listening on secretory immunoglobulin A, cortisol and emotional state. *Journal of Behavioral Medicine*, 27(6), 623-635.
- Kreutz, G., Ginsborg, J., & Williamon, A. (2009). Health-promoting behaviours in conservatoire students. *Psychology of Music*, 37(1), 47-60. DOI: 10.1177/0305735607086047
- Krout, R. E. (2007). Music listening to facilitate relaxation and promote wellness: Integrated aspects of our neurophysiological responses to music. *The Arts in Psychotherapy*, 34, 134-141. DOI: 10.1016/j.aip.2006.11.001
- Kuhn, D. (2002). The effects of active and passive participation in musical activity on the immune system as measured by salivary immunoglobulin A (SigA). *Journal of Music Therapy*, 39(1), 30-39.
- Lai, H.-L., Chang, E.-T., Li, Y.-M., Huang, C.-Y., Lee, L.-H., & Wang, H.-M. (2015). Effects of music videos on sleep quality in middle-aged and older adults with chronic insomnia: A randomized controlled trial. *Biological Research for Nursing*, 17(3), 340-347. DOI: 10.1177/1099800414549237
- \* Laiho, S. (2004). The psychological functions of music in adolescence. *Nordic Journal of Music Therapy*, 13(1), 47-63.
- \* Lally, E. (2009). 'The power to heal us with a smile and a song': Senior well-being, music-based participatory arts and the value of qualitative evidence. *Journal of Arts and Communities*, 1(1), 25-44. DOI: 10.1386/jaac.1.1.25/1
- Lane, J. S. (2012). Engaging preservice music teachers with adult amateur musicians. *International Journal of Community Music*, 5(3), 317-322. DOI: 10.1386/ijcm.5.3.317\_1
- \* Laukka, P. (2007). Uses of music and psychological well-being among the elderly. *Journal of Happiness Studies*, 8, 215-341. DOI: 10.1007/s10902-006-9024-3
- \* Lehmborg, L. J., & Fung, C. V. (2010). Benefits of music participation for senior citizens: A review of the literature. *Music Education Research International*, 4, 19-30.
- Lenette, C., & Sunderland, N. (2016). "Will there be music for us?" Mapping the health and well-being potential of participatory music practice with asylum seekers and refugees across contexts of conflict and refuge. *Arts & Health: An International Journal for Research, Policy and Practice*, 8(1), 32-49. DOI: 10.1080/17533015.2014.961943

- \* Livesey, L., Morrison, I., Clift, S., & Camic, P. (2012). Benefits of choral singing for social and mental wellbeing: Qualitative findings from a cross-national survey of choir members. *Journal of Public Mental Health, 11*(1), 10-26. DOI: 10.1108/17465721211207275
- Longhi, E., & Pickett, N. (2008). Music and well-being in long-term hospitalized children. *Psychology of Music, 36*(2), 247-256. DOI: 10.1177/0305735607082622
- Longhi, E., Pickett, N., & Hargreaves, D. J. (2015). Wellbeing and hospitalized children: Can music help? *Psychology of Music, 43*(2), 188-196. DOI: 10.1177/0305735613499781
- \* Lonsdale, A. J., & North, A. C. (2011). Why do we listen to music? A uses and gratifications analysis. *British Journal of Psychology, 102*(1), 108-134. DOI: 10.1348/000712610X506831
- \* Lord, V. M., Cave, P., Hume, V. J., Flude, E. J., Evans, A., Kelly, J. L., Hopkinson, N. S. (2010). Singing teaching as a therapy for chronic respiratory disease - A randomised controlled trial and qualitative evaluation. *BMC Pulmonary Medicine, 10*, 41.
- \* Louhivuori, J., Salminen, V.-M., & Lebaka, E. (2005). "Singing together" – A cross-cultural approach to the meaning of choirs as a community. In P. S. Campbell, J. Drummond, P. Dunbar-Hall, K. Howard, H. Schippers, & T. Wiggins (Eds.), *Cultural diversity in music education: Directions and challenges for the 21st century* (81-92). Bowen Hills, Qld: Australian Academic Press.
- MacDonald, R. A. R. (2013). Music, health, and well-being: A review. *International Journal of Qualitative Studies on Health and Well-being, 8*, 20635. DOI: 10.3402/qhw.v8i0.20635
- Magee, W. L. (2011). Music technology for health and well-being: The bridge between the arts and science. *Music and Medicine, 3*(3), 131-133. DOI: 10.1177/1943862111411719
- Magee, W. L., & Davidson, J. W. (2004). Music therapy in multiple sclerosis: Results of a systematic qualitative analysis. *Music Therapy Perspectives, 22*, 39-51.
- Magee, W. L., & Davidson, J. W. (2004). Singing in therapy: Monitoring disease process in chronic degenerative illness. *British Journal of Music Therapy, 18*(2), 65-77.
- Mantie, R. (2012). A study of community band participants: Implications for music education. *Bulletin of the Council for Research in Music Education, 191*, 21-43.
- \* McFerran, K. S., & Saarikallio, S. (2014). Depending on music to feel better: Being conscious of responsibility when appropriating the power of music. *The Arts in Psychotherapy, 41*, 89-97. DOI: 10.1016/j.aip.2013.11.007
- \* McQueen, H., Hallam, S., Creech, A., & Varvarigou, M. (2013). A philosophical perspective on leading music activities for the over 50s. *International Journal of Lifelong Education, 32*(3), 353-377. DOI: 10.1080/02601370.2012.738432
- \* Michalos, A. C. (2005). Arts and the quality of life: An exploratory study. *Social Indicators Research, 71*, 11-59. DOI: 10.1007/s11205-004-8013-3

- \* Murcia, C. Q., Kreutz, G., Clift, S., & Bongard, S. (2010). Shall we dance? An exploration of the perceived benefits of dancing on well-being. *Arts and Health: An International Journal for Research, Policy and Practice*, 2(2), 149-163. DOI: 10.1080/17533010903488582
- Myskja, A., & Nord, P. G. (2008). "The Day the Music Died": A pilot study on music and depression in a nursing home. *Nordic Journal of Music Therapy*, 17(1), 30-40. DOI: 10.1080/08098130809478194
- Nyqvist, F., Finnäs, F., Jakobsson, G., & Koskinen, S. (2008). The effect of social capital on health: The case of two language groups in Finland. *Health and Place*, 14, 347-360. DOI: 10.1016/j.healthplace.2007.09.001
- O'Shea, H. (2012). 'Get back to where you once belonged!' The positive creative impact of a refresher course for 'baby-boomer' rock musicians. *Popular Music*, 31(2), 199-215. DOI: 10.1017/S0261143012000025
- O'Connor, D. W., Ames, D., Gardner, B., & King, M. (2009). Psychosocial treatments of behavior symptoms in dementia: A systematic review of reports meeting quality standards. *International Psychogeriatrics*, 21(2), 225-240. DOI: 10.1017/S1041610208007588
- O'Connor, D. W., Ames, D., Gardner, B., & King, M. (2009). Psychosocial treatments of psychological symptoms in dementia: A systematic review of reports meeting quality standards. *International Psychogeriatrics*, 21(2), 241-251. DOI: 10.1017/S1041610208008223
- \* Packer, J., & Ballantyne, J. (2011). The impact of music festival attendance on young people's psychological and social well-being. *Psychology of Music*, 39(2), 164-181. DOI: 10.1177/0305735610372611
- \* Palmer, R. (2008). Questions arising from the views of some members of four amateur classical music organizations. *International Journal of Community Music*, 1(2), 203-216. DOI: 10.1386/ijcm.1.2.203/1
- Patteson, A. (2000). Singing a woman's life: How singing lessons transformed the lives of nine women. In B. A. Roberts & A. Rose (Eds.), *Sharing the voices: The phenomenon of singing 2. Proceedings of the International Symposium*. St. John's, Newfoundland, Canada, Memorial University of Newfoundland.
- \* Pavlakou, M. (2009). Benefits of group singing for people with eating disorders: Preliminary findings from a non-clinical study. *Approaches: Music Therapy and Special Music Education*, 1(1), 30-48.
- Pickles, V. (2003). Music and the third age. *Psychology of Music*, 31(4), 415-423.
- Pitts, S. E. (2004). 'Everybody wants to be Pavarotti': The experience of music for performers and audience at a Gilbert and Sullivan festival. *Journal of the Royal Musical Association*, 129(1), 143-160.
- \* Pothoulaki, M., MacDonald, R. A. R., & Flowers, P. (2012). An interpretative phenomenological analysis of an improvisational music therapy program for cancer patients. *Journal of Music Therapy*, 49(1), 45-67.
- Pothoulaki, M., MacDonald, R. A. R., & Flowers, P. (2012). The use of music in chronic illness: Evidence and arguments. In R. A. R. MacDonald, G. Kreutz, & L. Mitchell (Eds.), *Music, health, and wellbeing* (pp. 239-256). Oxford, UK: Oxford University Press.

- Racette, A., Bard, C., & Peretz, I. (2006). Making non-fluent aphasics speak: Sing along! *Brain*, *129*, 2571-2584. DOI: 10.1093/brain/awl250
- Raglio, A., & Oasi, O. (2015). Music and health: what interventions for what results? *Frontiers in Psychology*, *6*, 230. DOI: 10.3389/fpsyg.2015.0023
- Raglio, A., Filippi, S., Bellandi, D., & Stramba-Badiale, M. (2014). Global music approach to persons with dementia: Evidence and practice. *Clinical Interventions in Aging*, *9*, 1669-1676.
- Rohwer, D. (2008). Health and wellness issues for adult band musicians. *Medical Problems of Performing Artists*, *23*, 54-58.
- \* Rohwer, D. (2009). Perceived instructional needs and desires of 8th grade and senior citizen band members. *Journal of Music Teacher Education*, *18*(2), 1-12.
- \* Rohwer, D. (2010). Church musicians' participation perceptions: Applications to community music. *Research and Issues in Music Education*, *8*(1), n.p. Retrieved from <http://files.eric.ed.gov/fulltext/EJ894767.pdf>
- Rohwer, D. (2010). Understanding adult interests and needs: The pitfalls in wanting to know. *International Journal of Community Music*, *3*(2), 203-212. DOI: 10.1386/ijcm.3.2.203\_1
- \* Rohwer, D. (2012). How participants envision community music in Welsh men's choirs. *Research and Issues in Music Education*, *10*(1), n.p. Retrieved from <http://www.stthomas.edu/rimeonline/vol10/rohwer.htm>
- \* Rohwer, D. (2013). *Making music as an adult: What do the spouses think?* Texas Music Education Research. Austin, TX: Texas Music Education Research.
- \* Rohwer, D., & Rohwer, M. (2009). A content analysis of choral students' participation perceptions: Implications for lifelong learning. *International Journal of Community Music*, *2*, 255-262. DOI: 10.1386/ijcm.2.2&3.255/1
- Rohwer, D., Coffman, D. D., & Raiber, M. (2013). Random or non-random thoughts: What senior adults think about within their ensemble settings. *International Journal of Community Music*, *5*(3), 289-302. DOI: 10.1386/ijcm.5.3.289\_1
- \* Saarikallio, S. (2012). Cross-cultural approaches to music and health. In R. A. R. MacDonald, G. Kreutz, & L. Mitchell (Eds.), *Music, health, and wellbeing* (pp. 477-490). Oxford, UK: Oxford University Press.
- Sandgren, M. (2009). Health anxiety instead of performance anxiety among opera singers. In J. Louhivuori, T. Eerola, S. Saarikallio, T. Himberg, & P.-S. Eerola (Eds.), *Proceedings of the 7th triennial conference of European Society for the Cognitive Sciences of Music*. Jyväskylä, Finland: ESCOM.
- Särkämö, T., Tervaniemi, M., Laitinen, S. L., Numminen, A., Kurki, M., Johnson, J. K., & Rantanen, P. (2013). Cognitive, emotional, and social benefits of regular musical activities in early dementia: Randomized controlled study. *The Gerontologist*, *54*(4), 634-650. DOI: 10.1093/geront/gnt100
- \* Schäfer, T., & Sedlmeier, P. (2009). From the functions of music to music preference. *Psychology of Music*, *37*(3), 279-300. DOI: 10.1177/0305735608097247
- \* Schäfer, T., & Sedlmeier, P. (2010). What makes us like music? Determinants of music preference. *Psychology of Aesthetics, Creativity, and the Arts*, *4*(4), 223-234. DOI: 10.1037/a0018374

- \* Schäfer, T., Sedlmeier, P., Städtler, C., & Huron, D. (2013). The psychological functions of music listening. *Frontiers in Psychology*, 4, 311. DOI: 10.3389/fpsyg.2013.00511
- \* Schäfer, T., Tipandjan, A., & Sedlmeier, P. (2012). The functions of music and their relationship to music preference in India and Germany. *International Journal of Psychology*, 47(5), 370-380. DOI: 10.1080/00207594.2012.688133
- \* Shansky, C. (2010). Adult motivations in community orchestra participation: A pilot case study of the Bergen philharmonic orchestra (New Jersey). *Research and Issues in Music Education*, 8(1), n.p. Retrieved from <http://www.stthomas.edu/rimeonline/vol8/>
- \* Sheldon, D. A. (1998). Participation in community and company bands in Japan. *Applications of Research in Music Education*, 17(1), 21-24. DOI: 10.1177/875512339801700105
- \* Silber, L. (2005). Bars behind bars: the impact of a women's prison choir on social harmony. *Music Education Research*, 7(2), 251-271. DOI: 10.1080/14613800500169811
- \* Sixsmith, A., & Gibson, G. (2007). Music and the wellbeing of people with dementia. *Ageing and Society*, 27, 127-145. DOI: 10.1017/S0144686X06005228
- Skingley, A., & Vella-Burrows, T. (2010). Therapeutic effects of music and singing for older people. *Nursing Standard*, 24(19), 35-41.
- Skingley, A., Clift, S. M., Coulton, S. P., & Rodriguez, J. (2011). The effectiveness and cost-effectiveness of a participative community singing programme as a health promotion initiative for older people: Protocol for a randomised controlled trial. *BioMedCentral Public Health*, 11, 142.
- Sluming, V., Brooks, J., Howard, M., Downes, J. J., & Roberts, N. (2007). Broca's area supports enhanced visuospatial cognition in orchestral musicians. *The Journal of Neuroscience*, 27(14), 3799-3806. DOI: 10.1523/JNEUROSCI.0147-07.2007
- \* Smithrim, K. L. (1998). Still singing for our lives: Singing in the everyday lives of women through this century. In B. A. Roberts (Ed.), *Sharing the voices: The phenomenon of singing. Proceedings of the International Symposium*. St. John's, Newfoundland, Canada: Memorial University of Newfoundland.
- Sonke, J., Rollins, J., Brandman, R., & Graham-Pole, J. (2009). The state of the arts in healthcare in the United States. *Arts & Health: An International Journal for Research, Policy and Practice*, 1(2), 107-135. DOI: 10.1080/17533010903031580
- \* Southcott, J. E. (2009). 'And as I go, I love to sing': the Happy Wanderers, music and positive aging. *International Journal of Community Music*, 2(2&3), 143-156. DOI: 10.1386/ijcm.2.2&3.143/1
- Staricoff, R. L. (2004). *Arts in health: A review of the medical literature*. London: Arts Council England.
- Steurer, M., Simak, S., Denk, D. M., & Kautzky, M. (1998). Does choir singing cause noise-induced hearing loss? *Audiology*, 37(1), 38-51.
- Svansdottir, H. B., & Snaedal, J. (2006). Music therapy in moderate and severe dementia of Alzheimer's type: A case-control study. *International Psychogeriatrics*, 18(4), 613-621. DOI: 10.1017/S1041610206003206



- \* Tonneijck, H. I. M., Kinébanian, A., & Josephsson, S. (2008). An exploration of choir singing: Achieving wholeness through challenge. *Journal of Occupational Science, 15*(3), 173-180. DOI: 10.1080/14427591.2008.9686627
- \* Tsugawa, Samuel (2009). *Senior adult music learning, motivation, and meaning construction in two New Horizons ensembles* (Doctoral dissertation, Arizona State University).
- Unwin, M. M., Kenny, D. T., & Davis, P. J. (2002). The effects of group singing on mood. *Psychology of Music, 30*, 175-185.
- Valentine, E., & Evans, C. (2001). The effects of solo singing, choral singing and swimming on mood and physiological indices. *British Journal of Medical Psychology, 74*, 115-120.
- van Wijck, F., Knox, D., Dodds, C., Cassidy, G., Alexander, G., & MacDonald, R. A. R. (2011). Making music after stroke: Using musical activities to enhance arm function. *Annals Of The New York Academy Of Sciences, The Neurosciences and Music IV: Learning and Memory, 1252*, 305-311. DOI: 10.1111/j.1749-6632.2011.06403.x
- VanWeelden, K., & Cevasco, A. M. (2009). Geriatric clients' preferences for specific popular songs to use during singing activities. *Journal of Music Therapy, 46*(2), 147-159.
- \* Varvarigou, M., Creech, A., Hallam, S., & McQueen, H. (2011). Bringing different generations together in music-making – An intergenerational music project in East London. *International Journal of Community Music, 4*(3), 207-220. DOI: 10.1386/ijcm.4.3.207\_1
- \* Varvarigou, M., Hallam, S., Creech, A., & McQueen, H. (2012). Benefits experienced by older people who participated in group music-making activities. *Journal of Applied Arts and Health, 3*(2), 183-198.
- \* Varvarigou, M., Hallam, S., Creech, A., & McQueen, H. (2013). Different ways of experiencing music-making in later life: Creative music sessions for older learners in East London. *Research Studies in Music Education, 35*(1), 103-118. DOI: 10.1177/1321103X13478863
- Västfjäll, D., Juslin, P. N., & Hartig, T. (2012). Music, subjective wellbeing and health: The role of everyday emotions. In R. A. R. MacDonald, G. Kreutz, & L. Mitchell (Eds.), *Music, health, and wellbeing* (pp. 405-423). Oxford, UK: Oxford University Press.
- Vergheze, J., Lipton, R. B., Katz, M. J., Hall, C. B., Derby, C. A., Kuslansky, G.,...Buschk, H. (2003). Leisure activities and the risk of dementia in the elderly. *The New England Journal of Medicine, 348*, 2508-2516.
- \* von Lob, G., Camic, P., & Clift, S. (2010). The use of singing in a group as a response to adverse life events. *International Journal of Mental Health Promotion, 12*(3), 45-53. DOI: 10.1080/14623730.2010.9721818
- Wade, L. M. (2002). A comparison of the effects of vocal exercises/ singing versus music-assisted relaxation on peak expiratory flow rates of children with asthma. *Music Therapy Perspectives, 20*, 31-37.
- Young, L. (2008). The potential health benefits of community based singing groups for adults with cancer. *Canadian Journal of Music Therapy ∞ Revue canadienne de musicothérapie, 15*(1), 11-27.

Zelazny, C. M. (2001). Therapeutic instrumental music playing in hand rehabilitation for older adults with osteoarthritis: Four case studies. *Journal of Music Therapy, 38*(2), 97-113.

Table 1.  
*Works Reporting on Correlates of Music Participation*

Category	Correlate items included?		
	No	Yes	Total
Broad activity type			
Music making activities (involving at least voice and/or instruments)	62	68	130
Music listening	7	14	21
Leisure (e.g., participatory arts programs, attending cultural events)	9	2	11
Other (e.g., dancing, music therapy, composition, etc.)	13	13	26
Not clearly specified	14	0	14
<b>Total</b>	<b>105</b>	<b>97</b>	<b>202</b>
Type of music-making activity <sup>a</sup>			
Voice only	26	37	63
Instrument only	15	12	27
Mixed voice and instrument (or where the activity was broadly stated as music-making without further details)	7	12	19
Music making with additional activities in conjunction (such as activities that included making music in addition to therapy or dancing or listening)	19	7	26
<b>Total</b>	<b>67</b>	<b>68</b>	<b>135</b>

<sup>a</sup> Given the number of studies concerning music-making activities, we divided this category into four specific subgroupings.

Table 2.  
*Details of the Principal Axis Factor Analysis Concerning the Measure Items*

Item	Factor <sup>a</sup>				
	1	2	3	4	5
I feel uplifted.	0.92				
It improves/ enhances my mood.	0.88				
It makes me feel better.	0.77				
It helps me cope with stresses and everyday problems.	0.71				
It relieves stress/tension.	0.70				
It helps my psychological well-being.	0.67				
It puts me on a high.	0.64				
It gives me a peak experience.	0.54				
It gives me a sense of inner contentment.	0.52				
It offers me spiritual growth and healing.	0.43			0.41	
It adds meaning/ purpose to my life.	0.42				
It does (not) help me feel more positive about life.		0.85			
It does (not) give me a sense of belonging.		0.70	0.37		
It does (not) give me happiness.		0.62			
It does (not) enhance my quality of life.		0.60			
It does (not) give me a sense of playing a valued and vital role within a community.		0.60			
It does (not) make me feel connected to others.		0.51	0.46		
I do (not) feel connected to my audience.		0.43			0.33
It does (not) help keep me healthy.		0.42			
It does (not) increase my self-esteem.		0.42			
It does (not) allow me to be creative.		0.40			0.31
It does (not) help me feel more positive about myself.	0.34	0.39			
I develop friendships.			0.84		
I develop bonds with other people.			0.80		

It provides a feeling of community.				0.74	
It connects me to the community.	0.30			0.71	
I have positive feelings about other people.				0.49	
It does (not) help me to think about who I am.					0.70
It does (not) help me learn about myself.					0.63
It does (not) help me better understand my thoughts and emotions.					0.62
It allows me to communicate non-verbally.					0.43
It does (not) help me with my memory recall.	0.31				0.33
It does (not) push me to achieve my highest potential.					0.70
It does (not) give me a social group identity.			0.41		0.45
I do (not) experience self-improvement.					0.41
It does (not) generate/ increase my self confidence.					0.38
It encourages me to look after my physical health.					
It contributes/ improves my physical health.					
It does (not) improve my motor coordination.					
It gives me control over my life.					
Eigenvalue	13.02	3.14	1.92	1.44	0.84
% of variance	32.56	7.84	4.80	3.61	2.11
Cronbach's alpha	0.90	0.89	0.87	0.75	0.79

*Note.* Factor loadings < .30 are not shown.

<sup>a</sup> Factors 1-5 were labeled as “mood and coping,” “esteem and worth,” “socializing,” “cognitive,” and “self-actualization” respectively.