

## Two-electron interference in angular resolved double photoionization of Mg

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2014 J. Phys.: Conf. Ser. 488 022006

(<http://iopscience.iop.org/1742-6596/488/2/022006>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 134.7.93.129

This content was downloaded on 08/04/2016 at 04:13

Please note that [terms and conditions apply](#).

## Two-electron interference in angular resolved double photoionization of Mg

E. Sokell\*, P. Bolognesi†, A. Kheifets‡, Igor Bray#, S Safgren\* and L.Avaldi†<sup>1</sup>

\* School of Physics, UCD Science Centre, Belfield, Dublin 4, Ireland

† CNR-IMIP, Area della Ricerca di Roma 1, 00015 Monterotondo Scalo, Italy

‡ RSPE, The Australian National University, Canberra ACT 0200, Australia

# ARC Centre for Matter-Antimatter Studies, Curtin University, WA 6845 Perth, Australia


**Synopsis** We observe a two-electron interference effect in the symmetrized amplitude of the resonant double photoionization of Mg which was predicted theoretically, but never before observed experimentally. This observation is based on our experimental study of angle-resolved double photoionization of Mg at the photon energy of 55.49 eV ( $2p \rightarrow 3d$  resonance) under equal and unequal energy sharing conditions.

Over the past 20 years, investigation of double photoionization (DPI) in He had challenged both experimentalists and theorists. It was established that correlated dynamics of the two-electron escape is strongly constrained by its own symmetry and the Coulomb repulsion. Alkaline-earth-metal atoms are “quasi two-electron” systems and represent the most suitable candidates for extending the investigation of DPI beyond He.

We have undertaken an investigation of DPI in Mg and measured the triply-differential cross-section (TDCS) at 55.49 eV (the excitation of the  $2p \rightarrow 3d$  resonance) under both equal and unequal energy sharing conditions. Under the unequal energy sharing, the two complementary kinematics ( $E_1 \leftrightarrow E_2 = 10.4 \leftrightarrow 22.4$  eV) have been investigated. The experiments have been performed with the multi-coincidence endstation of the Gas Phase beamline at Elettra. The results have been compared with theoretical calculations using the CCC formalism, where the effect of the resonant excitation was incorporated semi-empirically [1].

From the experimental TDCS, the gerade and ungerade amplitudes have been extracted [2] and compared with the theoretical predictions. The main result [1] is the observation of a gerade amplitude in the equal energy sharing case which deviates strongly from a Gaussian shape. The whole of the amplitude can be better described by the di-Gaussian parameterization proposed in [3] and associated with the shape in the coordinate space of the  $Mg^+ 3s$  orbital (Fig. 1). The two Gaussians are associated with two main areas in the target coordinate space. Conversely, the  $He^+ 1s$  radial orbital has only one area of charge localization and thus, the DPI amplitude is represented by a single Gaussian. This is the first observation of a strong modification of the angular correlation pattern in the

DPI of a  $ns$ -orbital with respect to the one of He. The origin is in the structure of the initial state wavefunction and can be interpreted as a two-electron interference [3,1]. Similar to the electron “two-slit” experiments, where the wave functions of the electrons, “emitted” from different spatial positions, interfere, here the wave functions of two-electron pairs “originated” from two different regions of the charge density interfere.



**Figure 1.** The gerade amplitudes of DPI of Mg (top) and He (middle) at  $E_1 = E_2$  versus the mutual photoelectron angle  $\theta_{12}$ . The CCC calculation (red solid line) is fitted with the Gaussian and di-Gaussian parameterization (blue and green dashed lines). The experimental amplitude is shown with error bars. Bottom: the radial orbitals for  $Mg^+ 3s$  and  $He^+ 1s$  (red solid and green dashed lines, respectively)

### References

- [1] E Sokell *et al* 2013 *Phys. Rev. Lett.* **110** 083001
- [2] P Bolognesi *et al* 2003 *J. Phys. B* **36** L241
- [3] A S Kheifets *et al* 2011 *J. Phys. B* **44** 011002

<sup>†1</sup> E-mail: [Lorenzo.Avaldi@imip.cnr.it](mailto:Lorenzo.Avaldi@imip.cnr.it)

