

NOTICE: this is the author's version of a work that was accepted for publication in Organic Geochemistry. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Organic Geochemistry, Vol.61, 2013.  
DOI: 10.1016/j.orggeochem.2013.05.006

1 **A pyrolysis and stable isotopic approach to investigate the**  
2 **origin of methyltrimethyltridecylchromans (MTTCs)**

3 **Svenja Tulipani<sup>a\*</sup>, Kliti Grice<sup>a\*</sup>, Paul Greenwood<sup>a,b</sup>, Lorenz Schwark<sup>a,c</sup>**

4 *<sup>a</sup> WA Organic and Isotope Geochemistry Centre, Department of Chemistry, Curtin*  
5 *University, GPO Box U1987, Perth, WA 6845, Australia.*

6 *<sup>b</sup> Centre for Exploration Targeting and WA Biogeochemistry Centre, The*  
7 *University of Western Australia 35 Stirling Highway, Crawley WA 6009*  
8 *Australia*

9 *<sup>c</sup> Institute of Geoscience, Kiel University, Ludewig-Meyn Str. 10, 24118 Kiel,*  
10 *Germany.*

11 \*Corresponding author: email: [s.tulipani@curtin.edu.au](mailto:s.tulipani@curtin.edu.au); [k.grice@curtin.edu.au](mailto:k.grice@curtin.edu.au)

12 tel: +61 (0)8 9266 2474

13 **Abstract**

14 Methyltrimethyltridecylchromans (MTTCs) have been widely detected in  
15 sediments and crude oils from various depositional settings and are established  
16 markers for palaeosalinities. A likely origin of these compounds, which show a  
17 distinctive isoprenoid substituted aromatic structure, seems to be condensation  
18 reactions of phytol with higher plant-derived alkyl phenols during early  
19 diagenesis. However, a direct biological origin from phytoplanktonic organisms  
20 cannot be excluded. To further investigate the potential origin from condensation  
21 reactions, an online pyrolysis-gas chromatography- isotope ratio mass  
22 spectrometry (PY-GC-irMS) method with the capacity to measure  $\delta^{13}\text{C}$  in  
23 fragments (trimethylphenol and pristenes) generated from 5,7,8-trimethyl-  
24 MTTC was developed in this study. This straight forward technique poses a

25 great potential for the elucidation of chroman formation in geological samples as  
26 it possibly enables the distinction between the different proposed sources of  
27 isoprenoid and alkyl-phenol fragments (mainly phytoplankton and higher plants,  
28 respectively) based on their stable isotopic compositions. Furthermore, it might  
29 be useful for the investigation of products generated from MTTCs during  
30 thermal maturation of geological samples.

31 **Keywords:** Flash- pyrolysis, CSIA, palaeosalinity, phenols

## 32 **1. Introduction**

33 Methylated 2-methyl-2-trimethyltridecylchromans (MTTCs, **I** in **Figure 1**) in  
34 sediments or crude oils generally occur as distinct isomers of monomethyl,  
35 dimethyl and trimethyl homologues. They were first identified in geological  
36 samples by Sinninghe-Damsté et al. (1987), who also introduced them as  
37 palaeosalinity indicators. MTTCs have since been reported in a great variety of  
38 geological samples and the “chroman ratio” (5,7,8-trimethyl MTTC/total MTTCs)  
39 has been established as a powerful tool in salinity reconstructions (e.g. Schwark  
40 and Püttmann, 1990; Sinninghe Damsté et al., 1993; Grice et al., 1998; Schwark  
41 et al., 1998). However, their origin and geological formation pathway remain  
42 debated (Sinninghe Damsté et al., 1993; Li and Larter, 1995; Li et al., 1995,  
43 Sinninghe-Damsté and De Leeuw, 1995). Based on correlation of abundances  
44 and chroman ratios with other geological parameters and as an explanation for  
45 the limited number of naturally occurring isomers, a biosynthetic origin of  
46 MTTCs from phytoplanktonic organisms has been suggested (e.g. Sinninghe  
47 Damsté et al., 1993), although to date MTTCs or suitable direct precursors have  
48 not been found in organisms. An origin from higher plant tocopherols (**II**, **Figure**

1), which bear a strong structural similarity, has been ruled out due to their comparatively low abundances in the geosphere and the presence of a phenolic hydroxyl group at C-6 (Sinninghe Damsté et al., 1993; Li et al., 1995). Li et al. (1995) alternatively proposed that MTTCs might form via early diagenetic condensation reactions of the phytol side chain in chlorophylls with higher plant derived phenols, which would imply largely different source organisms for the isoprenoid and alkylphenol moiety of geological chromans. To further investigate this potential formation pathway, we developed a pyrolysis-stable isotope analytical method for  $\delta^{13}\text{C}$  determination in isoprenoid and alkylphenol fragments generated from MTTCs, which could possibly be used to establish the relationship to the different proposed source organisms of these fragments on a stable isotopic basis. Furthermore, tocopherols and MTTCs have been suggested as an additional source of pristane in more mature sediments/crude oils (Goossens et al., 1984; Li et al., 1995), which could also possibly be explored with this technique. The method was initially investigated by thermal degradation of an authentic 5,7,8-trimethyl-MTTC (triMeMTTC) standard in order to establish the stable isotopic relationship between the parent compound and the distinctive degradation products. Subsequently, chroman isolates from three Middle to Upper Devonian sediments (Canning Basin, WA) were analysed to demonstrate the applicability of the method in natural samples. Although pyrolysis products of natural and artificial MTTCs and related compounds have been thoroughly investigated by Li et al. (1995), there have been no previous isotopic based studies of these compounds to establish the formation mechanism of MTTCs.

## 72 2. Experimental

73 The authentic 5,7,8-trimethyl MTTC standard was synthesised from 2,3,5-  
74 trimethylphenol and phytol according to Sinninghe-Damsté et al. (1987).

75 Sediments (MWR-30.7 m, MWR-40.7 m and MWR-41.2 m) with high triMeMTTC  
76 abundances and exceptionally low maturities (e.g.  $T_{\max}$  405–413 °C; unpublished  
77 data) originated from basin facies associated with Middle to Upper Devonian reef  
78 systems in the Canning Basin, Western Australia. The powdered rock was  
79 Soxhlet extracted and the total lipid extract fractionated by silica gel column  
80 chromatography (for details see Grice et al., 2005; Supplementary material).

81 Unsaturated compounds were separated from the aliphatic fraction by  $\text{AgNO}_3$   
82 silica column chromatography (10%) using hexane (saturated compounds) and  
83 DCM (unsaturated compounds) as eluents. *n*-Alkanes were subsequently  
84 removed with ZSM5 molecular sieve (e.g. Audino et al., 2001) to obtain a  
85 branched and cyclic fraction. 5,7,8-trimethyl MTTC was further isolated from the  
86 aromatic fractions of MWR-40.7 m and MWR-41.2 m by  $\text{AgNO}_3$  thin layer  
87 chromatography (Eglinton and Murphy, 1969) using hexane as developer and the  
88 authentic chroman standard (visualized with rhodamine spray under UV-light)  
89 as reference. The MTTC containing silica band was scraped off, extracted with  
90 DCM and filtered through a glass sinter funnel under vacuum. The low amount  
91 of aromatic compounds in the MWR-30.7 m sample precluded a TLC isolate  
92 being obtained.

93 For bulk  $\delta^{13}\text{C}$  analysis a Delta V Plus mass spectrometer connected to a Thermo  
94 Flush 1112 via Conflow IV (Thermo-Finnigan/Germany) was used. Analytes  
95 were combusted at 1020 °C.

96 Gas chromatography-mass spectrometry (GC-MS) was performed on an Agilent  
97 5973 GC-MS equipped with a HP 6890 auto-sampler and a DB-5MS capillary  
98 column. The GC oven was heated from 40–310 °C or 325 °C at 3 °C/min with  
99 initial and final hold times of 1 min and 30 min, respectively. A CDS 5350 Auto-  
100 pyroprobe was used for flash pyrolysis (PY)-GC-MS. The pyrolysis chamber and  
101 injector were held at 300 °C and pyrolysis was separately performed at  
102 temperatures of 550 °C, 650 °C or 750 °C applied for 20 s. The pyrolysates were  
103 analysed with a 60:1 split. He carrier gas at a constant pressure of 17.5 psi was  
104 used and the GC oven was temperature programmed from -20 °C to 40 °C at 8  
105 °C/min and then to 320 °C at 4 °C/min with initial and final hold times of 1 and  
106 25 min, respectively. All other settings remained unchanged.

107 Gas chromatography-isotope ratio mass spectrometry (GC-irMS) was performed  
108 on a Micromass IsoPrime irMS interfaced to an Agilent 6890N GC fitted with a  
109 HP 7683 autosampler. GC parameters were similar to those used for GC-MS. For  
110 PY-GC-irMS a CDS-Pyroprobe 5000 was mounted directly on the vaporisation  
111 injector of the GC-irMS system. The pyrolysis chamber and injector were set to  
112 300 °C. Analytes were pyrolysed at 650 °C for 20 s, injected with a 30:1 split or  
113 splitless (for increased sensitivity) and trapped in liquid nitrogen until the end of  
114 pyrolysis. The GC oven was programmed from 40–325 °C at 4 °C/min with initial  
115 and final hold times of 2 and 15 min, respectively. GC column and all other  
116 settings remained unchanged. Reference standards of known isotopic  
117 composition were regularly analysed to confirm accuracy of isotope analysis. All  
118  $\delta^{13}\text{C}$  values reported in this study are the average of at least two replicates and  
119 standard deviations were reported.

120 Further details about typical injector, carrier gas and MS/irMS settings as well  
121 as GC column, interface (for GC-irMS) and instrument software used for GC-  
122 MS/irMS can be found in Supplementary materials of Melendez et al. (2013).

### 123 **3. Results and discussion**

124 The aim of this study was to develop an online flash pyrolysis-GC-irMS method  
125 which would allow stable isotopic correlation of MTTCs and related lower  
126 molecular weight products for the elucidation of their sources and formation  
127 pathways in geological samples. An authentic 5,7,8-triMeMTTC standard (often  
128 the most abundant natural chroman) was first analysed by PY-GC-MS to  
129 identify major degradation products of the parent structure and investigate  
130 pyrolysis efficiency at different temperatures (550 °C, 650 °C and 750 °C) in  
131 separate pyrolysis experiments. The major pyrolysates in all analyses were 2,3,5-  
132 trimethylphenol (see appendix for compound identification) and pristenes as well  
133 as the intact chroman (e.g. **Figure 2a**). The extent of pyrolytic degradation was  
134 inferred from the ratio between the abundance (peak area) of the  
135 trimethylphenol and all pristene products relative to the original chroman in  
136 four replicate analyses. The highest degradation efficiency was achieved at a  
137 pyrolysis temperature of 650 °C (ratios of 0.8, 1.6, 1.2 for 550 °C, 650 °C and 750  
138 °C, respectively), which therefore was used in all subsequent analyses. However,  
139 the replicates generally showed some variability which is typical of many  
140 analytical pyrolysis studies. Li et al. (1995) conducted offline pyrolysis over 65 h  
141 at 350 °C on isolates of 5,7,8-triMeMTTC which similarly showed high amounts  
142 of pristenes, but contrary to present data generated tetramethylphenol instead of  
143 trimethylphenol. This was also the main product we generated in preliminary

144 and unpublished pyrolysis experiments of the 5,7,8-triMeMTTC in sealed glass  
145 tubes at temperatures of 330 and 360 °C over 72h. The different product  
146 obtained from flash pyrolysis may be the result of the elevated pyrolysis  
147 temperatures leading to a different bond cleavage in the chroman. In an earlier  
148 study, tocopherols have also been shown to generate significant amounts of  
149 pristenes during pyrolysis and have therefore been suggested as a contributor to  
150 pristane in geological samples (Goossens et al., 1984).

151 Precision and accuracy of  $\delta^{13}\text{C}$  values measured by PY-GC-irMS were tested with  
152 five replicate analyses of the 5,7,8-triMeMTTC standard (using split and  
153 splitless injections). Standard deviations between 0.2‰ and 0.4‰ for all  
154 measured compounds confirmed an excellent precision (Table 1).  $\delta^{13}\text{C}$  values  
155 reported for prist-1-ene include a coeluting pristene isomer (*cf.* Figure 2a and  
156 Figure 3a, b). Apart from that, good baseline separations, essential for GC-irMS  
157 analysis, were achieved for all remaining products.  $\delta^{13}\text{C}$  values of  
158 trimethylphenol, pristenes and triMeMTTC in pyrolysates were comparable to  
159 reference values obtained by elemental analysis (EA)-irMS of the chroman  
160 standard as well as the phytol and trimethylphenol utilised for its synthesis  
161 (Table 2). This confirmed both the accuracy of the data and the preservation of  
162 the  $\delta^{13}\text{C}$  signature of source compounds during condensation reactions and  
163 pyrolysis. The slight systematic depletion of  $\delta^{13}\text{C}$  values obtained from EA-irMS  
164 in comparison to corresponding values measured on the GC-irMS system (in  
165 pyrolysis products as well as in the chroman standard analysed by conventional  
166 liquid injection; Table 2) can be attributed to instrumental bias. Similar  
167 systematic variations between different systems for EA- and GC-irMS have



168 previously been reported (e.g. Zwank et al., 2003). Nevertheless, values obtained  
169 from both methods are in accordance with the following mass balance equation:

170 
$$\delta^{13}\text{C}_{\text{triMeMTTC}} = \frac{9 \times \delta^{13}\text{C}_{\text{trimethylphenol}} + 20 \times \delta^{13}\text{C}_{\text{phytol/pristenes}}}{29}$$

171 where “ $\delta^{13}\text{C}_{\text{phytol/pristenes}}$ ” stands for  $\delta^{13}\text{C}$  of phytol (for bulk-irMS) or average  $\delta^{13}\text{C}$   
172 of all pristenes (for PY-GC-irMS). The calculated  $\delta^{13}\text{C}$  values for triMeMTTC of -  
173 32.5‰ and -33.3‰ for PY-GC-irMS and EA-irMS, respectively, are almost  
174 identical to the measured values (Table 1 and 2).

175 PY-GC-irMS was applied to the TLC-isolates from MWR-40.7 m and MWR-41.2  
176 m (Figure 2c) and the whole aromatic fraction of MWR-30.7 m (containing  
177 abundant triMeMTTC – **Figure 2b**). Figure 3c shows a typical GC-irMS trace of  
178 pyrolysates obtained from these samples. Notable differences to the pyrolysate  
179 distribution of the chroman standard include the absence of trimethylphenol and  
180 prist-2-ene, which can probably be attributed to matrix effects, i.e. other  
181 compounds present in the TLC isolates/aromatic fraction influencing thermal  
182 behaviour, which can alter flash pyrolysis product distributions (e.g. Greenwood  
183 et al., 2006). Further optimisation of the pyrolysis conditions for the challenges  
184 of geological samples would be useful, but was not done here due to the limited  
185 quantity of these samples. The  $\delta^{13}\text{C}$  values of pristene (most likely prist-1-ene  
186 and a second co-eluting isomer) measured by PY-GC-irMS of the three samples  
187 was consistently similar to the corresponding values of pristane and phytane  
188 obtained from traditional liquid injection GC-irMS. This correlation strongly  
189 points to a common source for these products, most likely the phytol side chain in  
190 chlorophylls (Table 3). Furthermore, the traditionally measured  $\delta^{13}\text{C}$  values of

191 triMeMTTC were also similar to the isotopic signatures of these products,  
192 although a very small  $^{13}\text{C}$  enrichment was notable (Table 3). However, since the  
193  $\delta^{13}\text{C}$  value of the alkylphenol moiety of the chroman standard could not be  
194 measured, the suggested formation of chromans by biosynthesis in  
195 phytoplanktonic organisms (Sinninghe Damsté et al., 1993) cannot be discounted  
196 based on these results.

#### 197 **4. Conclusions and outlook**

198 An online PY-GC-irMS method which enables  $\delta^{13}\text{C}$  analysis of major thermal  
199 breakdown products of triMe-MTTC (trimethylphenol and pristenes) was  
200 developed. Initial application to a triMeMTTC standard confirmed high precision  
201 and accuracy of the  $\delta^{13}\text{C}$  data. Furthermore, the isotopic relationship of major  
202 pyrolysis products to the parent chroman as well as to the corresponding source  
203 compounds used for synthesis of the standard was established. Similar analyses  
204 of triMeMTTC in isolates from immature sediments also generated a pristene  
205 peak, however, trimethylphenol and prist-2-ene, which were obtained from the  
206 standard in the previous analyses, were lacking. A more complete suite of MTTC  
207 pyrolysis markers should be achievable with further optimisation of pyrolysis  
208 conditions. Nevertheless, the few MTTC products detected in these initial  
209 analyses of geological material show a great potential for the application of this  
210 analytical method to probe the origin of MTTCs in geological samples.

#### 211 **Acknowledgements**

212 We would like to acknowledge Geoff Chidlow and Steven Clayton for technical  
213 support. Curtin University is thanked for providing a Curtin Strategic  
214 International Research Scholarship (CSIRS) to S.T. The Institute for Geoscience

215 Researcher (TiGER) is thanked for a top-up scholarship. K.G. acknowledges the  
216 Australian Research Council (ARC) for a Discovery QEII grant supporting this  
217 research, and the John de Laeter Centre for Isotope Research and ARC LIEF  
218 grants which provided the infrastructure support. Chevron (Ted Playton), the  
219 Geological Survey of Western Australia and the Minerals and Energy Research  
220 Institute of Western Australia (M396) are thanked for sponsoring an ARC  
221 Linkage grant (LP0883812) which supported the fieldwork and research.  
222 Wundargoodie, Mimbi Community and Mount Pierre Station are thanked for  
223 supporting the field work. All authors would like to thank Dr. Maowen Li, an  
224 anonymous reviewer and Associate Editor Prof. Simon George for constructive  
225 and helpful comments.

## 226 **Figure captions**

227 **Figure 1.** Chemical structures referred to in the text

228 **Figure 2.** Total ion chromatograms of a typical pyrolysate obtained from an  
229 authentic 5,7,8-trimethyl-2-methyl-2-trimethyltridecylchroman (triMeMTTC)  
230 standard (a), and triMeMTTC in isolates from natural samples (MWR-30.7 m,  
231 aromatic fraction; b, and MWR-40.7 m thin layer chromatography isolate from  
232 aromatic fraction (c), which were pyrolysed in subsequent experiments. TMP =  
233 trimethylphenol; \* = Impurities in triMeMTTC standard.

234 **Figure 3.** Pyrolysis-gas chromatography-isotope ratio mass spectrometry (PY-  
235 GC-irMS) chromatograms of authentic 5,7,8-trimethyl-2-methyl-2-  
236 trimethyltridecylchroman (triMeMTTC) standard (a and b) and MTTC-isolate  
237 from the MWR-40.7 m natural sample (c).

## 238 **Table captions**

239 Table 1.  $\delta^{13}\text{C}$  values of compounds in the pyrolysate obtained from five replicate  
 240 analyses of an authentic chroman standard including average  $\delta^{13}\text{C}$  values  $\pm$   
 241 standard deviation. <sup>a</sup> injection with 30:1 split; <sup>b</sup> splitless injection; \* joined peak  
 242 of prist-1-ene and second, less abundant pristene isomer

	$\delta^{13}\text{C}$ [‰ VPDB]					Average
	run 1 <sup>a</sup>	run 2 <sup>a</sup>	run 3 <sup>b</sup>	run 4 <sup>b</sup>	run 5 <sup>b</sup>	
<b>trimethylphenol</b>	-29.6	-29.9	-29.7	-29.4	-29.6	-29.6 $\pm$ 0.2
<b>prist-1-ene*</b>	-33.5	-34.2	-34.1	-33.7	-33.9	-33.9 $\pm$ 0.3
<b>prist-2-ene</b>	-33.4	-34.2	-33.9	-33.4	-34.0	-33.8 $\pm$ 0.4
<b>5,7,8-triMeMTTC</b>	-32.2	-32.8	n.d.	n.d.	n.d.	-32.5 $\pm$ 0.4

243

244 Table 2.  $\delta^{13}\text{C}$  values of the synthesized chroman standard and source compounds  
 245  $\pm$  standard deviations between 3 (a) or 2 (b) replicates obtained by elemental  
 246 analysis-isotope ratio mass spectrometry (EA-irMS) and gas chromatography  
 247 (GC)-irMS. n.d. = not determined

	$\delta^{13}\text{C}$ [‰ VPDB]		
	2,3,5-trimethylphenol	phytol	5,7,8-triMeMTTC
<b>EA-irMS</b>	-30.6 $\pm$ 0.0 <sup>a</sup>	-34.5 $\pm$ 0.0 <sup>b</sup>	-33.4 $\pm$ 0.2 <sup>b</sup>
<b>GC-irMS</b>	n.d.	n.d.	-32.9 $\pm$ 0.1 <sup>a</sup>

248

249 Table 3.  $\delta^{13}\text{C}$  [‰ VPDB] of selected hydrocarbons in the aliphatic and aromatic  
 250 fractions as well as pristenes generated by flash pyrolysis of the aromatic  
 251 fraction (a) or isolated chroman (b)  $\pm$  standard deviation of 2 replicate  
 252 measurements. \*Only measured once due to limited sample material.

Sample i.d.	GC-irMS			Py-GC-irMS
	pristane	phytane	5,7,8-triMeMTTC	Pristenes
<b>MWR-30.7 m</b>	-31.3 $\pm$ 0.2	-29.9 $\pm$ 0.4	n.d.	-31.2a*
<b>MWR-40.7 m</b>	-33.2 $\pm$ 0.1	-32.9 $\pm$ 0.4	-32.7 $\pm$ 0.2	-33.0 $\pm$ 0.1
<b>MWR-41.2 m</b>	-32.7 $\pm$ 0.1	-32.6 $\pm$ 0.0	-32.1 $\pm$ 0.0	-32.4 $\pm$ 0.0

253

254 **References**

- 255 Alexander, R., Berwick, L., Pierce, K., 2011. Single carbon surface reactions of 1-  
256 octadecene and 2,3,6-trimethylphenol on activated carbon: Implications for  
257 methane formation in sediments. *Organic Geochemistry* 42, 540-547.
- 258 Audino, M., Grice, K., Alexander, R., Boreham, C.J., Kagi, R.I., 2001. Unusual  
259 distribution of monomethylalkanes in *Botryococcus braunii*-rich samples:  
260 Origin and significance. *Geochimica et Cosmochimica Acta* 65, 1995-2006.
- 261 Bastow, T.P., van Aarssen, B.G.K., Alexander, R., Kagi, R.I., 2005. Origins of  
262 alkylphenols in crude oils: Hydroxylation of alkylbenzenes. *Organic*  
263 *Geochemistry* 36, 991-1001.
- 264 Eglinton, G., Murphy, M.T.J., 1969. *Organic Geochemistry: Methods and*  
265 *Results*. Springer-Verlag, Berlin.
- 266 Goossens, H., de Leeuw, J.W., Schenck, P.A., Brassell, S.C., 1984. Tocopherols as  
267 likely precursors of pristane in ancient sediments and crude oils. *Nature* 312,  
268 440-442.
- 269 Greenwood, P.F., Leenheer, J.A., McIntyre, C., Berwick, L., Franzmann, P.D.,  
270 2006. Bacterial biomarkers thermally released from dissolved organic matter.  
271 *Organic Geochemistry* 37, 597-609.
- 272 Grice, K., Cao, C., Love, G.D., Böttcher, M.E., Twitchett, R.J., Grosjean, E.,  
273 Summons, R.E., Turgeon, S.C., Dunning, W., Jin, Y., 2005. Photic zone euxinia  
274 during the Permian-Triassic superanoxic event. *Science* 307, 706-709.
- 275 Grice, K., Schouten, S., Nissenbaum, A., Charrach, J., Sinninghe Damsté, J.S.,  
276 1998. A remarkable paradox: Sulfurised freshwater algal (*Botryococcus braunii*)

277 lipids in an ancient hypersaline euxinic ecosystem. *Organic Geochemistry* 28,  
278 195-216.

279 Li, M., Larter, S.R., 1995. Reply to comments by Sinninghe Damsté and De  
280 Leeuw (1995) on Li et al. (1995), *Organic Geochemistry* 23, 159–167. *Organic*  
281 *Geochemistry* 23, 1089-1093.

282 Li, M., Larter, S.R., Taylor, P., Jones, D.M., Bowler, B., Bjorøy, M., 1995.  
283 Biomarkers or not biomarkers? A new hypothesis for the origin of pristane  
284 involving derivation from methyltrimethyltridecylchromans (MTTCs) formed  
285 during diagenesis from chlorophyll and alkylphenols. *Organic Geochemistry* 23,  
286 159-167.

287 Melendez, I., Grice, K., Trinajstic, K., Ladjavardi, M., Greenwood, P., Thompson,  
288 K., 2013. Biomarkers reveal the role of photic zone euxinia in exceptional fossil  
289 preservation: An organic geochemical perspective. *Geology*, 41 123-126.

290 Schwark, L., Püttmann, W., 1990. Aromatic hydrocarbon composition of the  
291 Permian Kupferschiefer in the Lower Rhine Basin, NW Germany. *Organic*  
292 *Geochemistry* 16, 749-761.

293 Schwark, L., Vliex, M., Schaeffer, P., 1998. Geochemical characterization of  
294 Malm Zeta laminated carbonates from the Franconian Alb, SW-Germany (II).  
295 *Organic Geochemistry* 29, 1921-1952.


296 Sinninghe-Damsté, J.S., De Leeuw, J.W., 1995. Comments on “Biomarkers or not  
297 biomarkers. A new hypothesis for the origin of pristane involving derivation  
298 from methyltrimethyltridecylchromans (MTTCs) formed during diagenesis from  
299 chlorophyll and alkylphenols” from M. Li, S.R. Larter, P. Taylor, D.M. Jones, B.  
300 Bowler and M. Bjorøy. *Organic Geochemistry* 23, 1085-1087.

301 Sinninghe-Damsté, J.S., Kock-Van Dalen, A.C., De Leeuw, J.W., Schenck, P.A.,  
 302 Guoying, S., Brassell, S.C., 1987. The identification of mono-, di- and trimethyl  
 303 2-methyl-2-(4,8,12-trimethyltridecyl)chromans and their occurrence in the  
 304 geosphere. *Geochimica et Cosmochimica Acta* 51, 2393-2400.

305 Sinninghe Damsté, J.S., Keely, B.J., Betts, S.E., Baas, M., Maxwell, J.R., de  
 306 Leeuw, J.W., 1993. Variations in abundances and distributions of isoprenoid  
 307 chromans and long-chain alkylbenzenes in sediments of the Mulhouse Basin: A  
 308 molecular sedimentary record of palaeosalinity. *Organic Geochemistry* 20,  
 309 1201-1215.


310 Zwank, L., Berg, M., Schmidt, T.C., Haderlein, S.B., 2003. Compound-specific  
 311 carbon isotope analysis of volatile organic compounds in the low-microgram per  
 312 liter range. *Analytical Chemistry* 75, 5575-5583.

313 **Figures**




314


(a) Pyrolysate of triMeMTTC



(b) MWR-30.7 m




(c) MWR-40.7 m



315

*triMeMTTC standard (30:1 split)*



316  
317

## 318 Appendix

### 319 Identification of 2,3,5-trimethylphenol

320 The identity of 2,3,5-trimethylphenol (TMP) generated by pyrolysis of the 5,7,8  
321 triMe-MTTC standard was confirmed by the comparison of retention times with  
322 an authentic 2,3,5-TMP standard. For this purpose a CDS-Pyroprobe 5000 was



323 mounted directly on the vaporisation injector of the GC-MS system described in  
324 the experimental section. Except for increasing the initial hold time at -20 °C to  
325 2 min and the utilization of a different GC-column (ZB-5; Phenomenex) all GC-  
326 MS conditions were the same as described in the experimental section. Previous  
327 studies have shown that 2,3,5-TMP did not co-elute with other TMP isomers at  
328 comparable GC conditions (Bastow et al., 2005; Alexander et al., 2011), which  
329 enables an unequivocal identification of the generated TMP using this standard.  
330 The 5,7,8 triMe-MTTC standard was pyrolysed at 650 °C for 20s. For the  
331 analysis of the 2,3,5-TMP standard the pyrolysis chamber was kept at 300 °C for  
332 20s. Total ion chromatograms (TIC) of the 2,3,5-TMP standard and the MTTC  
333 pyrolysis product are displayed in Fig. A1. The mass spectrum of the TMP  
334 generated from MTTC pyrolysis is shown in Fig. A2.

### 335 **Figure Captions Appendix**

336 Figure A1:

337 Overlain TIC chromatograms of the 2,3,5-trimethylphenol (TMP) standard and  
338 the TMP in the pyrolysate of 5,7,8 trimethylmethyltrimethyltridecylchroman  
339 (triMeMTTC) analysed under the same GC-conditions confirming the identity of  
340 the latter

341


342 Figure A2:

343 Mass spectrum of the 2,3,5-trimethylphenol in the pyrolysate of 5,7,8  
344 trimethylmethyltrimethyltridecylchroman (MTTC)

345


### 346 **Figures Appendix**

347 **Figure A1**



348

349 **Figure A2**



350