

Public Health

Correlations between Control of COVID-19 Transmission and Influenza Occurrences in Malaysia

--Manuscript Draft--

Manuscript Number:	PUHE-D-21-01168R1
Article Type:	COVID-19 Original Research
Keywords:	COVID-19; influenza; SARS-CoV-2; movement control; physical distancing; sanitization
Corresponding Author:	Kuok Ho Daniel Tang, Ph.D. Beijing Normal University-Hong Kong Baptist University United International College Zhuhai, CHINA
First Author:	Kuok Ho Daniel Tang, Ph.D.
Order of Authors:	Kuok Ho Daniel Tang, Ph.D. Bridgid Lai Fui Chin
Abstract:	<p>Objectives: The transmission of COVID-19 has sent Malaysia into cycles of tightening and relaxation of movement control which are still continuing currently in line with local fluctuations of new COVID-19 cases. During movement control, measures comprising physical distancing, hand cleaning or sanitizing, and sanitization of premises are consistently implemented while self-isolation and travel restrictions are adaptively enforced. This study aims to examine if the control of COVID-19 transmission has an effect on the national influenza occurrences as some measures for COVID-19 control are similar to those for influenza.</p> <p>Study design: This study drew the data of weekly new cases of influenza and COVID-19 from official platforms for non-parametric statistical analysis.</p> <p>Methods: This study compared the influenza occurrences before and after the onset of COVID-19 using the Mann-Whitney U Test and explored the Spearman's correlations between COVID-19 and influenza incidences after the onset of COVID-19.</p> <p>Results: It shows that influenza incidences before and after the onset of COVID-19 were significantly different where influenza cases have significantly reduced after the onset of COVID-19. The weekly cases of influenza and COVID-19 were significantly and negatively correlated.</p> <p>Conclusions: This study underscores the co-benefits of COVID-19 control measures and alleviates the concern for the risk of COVID-19 and influenza coinfection.</p>

Correlations between Control of COVID-19 Transmission and Influenza Occurrences in Malaysia

Kuok Ho Daniel Tang^{1*}, Bridgid Lai Fui Chin

^{1*}Environmental Science Program, Division of Science and Technology, Beijing Normal University-Hong Kong Baptist University United International College, 2000 Jintong Road, Tangjiawan, Zhuhai, GD 519087, China. Email: daniel.tangkh@yahoo.com

²Department of Chemical Engineering, Faculty of Engineering and Science, Curtin University Malaysia, CDT 250, 98009 Miri, Sarawak, Malaysia

Correlations between Control of COVID-19 Transmission and Influenza Occurrences in Malaysia

Abstract

Objectives: The transmission of COVID-19 has sent Malaysia into cycles of tightening and relaxation of movement control which are still continuing currently in line with local fluctuations of new COVID-19 cases. During movement control, measures comprising physical distancing, hand cleaning or sanitizing, and sanitization of premises are consistently implemented while self-isolation and travel restrictions are adaptively enforced. This study aims to examine if the control of COVID-19 transmission has an effect on the national influenza occurrences as some measures for COVID-19 control are similar to those for influenza.

Study design: This study drew the data of weekly new cases of influenza and COVID-19 from official platforms for non-parametric statistical analysis.

Methods: This study compared the influenza occurrences before and after the onset of COVID-19 using the Mann-Whitney U Test and explored the Spearman's correlations between COVID-19 and influenza incidences after the onset of COVID-19.

Results: It shows that influenza incidences before and after the onset of COVID-19 were significantly different where influenza cases have significantly reduced after the onset of COVID-19. The weekly cases of influenza and COVID-19 were significantly and negatively correlated.

Conclusions: This study underscores the co-benefits of COVID-19 control measures and alleviates the concern for the risk of COVID-19 and influenza coinfection.

Keywords: COVID-19, Influenza, SARS-CoV-2, movement control, physical distancing, sanitization

Introduction

COVID-19 was first detected on 31 December 2019 in Wuhan, China and in less than a month, its arrival in Malaysia involving three Chinese citizens was reported on 25 January 2020 ⁽¹⁾. Malaysia recorded a total of 8 confirmed cases of COVID-19 on 30 January 2020, all involving Chinese citizens. On 4 February 2020, the first local infection was reported. Since 27 January 2020, a series of travel restrictions has been imposed by the Malaysian Government ⁽¹⁾. The Movement Control Order (MCO) was enforced on 16 March 2020 as the cases of COVID-19 mounted. Since then, Malaysia has gone through a few cycles of local tightening and relaxation in its combat against COVID-19 in line with local fluctuations of COVID-19 cases ⁽¹⁾. To date, the Conditional Movement Control Order (CMCO) as a less stringent variant of the MCO is still implemented in most regions of Malaysia with resurging cases of COVID-19 ⁽²⁾. The implementation of MCO has seen the compulsory adoption of precautionary measures such as self-isolation, physical distancing, mask-wearing, hand-cleaning and sanitization of premises in tandem with strict travel restrictions ⁽³⁾. Many of the measures particularly physical distancing, hand-cleaning and sanitization of premises are still practiced in the continual combat against COVID-19, while self-isolation and travel restrictions are situationally tightened and relaxed, often seen in the cycles of closure and reopening of certain non-essential services as well as adaptive rules on social gathering and cross-district or cross-state travels within the country ⁽²⁾⁽³⁾.

1 Coincidentally, COVID-19 and influenza are caused by contagious viruses which attack the respiratory
2 system, and they transmit in similar ways through droplets, body contacts and contacts with surfaces
3 containing the viruses ⁽⁴⁾⁽⁵⁾. However, COVID-19 has an overall higher transmission rate than influenza
4 and could result in more serious symptoms in certain individuals including chest pain, and breathing
5 difficulty which often prompts the use of ventilators ⁽⁴⁾. Influenza typically has an incubation period of
6 1 to 4 days whereas COVID-19 has a longer incubation period of 2 to 14 days ⁽⁴⁾. A median incubation
7 period of 5.1 days for COVID-19 has been reported ⁽⁶⁾. The precautionary measures to control the
8 transmission of COVID-19 and influenza are similar in certain aspects such as physical distancing,
9 mask-wearing and hand-sanitizing ⁽⁷⁾. Sanitization of premises is less prevalent in the control of
10 influenza probably because influenza is generally less severe than COVID-19 and the vaccines for
11 influenza are more established. Influenza is caused by the influenza virus and there are four types of
12 influenza virus, namely A, B, C and D ⁽⁸⁾. Influenza types A and B are responsible for seasonal epidemics
13 while influenza type C infects children mainly, causing mild symptoms. Influenza type D is not known
14 to infect or cause illness among humans ⁽⁸⁾. In Malaysia, influenza types A and B are common. The
15 A(H1N1) pdm09 virus emerged in mid-2009 is still prevalent in Malaysia and has been reported to co-
16 circulate with subtype A(H3) and type B viruses ⁽⁹⁾.

21 With similarities in the transmission means of COVID-19 and influenza, certain precautionary
22 measures against COVID-19 spread have also been employed for influenza even before the rise
23 of COVID-19. In view of their common clinical manifestations, similar means of transmission
24 prompting similar precautions, co-existence and the possibility of co-infection, it is therefore, of
25 interest to examine if the combat against COVID-19 in Malaysia has an effect on the national
26 influenza cases and whether it yields any co-benefits in controlling influenza. Influenza is known to
27 exist all year round in tropical countries like Malaysia where its seasonal variation is less obvious. It
28 burdens the healthcare system of the country to a certain extent ⁽¹⁰⁾. As such, changes in the
29 incidence of influenza due to the implementation of COVID-19 control measures would have
30 implications on the national healthcare.

34 Currently, there is no regional study conducted to examine the effects of COVID-19 control
35 on influenza cases. Such studies are few even on a global scale. Relevant studies mostly concern
36 influenza and COVID-19 coinfection as well as threats due to the co-existence of influenza and
37 COVID-19 ⁽⁴⁾⁽¹¹⁾⁽¹²⁾. There are also studies related to influenza vaccination to reduce influenza
38 infection and coinfection with COVID-19 ⁽¹³⁾⁽¹⁴⁾. There is one study examining the potential decrease
39 of influenza in the United States, Australia, Chile and South Africa as the COVID-19 pandemic
40 unfolded in 2020 but the study lacked statistical analyses ⁽⁷⁾. In view of the concern for COVID-19
41 and influenza coinfection, a study on the incidence of influenza during the COVID-19 era would help
42 to address the concern.

48 **Methods**

51 Data of daily new COVID-19 cases in Malaysia were sourced from the data platform maintained by
52 the Centre for System Science and Engineering (CSSE) of John Hopkins University ⁽¹⁵⁾. COVID-19
53 was detected in Malaysia on 25 January 2020 and Malaysia entered the first phase of movement
54 control on 18 March 2020 with steadily increasing cases of COVID-19 ⁽¹⁾. During movement control,
55 the public were ordered to self-isolate and practice physical distancing and there was strong
56 advice on the wearing of masks and hand-sanitizing. Sanitization of premises was also vigorously
57 conducted ⁽¹⁾. More than a year after the outbreak, while self-isolation has been lifted, physical
58 distancing, the wearing of masks, the upkeep of personal hygiene and disinfection are still
59 frequently implemented and are claimed to be the new norms ⁽³⁾. The data for COVID-19 cases
60 from Week 6 of 2020 starting on 3 February 2020 till Week 14 of 2021 ending on 4 April 2021
61 were extracted from the database to
62
63
64
65

demonstrate the progression of COVID-19 since the early stage of its onset in Malaysia (also see Figure 1 for the timeline).




Figure 1. Timeline of COVID-19 Pandemic in Malaysia and the Data Periods Analysed

Data for weekly number of influenza cases were obtained from the Flunet fed by the Global Influenza Surveillance and Response System (GISRS) maintained by the World Health Organization (WHO). Two sets of weekly influenza data were sourced ⁽¹⁶⁾. One set of the data comprised weekly number of influenza cases from Week 49 of 2018 commencing on 3 December 2018 to Week 5 of 2020 ending on 2 February 2020 while the other set of the data comprised the cases from Week 6 of 2020 to Week 14 of 2021 in parallel to the weekly COVID-19 cases (Figure 1). The former represented the number of influenza cases before the implementation of COVID-19 precautionary measures while the latter represented the number of influenza cases during the development, implementation and continual upkeep of COVID-19 precautionary practices in line with the unfolding of COVID-19 transmission in Malaysia and its progression to the current new norms in the COVID-19 era. The first COVID-19 cases were detected in Malaysia on Week 4 of 2020. With increase of cases in the subsequent week, flight suspension, travel restriction and precautionary measures were initiated. It is therefore deemed that Week 6 of 2020 is sufficiently representative of the commencement and development of COVID-19 precautions in Malaysia ⁽¹⁾. The influenza cases were classified based on the types and subtypes of influenza viruses detected. There were two types of influenza viruses detected, i.e. Type A and Type B. The subtypes of influenza A viruses reported in the period of interest were A(H1N1) pdm09, A(H3) and other non-subtyped A viruses whereas those of influenza B viruses were labelled as lineage not determined. This is because, unlike influenza A viruses, the lineages of influenza B viruses were not differentiated during diagnostic tests.

Statistical analyses were conducted to determine the correlations between the data and whether the datasets pre-COVID-19 and after the onset of COVID-19 in Malaysia were significantly different. Normality test was conducted on the datasets to determine if parametric or non-parametric tests would be adopted for the analyses. Shapiro-Wilk Test of Normality showed $p < 0.05$, indicating that a lack of normal distribution of the data ⁽¹⁷⁾. Therefore, non-parametric analyses were conducted where Spearman correlation was employed for correlational analysis between the COVID-19 cases and cases of influenza after the onset of COVID-19 (Week 6 2020 to Week 14 2020) to identify if patterns of variations in weekly COVID-19 and influenza incidences during this period were linked, without any

assumption made on the directions of variations. The correlational analysis did not aim to show the causal relations of any significant variation patterns of the data and it mainly served as a supplementary analysis in identification and explanation significant trends. More importantly, Mann-Whitney U Test was employed to determine if the datasets before and after the onset of COVID-19 were significantly different.

Results

Table 1 shows the descriptive statistics of the weekly COVID-19 cases from Week 6 of 2020 to the Week 14 of 2021 as well the cases of influenza by subtype within the same period and from Week 49 of 2018 to Week 5 of 2020. The weekly new cases of COVID-19 averaged at 821.9 over the 61 weeks tracked during which all influenza cases were consistently lower than 1 in comparison to the 61 weeks prior to the onset of COVID-19. Skewness and kurtosis provide an indication of normality, with a skewness of 0 representing a symmetrical dataset and a kurtosis of 3 representing sizes of two tails mirroring normal distribution ⁽¹⁷⁾. Extent of deviation from a skewness of 0 and a kurtosis of 3 implies the extent of deviation from normal distribution. Table 1 shows varying levels of deviation from normality which agree with the results Shapiro-Wilk Test of Normality revealing a lack of normal distribution, hence the appropriateness of non-parametric tests for this study.

Table 1: Descriptive Statistics of the Weekly New Cases of COVID-19 and Influenza

Weekly New Cases	n	Mean ± SD	Skewness	Kurtosis
From Week 6 2020 to Week 14 2021				
COVID-19	61	821.90 ± 1135.26	1.52	1.64
A(H1N1) pdm09	61	0.77 ± 2.21	2.95	7.74
A(H3)	61	0.64 ± 1.80	2.88	7.51
A(not subtyped)	61	0.16 ± 0.76	5.44	31.26
B(lineage not determined)	61	0.25 ± 1.03	4.82	23.60
From Week 49 2018 to Week 5 2020				
A(H1N1) pdm09	61	7.41 ± 6.59	1.26	1.06
A(H3)	61	3.16 ± 2.60	0.62	-0.48
A(not subtyped)	61	2.23 ± 3.48	2.12	4.42
B(lineage not determined)	61	9.08 ± 7.33	1.13	1.14

Figure 2 shows the weekly cases of COVID-19 from Week 6 2020 to Week 14 2021. The weekly new cases were consistently below 250 until Week 41 2020 (equivalent to Week 36 in Figure 2) in which the new cases shot to 691.57. The weekly new cases increased progressively after that, peaking at 4451.14 in Week 5 2021 (equivalent to Week 52 in Figure 2) and signalling a new wave of COVID-19 transmission. The spike of COVID-19 cases since Week 41 or October 2020 could be attributed to the relaxation of the CMCO which was set to replace the MCO implemented between 18 March 2020 to 3 May 2020 (Figure 1) ⁽¹⁸⁾. The CMCO came into force on 4 May 2020 and ended on 9 June, after which the RMCO took over. Implementation of the RMCO saw the lifting of a number of restrictions such as those on interstate travel, religious activities, meeting and events as well as non-essential services such as wellness and entertainment centres ⁽¹⁸⁾. The RMCO was extended until 31 March 2021 with re-implementation of MCO and CMCO in certain states where the COVID-19 cases were high (Figure 1) ⁽¹⁸⁾. In Week 5 2021 (Week 52 in Figure 2), there was a mix of RMCO, MCO and CMCO implemented throughout Malaysia with MCO effected in states with alarming new COVID-19 cases ⁽²⁾⁽¹⁸⁾. This

resembles the suppression and relaxation model of COVID-19 control practiced in many other regions of the world ⁽¹⁹⁾.




Figure 2: Weekly New COVID-19 Cases from Week 6 2020 to Week 14 2021 (Note: Week 6 2020 is denoted as Week 1 in the graph)

Figure 3a shows a sharp drop of influenza cases as early as Week 11 of 2020 (equivalent to Week 6 in Figure 3a) while cases of A(H1N1) pdm09 and A(H3) decreased to zero on Week 14 of 2020 (equivalent to Week 9 in Figure 3a), almost two weeks after the commencement of MCO. The influenza cases have remained at zero most of the time since then until Week 14, 2021. In contrast, the influenza cases during the pre-COVID-19 period from Week 49 2018 to Week 5 2020 fluctuated, with A(H1N1) pdm09 and B seemed to be more prevalent. Influenza B cases peaked in Week 25 of 2019 (equivalent to Week 29 in Figure 3b) and remained high in the next 10 weeks whereas A(H1N1) pdm09 cases rose in Week 51 of 2019 (equivalent to Week 55 in Figure 3b) and were high in the following 5 weeks.





Figure 3: Weekly New Influenza Cases from (a) Week 6 2020 – Week 14 2021, (b) Week 49 2018 – Week 5 2020 (Note: The beginning of the study period is denoted as Week 1)

Mann-Whitney U Test was performed on the influenza datasets before and after the onset of COVID-19 to determine if implementation of the COVID-19 control measures had any significant effects on the influenza cases. The Mann-Whitney U Test in Table 2 shows that the asymptotic significance, p , of all the paired datasets of influenza subtypes was <0.001 , demonstrating significant differences.

Table 2: Mann-Whitney U Test for Influenza Cases before and after the Onset of COVID-19

Pair of Influenza Cases*	Mann-Whitney U	Wilcoxon W	Z	Asymptotic Significance (2-tailed)
A1(H1N1) pdm09	371.50	2262.50	-8.05	<0.001
A(H3)	610.00	2501.00	-6.91	<0.001

A(not subtyped)	989.50	2880.50	-5.54	<0.001
B(lineage not determined)	146.50	2037.50	-9.32	<0.001

* Week 6 2020 – Week 14 2021 and Week 49 2019 – Week 5 2020

Spearman’s correlation shows that weekly new COVID-19 cases have significant negative correlations with weekly new cases of A(H1N1) pdm09 ($r_s = -0.44$), A(H3) ($r_s = -0.39$) and A(not subtyped) ($r_s = -0.40$) at significance, $p = 0.01$. At $p = 0.05$, the negative correlation between weekly new COVID-19 cases and weekly new influenza B cases is significant. The negative correlations, albeit slight to moderate, are statistically significant, indicating that the influenza cases decreased with advancing COVID-19 and its transmission control. Significant positive correlations are observed among the influenza cases possibly due to the concerted decline in the cases of all the influenza subtypes reported. The positive correlation was particularly strong between A(H1N1) pdm09 and A(H3) ($r_s = 0.94$).

Table 3: Spearman’s Correlation between COVID-19 and Influenza Cases from Week 6 2020 to Week 14 2021

	COVID-19	A(H1N1) pdm09	A(H3)	A(not subtyped)	B(lineage not determined)
COVID-19	1.00				
A(H1N1) pdm09	-0.44**	1.00			
A(H3)	-0.39**	0.94**	1.00		
A(not subtyped)	-0.40**	0.69**	0.67**	1.00	
B(lineage not determined)	-0.27*	0.59**	0.61**	0.42**	1.00

**Correlation is significant at the 0.01 level (2-tailed)

*Correlation is significant at the 0.05 level (2-tailed)

Discussion

Since the onset of COVID-19, wearing of masks and upkeeping of hand hygiene have been practiced and these practices culminated when the MCO came into force ⁽¹⁾. During MCO, there were strict regulations on physical distancing, self-isolation, travel and sanitization of premises ⁽¹⁾. Even after the transition to CMCO and RMCO, physical distancing, wearing of masks, upkeep of hand hygiene and sanitization have been widely practiced while self-isolation and travel restrictions are situational and have been imposed to different extents ⁽²⁰⁾. The practices of physical distancing, wearing of masks and upkeeping of hand hygiene are collectively known as the new norm in the COVID-19 era, and their wide adoption might have contributed to the reduction in weekly new cases of all influenza subtypes reported, which is shown in Figure 3a ⁽³⁾.

Fluctuation in the influenza cases before the onset of COVID-19 (Figure 3b) seemed to imply seasonal variations. Roussel et al. investigated the correlation between weekly variations of influenza cases and climatic factors in a temperate setting ⁽²¹⁾. They found average temperature, absolute humidity, sunshine duration and relative humidity to have significant impacts on the weekly variations but singling out the factors were difficult. In Malaysia located in the tropics, the seasonal variation of influenza cases is less obvious and influenza cases are reported throughout the year ⁽⁹⁾. This aligns with Figure 3b that there were influenza cases all year round. Despite the spikes in Figure 3b, seasonal

1 variation of the influenza cases cannot be concluded from this study and is outside the scope of this
2 study.

3 While measures to combat COVID-19 have been implemented since its onset, the measures seem to
4 be more effective in bringing down influenza cases compared to COVID-19, likely because of the higher
5 reproductive number of COVID-19 which have resulted in its quicker transmission ⁽²²⁾. In addition, the
6 longer incubation period of COVID-19 compared to influenza and the lack of symptoms in certain
7 individuals also contribute to the difficulty in its detection at the early stage of infection ⁽⁴⁾. Therefore,
8 unlike influenza, resurgence of COVID-19 cases has occurred throughout the movement control period
9 in Malaysia.
10
11

12 Malaysia was generally well-prepared for COVID-19 since the WHO declared it a global
13 pandemic during which the Ministry of Health implemented a robust preparedness plan which
14 involved ramping up screening and sampling at entry points as well as health clinics and
15 hospitals ⁽²³⁾. Besides, the Malaysian government also collaborated with private laboratories in
16 the performance of COVID-19 rRT-PCR tests, with the combined ability of all 43 public and private
17 laboratories hitting 16,635 rRT-PCR tests per day. The increasing demand for COVID-19 tests did
18 not seem to significantly affect influenza testing ⁽²³⁾. During the pandemic period from Week 6
19 2020 to Week 14 2021, the total number of samples for influenza test received was 3478 and the
20 total number of samples processed was 3447, which is equivalent to a process rate of 99.1% ⁽²⁴⁾.
21 From Week 49 2018 to Week 5 2020 prior to the pandemic and MCO in Malaysia, the total number
22 of samples for influenza test received was 7224 and the number of samples processed was 6999,
23 hence a process rate of 96.9%. While the number of samples for influenza test received during the
24 pandemic period was down by slightly more than half, the sample process rate was in fact
25 marginally higher than before the pandemic ⁽²⁴⁾. Converting to the number of positive influenza
26 cases per 100 of samples processed, there were 19.2 between Week 49 2018 and Week 5 2020,
27 whereas there were only 3.2 between Week 6 2020 and Week 14 2021. Being in the tropics, the
28 influenza cases in Malaysia are less susceptible to seasonal variation than in the temperate
29 regions and they have been detected all year round (Figure 3b). In addition, the influenza cases in
30 2020 was the lowest compared to the previous three years, especially after the initiation of MCO ⁽²⁴⁾.
31
32
33
34
35
36
37

38 The study faces certain limitations due to its correlational nature. Though it shows that the influenza
39 cases before and after the onset of COVID-19 are significantly different, it does not establish the causal
40 relations for the reduction in influenza cases after the onset of COVID-19. The plausible causes of
41 influenza decline were deduced from the obvious implementation of COVID-19 control measures and
42 behavioral changes ensuing COVID-19 which is still evident to date. Due to limited data on the
43 frequency of sanitation, the use of hand sanitizers or the practice of hand hygiene, the use of masks
44 as well as the practices of physical distancing, this study does not examine how each of the factors
45 contribute to the reduction of influenza cases. Similarly, the lack of data for these variables does not
46 permit the comparison of whether the adoption or intensified adoption of these precautions would
47 significantly reduce influenza cases. Besides, this study may not have accounted for all confounding
48 factors such as changes in health-seeking behaviors during the COVID-19 era.
49
50
51

52 As such, future studies can aim to address the limitations by examining the causal relations between
53 the COVID-19 control measures and the decline in influenza cases. These control measures or factors
54 could be examined separately under controlled settings to investigate their effectiveness in controlling
55 influenza and the underlying mechanisms. The correlational studies could be extended to different
56 countries adopting similar COVID-19 control measures to confirm if the control measures were the
57 contributing factors of diminishing influenza cases.
58
59
60
61
62
63
64
65

Conclusion

This study reveals that there are significant differences in the weekly cases of influenza before and after the onset of COVID-19 in Malaysia attributed probably to the COVID-19 control measures consisting mainly of physical distancing, mask-wearing, keeping of hand hygiene and sanitization of premises which is still widely practiced currently. The implementation of self-isolation is situational depending on the local cases of COVID-19 and therefore, is constrained in accounting for the consistently low weekly new influenza cases throughout the COVID-19 era. This study highlights the co-benefit of implementing COVID-19 control measures, particularly in the significant reduction of influenza. The lower respiratory tract infection associated with influenza was reported to cause 2.6 deaths per 100,000 population in Malaysia in 2017 ⁽²⁵⁾. This study suggests that situational practices of physical distancing, mask-wearing and keeping of hand hygiene during influenza seasons post-COVID-19 could be beneficial and hand hygiene is a practice which could be continued in the control of influenza post-COVID-19.

Acknowledgments

The authors wish to thank BNU-HKBU United International College and Curtin University Malaysia for providing the necessary resources to complete this study.

Funding: None

Completing interest: None declared

Ethical approval: Not required as the study does not involve any human subjects.

References

1. Tang KHD. Movement control as an effective measure against Covid-19 spread in Malaysia: an overview. *J Public Health (Bangkok)* [Internet]. 2020; Available from: <https://doi.org/10.1007/s10389-020-01316-w>
2. Borneo Post Online. Specific movement control enforced in various zones - SDMC. 2021 Apr 21; Available from: <https://www.theborneopost.com/2021/04/21/specific-movement-control-enforced-in-various-zones-sdmc/>
3. Tang KHD. Controversies of The Post-Lockdown New Normal - It May Not be Entirely Normal [Internet]. Vol. 11, *Current Research Journal of Social Sciences and Humanities*. 2021. Available from: <http://journalofsocialsciences.org/article/60/>
4. Solomon DA, Sherman AC, Kanjilal S. Influenza in the COVID-19 Era. *JAMA* [Internet]. 2020 Oct 6;324(13):1342–3. Available from: <https://doi.org/10.1001/jama.2020.14661>
5. Tang KHD. A scoping review of studies on COVID-19. *Int J Sci Healthc Res*. 2020;5(2):205–14.
6. McAloon C, Collins Á, Hunt K, Barber A, Byrne AW, Butler F, et al. Incubation period of COVID-19: a rapid systematic review and meta-analysis of observational research. *BMJ Open* [Internet]. 2020 Aug 1;10(8):e039652. Available from: <http://bmjopen.bmj.com/content/10/8/e039652.abstract>
7. Olsen SJ, Azziz-Baumgartner E, Budd AP, Brammer L, Sullivan S, Pineda RF, et al. Decreased influenza activity during the COVID-19 pandemic—United States, Australia, Chile, and South

1 Africa, 2020. *Am J Transplant* [Internet]. 2020 Dec 1;20(12):3681–5. Available from:
2 <https://doi.org/10.1111/ajt.16381>

- 3 8. Suess T, Remschmidt C, Schink SB, Schweiger B, Heider A, Milde J, et al. Comparison of
4 Shedding Characteristics of Seasonal Influenza Virus (Sub)Types and Influenza
5 A(H1N1)pdm09; Germany, 2007–2011. *PLoS One* [Internet]. 2012 Dec 11;7(12):e51653.
6 Available from: <https://doi.org/10.1371/journal.pone.0051653>
7
- 8 9. Saha S, Chadha M, Al Mamun A, Rahman M, Sturm-Ramirez K, Chittaganpitch M, et al.
9 Influenza seasonality and vaccination timing in tropical and subtropical areas of southern and
10 south-eastern Asia. *Bull World Health Organ*. 2014;92:318–30.
11
- 12 10. Jamal I-CS. The burden of human influenza in Malaysia. *Med J Malaysia*. 2015;70(3):127–30.
13
- 14 11. Konala VM, Adapa S, Naramala S, Chenna A, Lamichhane S, Garlapati PR, et al. A Case Series
15 of Patients Coinfected With Influenza and COVID-19. *J Investig Med High Impact Case Reports*
16 [Internet]. 2020 Jan 1;8:2324709620934674. Available from:
17 <https://doi.org/10.1177/2324709620934674>
18
- 19 12. Ozaras R, Cirpin R, Duran A, Duman H, Arslan O, Bakcan Y, et al. Influenza and COVID-19
20 coinfection: Report of six cases and review of the literature. *J Med Virol* [Internet]. 2020 Nov
21 1;92(11):2657–65. Available from: <https://doi.org/10.1002/jmv.26125>
22
- 23 13. Grech V, Borg M. Influenza vaccination in the COVID-19 era. *Early Hum Dev* [Internet].
24 2020;148:105116. Available from:
25 <https://www.sciencedirect.com/science/article/pii/S0378378220303455>
26
- 27 14. Maltezou HC, Theodoridou K, Poland G. Influenza immunization and COVID-19. *Vaccine*
28 [Internet]. 2020/07/29. 2020 Sep 3;38(39):6078–9. Available from:
29 <https://pubmed.ncbi.nlm.nih.gov/32773245>
30
- 31 15. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time.
32 *Lancet Infect Dis*. 2020;20(5):533–4.
33
- 34 16. Hay AJ, McCauley JW. The WHO global influenza surveillance and response system (GISRS)—A
35 future perspective. *Influenza Other Respi Viruses* [Internet]. 2018 Sep 1;12(5):551–7.
36 Available from: <https://doi.org/10.1111/irv.12565>
37
- 38 17. Ghasemi A, Zahediasl S. Normality tests for statistical analysis: a guide for non-statisticians.
39 *Int J Endocrinol Metab* [Internet]. 2012/04/20. 2012;10(2):486–9. Available from:
40 <https://pubmed.ncbi.nlm.nih.gov/23843808>
41
- 42 18. FMT. Timeline of a year in lockdown. 2021 Mar 18; Available from:
43 <https://www.freemalaysiatoday.com/category/nation/2021/03/18/timeline-of-a-year-in-lockdown/>
44
- 45 19. Chowdhury R, Heng K, Shawon MSR, Goh G, Okonofua D, Ochoa-Rosales C, et al. Dynamic
46 interventions to control COVID-19 pandemic: a multivariate prediction modelling study
47 comparing 16 worldwide countries. *Eur J Epidemiol* [Internet]. 2020;35(5):389–99. Available
48 from: <https://doi.org/10.1007/s10654-020-00649-w>
49
- 50 20. The Straits Times. Malaysia extends recovery movement control order to March 31. 2021 Jan
51 1; Available from: <https://www.straitstimes.com/asia/se-asia/malaysia-extends-recovery-movement-control-order-to-mar-31>
52
- 53 21. Roussel M, Pontier D, Cohen J-M, Lina B, Fouchet D. Quantifying the role of weather on
54 seasonal influenza. *BMC Public Health* [Internet]. 2016;16(1):441. Available from:
55
56
57
58
59
60
61
62
63
64
65

<https://doi.org/10.1186/s12889-016-3114-x>

22. Billah MA, Miah MM, Khan MN. Reproductive number of coronavirus: A systematic review and meta-analysis based on global level evidence. *PLoS One* [Internet]. 2020 Nov 11;15(11):e0242128. Available from: <https://doi.org/10.1371/journal.pone.0242128>
23. Rahman F. The Malaysian response to COVID-19: building preparedness for 'surge capacity', testing efficiency and containment. *Eur Pharm Rev* [Internet]. 2020; Available from: <https://www.europeanpharmaceuticalreview.com/article/125084/the-malaysian-response-to-covid-19-building-preparedness-for-surge-capacity-testing-efficiency-and-containment/>
24. World Health Organization. Flunet [Internet]. 2021 [cited 2021 Jun 22]. Available from: <https://apps.who.int/flumart/Default?ReportNo=12>
25. Troeger CE, Blacker BF, Khalil IA, Zimsen SRM, Albertson SB, Abate D, et al. Mortality, morbidity, and hospitalisations due to influenza lower respiratory tract infections, 2017: an analysis for the Global Burden of Disease Study 2017. *Lancet Respir Med* [Internet]. 2019;7(1):69–89. Available from: <https://www.sciencedirect.com/science/article/pii/S221326001830496X>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65