

Candidate Tidal Disruption Event AT2019fdr Coincident with a High-Energy Neutrino

Simeon Reusch^{1,2,*} Robert Stein^{1,2,3} Marek Kowalski^{1,2,†} Sjoert van Velzen⁴ Anna Franckowiak^{1,5}
 Cecilia Lunardini⁶ Kohta Murase^{7,8} Walter Winter¹ James C. A. Miller-Jones⁹ Mansi M. Kasliwal³
 Marat Gilfanov^{10,11} Simone Garrappa^{1,2} Vaidehi S. Paliya¹² Tomás Ahumada¹³ Shreya Anand³
 Cristina Barbarino¹⁴ Eric C. Bellm¹⁵ Valéry Brinnel² Sara Buson¹⁶ S. Bradley Cenko^{17,18} Michael W. Coughlin¹⁹
 Kishalay De³ Richard Dekany²⁰ Sara Frederick¹³ Avishay Gal-Yam²¹ Suvi Gezari^{22,13} Marcello Giroletti²³
 Matthew J. Graham³ Viraj Karambelkar³ Shigeo S. Kimura²⁴ Albert K. H. Kong²⁵ Erik C. Kool¹⁴
 Russ R. Laher²⁶ Pavel Medvedev¹⁰ Jannis Necker^{1,2} Jakob Nordin² Daniel A. Perley²⁷ Mickael Rigault²⁸
 Ben Rusholme²⁶ Steve Schulze²⁹ Tassilo Schweyer¹⁴ Leo P. Singer³⁰ Jesper Sollerman¹⁴ Nora Linn Strotzjohann²¹
 Rashid Sunyaev^{10,11} Jakob van Santen¹ Richard Walters²⁰ B. Theodore Zhang⁸ and Erez Zimmerman²¹

¹Deutsches Elektronen Synchrotron DESY, Platanenallee 6, D-15738 Zeuthen, Germany

²Institut für Physik, Humboldt-Universität zu Berlin, D-12489 Berlin, Germany

³Division of Physics, Mathematics, and Astronomy, California Institute of Technology, 1200 East California Boulevard, MC 249-17, Pasadena, California 91125, USA

⁴Leiden Observatory, Leiden University, Postbus 9513, 2300 RA, Leiden, Netherlands

⁵Fakultät für Physik und Astronomie, Ruhr-Universität Bochum, D-44780 Bochum, Germany

⁶Department of Physics, Arizona State University, Tempe, Arizona 85287-1504, USA

⁷Department of Physics; Department of Astronomy and Astrophysics; Center for Multimessenger Astrophysics, Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, Pennsylvania 16802, USA

⁸Center for Gravitational Physics, Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto, Kyoto 606-8502, Japan

⁹International Centre for Radio Astronomy Research - Curtin University, GPO Box U1987, Perth, Washington 6845, Australia

¹⁰Space Research Institute (IKI), Russian Academy of Sciences, Profsoyuznaya ul. 84/32, Moscow 117997, Russia

¹¹Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, D-85741 Garching, Germany

¹²Aryabhata Research Institute of Observational Sciences (ARIES), Manora Peak, Nainital-263001 Uttarakhand, India

¹³Department of Astronomy, University of Maryland, College Park, Maryland 20742, USA

¹⁴Department of Astronomy, The Oskar Klein Centre, Stockholm University, SE-106 91 Stockholm, Sweden

¹⁵DIRAC Institute, Department of Astronomy, University of Washington, 3910 15th Avenue NE, Seattle, Washington 98195, USA

¹⁶Institut für Theoretische Physik and Astrophysik, Universität Würzburg, D-97074 Würzburg, Germany

¹⁷Astrophysics Science Division, NASA Goddard Space Flight Center, 8800 Greenbelt Road, Greenbelt, Maryland 20771, USA

¹⁸Joint Space-Science Institute, University of Maryland, College Park, Maryland 20742, USA

¹⁹School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA

²⁰Caltech Optical Observatories, California Institute of Technology,

1200 East California Boulevard, MC 249-17, Pasadena, California 91125, USA

²¹Department of Particle Physics and Astrophysics, Weizmann Institute of Science, 234 Herzl Street 76100 Rehovot, Israel

²²Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, Maryland 21218, USA

²³INAF - Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, via Gobetti 93/3, I-40129 Bologna, Italy

²⁴Frontier Research Institute for Interdisciplinary Sciences; Astronomical Institute, Tohoku University, Sendai 980-8574, Japan

²⁵Institute of Astronomy, National Tsing Hua University, No. 101 Section 2 Kuang-Fu Road, Hsinchu 30013, Taiwan

²⁶IPAC, California Institute of Technology, 1200 East California Boulevard, MC 249-17, Pasadena, California 91125, USA

²⁷Astrophysics Research Institute, Liverpool John Moores University, 146 Brownlow Hill, Liverpool L3 5RF, United Kingdom

²⁸Univ Lyon, Univ Claude Bernard Lyon 1, CNRS/IN2P3, UMR 5822, F-69622 Villeurbanne, France

²⁹The Oskar Klein Centre, Physics Department of Physics, Stockholm University, Albanova University Center, SE-106 91 Stockholm, Sweden

³⁰NASA Goddard Space Flight Center, University of Maryland, Baltimore County, Greenbelt, Maryland 20771, USA



(Received 16 December 2021; accepted 9 March 2022; published 3 June 2022)

The origins of the high-energy cosmic neutrino flux remain largely unknown. Recently, one high-energy neutrino was associated with a tidal disruption event (TDE). Here we present AT2019fdr, an exceptionally luminous TDE candidate, coincident with another high-energy neutrino. Our observations, including a bright dust echo and soft late-time x-ray emission, further support a TDE origin of this flare. The probability of finding two such bright events by chance is just 0.034%. We evaluate several models for neutrino production and show that AT2019fdr is capable of producing the observed high-energy neutrino, reinforcing the case for TDEs as neutrino sources.

DOI: 10.1103/PhysRevLett.128.221101

Neutrino astronomy is at a crossroads: While a flux of high-energy cosmic neutrinos has been firmly established through observations with the IceCube Neutrino Observatory [1–4], identifying their sources has been a challenge. The emission of cosmic neutrinos is a smoking-gun signature for hadronic acceleration (see Ref. [5] for a recent review), and discovering their sources will allow us to resolve long-standing questions about the production sites of high-energy cosmic rays.

Three sources have thus far been associated with neutrinos at post-trial significance of $\approx 3\sigma$, which can be considered evidence for a true association [6]. In 2017, the flaring blazar TXS 0506 + 056 was identified as the likely source of neutrino alert IC170922A [7]. This same source was also associated with a neutrino flare in 2014–2015 [8], occurring during a period without significant electromagnetic flaring activity [9]. In 2019, the tidal disruption event (TDE) AT2019dsg was identified as the likely source of IC191001A [10]. More recently, the IceCube Collaboration reported a clustering of neutrinos from the direction of the active galactic nucleus (AGN) in the nearby galaxy NGC 1068 [11]. AGN are galaxies with high levels of supermassive black hole (SMBH) accretion, and have been long proposed as high-energy neutrino sources [12–17]. These associations and other conceptual arguments suggest that the neutrino flux may arise from a mixture of different astrophysical populations [18–20], although AGN or another source class can still be dominant [21].

TDEs are rare transients that occur when stars pass close enough to SMBHs and get destroyed by tidal forces. The result of this destruction is a luminous electromagnetic flare with a timescale of \sim months. Theoretical studies have suggested that TDEs might be sources of high-energy neutrinos and ultrahigh-energy cosmic rays [22–39]. Some models consider emission from a relativistic jet, while others propose additional neutrino production scenarios, e.g., in a disk, disk corona, or wind (see Refs. [36,40]). In the case of AT2019dsg, radio observations confirmed long-lived nonthermal emission from the source [10,41–44], but generally disfavor those models relying on the presence of an on-axis relativistic jet [35] in the standard leptonic radio emission scenario.

TDEs and AGN flares are ultimately both modes of SMBH accretion. Some models highlight this potential similarity, and have developed common frameworks for neutrino emission from both cases (see, e.g., Ref. [36]). However, AGN flares are vastly more numerous than TDEs, injecting significantly more energy into the Universe. If TDEs nonetheless contribute significantly to the neutrino flux, they must be very efficient neutrino emitters. Whether there are particular characteristics of TDEs that enable efficient neutrino production, and whether these conditions are also present in particular classes of AGN accretion flares, remain open questions for neutrino astronomy.

Bridging these two astrophysical populations, we here report new observations of AT2019fdr, a candidate TDE in a Narrow-Line Seyfert 1 (NLSy1) active galaxy [45]. Similar to AT2019dsg, AT2019fdr was identified as a likely neutrino source by the neutrino follow-up program of the Zwicky Transient Facility (ZTF) [46–48]. AT2019fdr lies within the reported 90% localization region of the IceCube high-energy neutrino IC200530A [49]. The observations were processed by `nuztf`, our multimessenger analysis pipeline [50,51], which searches for extragalactic transients in spatial and temporal coincidence with high-energy neutrinos [10,52], and AT2019fdr was reported as a candidate [53].

AT2019fdr, a long-duration flare (see Fig. 1) of apparent nuclear origin, was first discovered by ZTF one year prior to the neutrino detection [45,54]. AT2019fdr reached a peak flux of $1.3 \times 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$ in the optical ZTF g band on August 10, 2019, before slowly fading. With a peak g -band luminosity of $L_{\text{peak}} = 2.9 \times 10^{44} \text{ erg s}^{-1}$, AT2019fdr was an extraordinarily luminous event. At the time of neutrino detection, it had decayed to $\sim 30\%$ of its peak flux, and was still detected by ZTF as of August 2021. Forced photometry using data from ZTF (up to 400 days prior to the flare) as well as from the Palomar Transient Factory (2010–2016) [55] shows no historical variability.

AT2019fdr was classified as a probable TDE, though an extreme AGN flare origin could not be ruled out [45]. High-resolution spectra yielded a redshift of $z = 0.267$. Using a spectrum from the Alhambra Faint Object Spectrograph and Camera (ALFOSC), on the Nordic Optical Telescope (NOT; PI: Sollerman), a virial black hole mass estimate of $M_{\text{BH}} = 10^{7.55 \pm 0.13} M_{\odot}$ was inferred; for further details refer to the Supplemental Material [57].

Though the classification of AT2019fdr based on early observations included the possibility of it being a Type II superluminous supernova (SLSN-II, see Ref. [176]) [177], leading to further studies [178], its subsequent spectroscopic and photometric evolution was not consistent with expectations for SLSNe. Frederick *et al.* [45] already disfavored the SLSN hypothesis based on the long-lived U-band and the UV emission, the flare’s longevity, emission at the blue end of the Balmer line profiles as well as its proximity to the nucleus of the galaxy. Here we add a late-time x-ray detection and the detection of a strong infrared echo, rendering a SLSN interpretation less likely (see below).

After discovery, AT2019fdr was also observed by the Ultraviolet and Optical Telescope (UVOT) [179] aboard the *Neil Gehrels Swift Observatory (Swift)* [45,180]. Additional observations continued up to June 7, 2020, including one epoch shortly after the neutrino detection. By that point, the transient had faded by 84% in the UVW1-band from its peak luminosity of $2.1 \times 10^{44} \text{ erg s}^{-1}$. AT2019fdr was not detected in any of the simultaneous x-ray observations by the




FIG. 1. The bottom plot shows the light curve in the optical ZTF g band, the infrared P200 Ks and WISE W1 band as well as the modeled dust echo (black line, dashdot), with the neutrino arrival time marked with a red dotted vertical line. The SRG/eROSITA x-ray measurements are also included. The shaded gray areas are averaged and their respective SEDs are shown in the top panels, including a fitted blue and a red blackbody (blue dashed and red dotted curve; lab frame), as well as the combined spectrum (black solid curve). The left axes all show νF_ν , where F_ν is the spectral flux density at frequency ν , while the right axes show νL_ν , where L_ν is the luminosity at frequency ν . Note: SRG/eROSITA data are given in units of integrated flux. The second epoch (middle plot on top) encompasses several months to include both WISE and P200 infrared data points. The global values for line-of-sight dust extinction are $A_V = 0.45^{+0.14}_{-0.14}$ mag, assuming $R_V = 3.1$ and the Calzetti attenuation law [56]. Note that the x-ray measurements were not included in the blackbody fits. The luminosities are given in the source rest frame.

Swift X-ray Telescope (XRT) [181], yielding a combined 3σ flux upper limit of $1.4 \times 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$ for all observations before neutrino arrival (corrected for absorption).

The position of AT2019fdr was also visited by the eROSITA telescope [182] aboard the *Spectrum-Roentgen-Gamma* (SRG) mission [183] four times. The first two visits did not detect an excess, with a mean flux upper limit of $2.7 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$ at the 95% confidence level. However, at the third visit on March 10–11, 2021, it detected late time x-ray emission from the transient with an energy flux of $6.2^{+2.7}_{-2.1} \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$ in the 0.3–2.0 keV band, thus showing temporal evolution in the x-ray flux (see Fig. 1). The detection displayed a very soft thermal spectrum with a best fit blackbody temperature of 56^{+32}_{-26} eV.

The softness of the spectrum provides further evidence for AT2019fdr being a TDE rather than regular AGN variability, where soft spectra are rare [184]. Though NLSy1 galaxies generally exhibit softer x-ray spectra, the temperature of AT2019fdr is atypically low even in this

context (lower than all NLSy1s in Refs. [185] and [186]). Furthermore, x-ray emission is rarely seen for SLSNe [187], with only the first SLSN ever observed, SCP 06F6 [188], possibly showing an x-ray flux exceeding the luminosity of AT2019fdr [189]. This provides more evidence against the SLSN classification.

AT2019fdr was further detected at midinfrared (MIR) wavelengths as part of routine NEOWISE survey observations [190] by the Wide-field Infrared Survey Explorer (WISE) [191]. Using pre-flare archival NEOWISE data as baseline, a substantial flux increase was detected in both W1 and W2 band. MIR emission reached a peak luminosity of $1.9 \times 10^{44} \text{ erg s}^{-1}$ on August 13, 2020, over one year after the optical and UV peak. Complementary near-infrared (NIR) measurements were taken with the P200 wide field infrared camera (WIRC) [192] in the J, H, and Ks band. After subtracting a synthetic host model (see Supplemental Material [57]), a fading transient infrared signal was detected in all three bands; see Fig. 1.

We modeled this light curve as a composite of two unmodified blackbodies (a “blue” and a “red” blackbody). We interpret the time-delayed infrared emission as a dust echo: The blue blackbody heats surrounding dust, which then starts to glow. The light curve of this dust echo was inferred using the method described in Ref. [193] and the corresponding fit is shown in Fig. 1. An optical or UV bolometric luminosity of $L = 1.4^{+0.1}_{-0.1} \times 10^{45}$ erg s⁻¹ at peak was derived. By integrating this component over time, we derived a total bolometric energy of $E_{\text{bol}} = 3.4 \times 10^{52}$ erg (the red blackbody was not added, as dust absorption is already accounted for through the extinction correction). This is almost twice the inferred bolometric energy of ASASSN-15lh, which was one of the brightest transients ever reported [194] and was suggested to be a TDE [195]. Furthermore, the energy budget, bolometric evolution, and luminous dust echo suggest that AT2019fdr belongs to a class of TDE candidates observed in AGN (similar to PS1-10adi [196], AT2017gbl [197] or Arp 299-B AT1 [198]). For details on the modeling methods, see Supplemental Material [57].

Following the neutrino detection, we performed radio observations of AT2019fdr with a dedicated Very Large Array (VLA) [199] director’s discretionary time (DDT) program three times over a period of four months, and obtained multifrequency detections. AT2019fdr shows a featureless power law spectrum consistent with optically thin synchrotron emission above ~ 1 GHz with no significant intrinsic evolution between the epochs (see Supplemental Material [57]). The peak flux density was 0.39 ± 0.03 mJy in the 1–2 GHz band. The lack of apparent evolution suggests that the radio emission is not related to the transient, but rather originated from the AGN host. An additional subdominant transient component could be present.

No γ rays were detected by the *Fermi* large area telescope (*Fermi*-LAT) [200] between the first detection of AT2019fdr and one year after neutrino detection, yielding an upper limit of 1.3×10^{-12} erg s⁻¹ cm⁻² (see Ref. [201]).

AT2019fdr is the second probable neutrino-TDE association found by ZTF. To calculate the probability of finding two such coincident events by chance, while accounting for the fact that some TDEs will not be spectroscopically classified, we developed a broader sample of photometrically selected “candidate TDEs.” We selected “nuclear” transients that are at least as bright as AT2019fdr from the sample of ZTF transients, and applied cuts to identify TDE-like rise times and decay times (see Supplemental Material [57] and Ref. [201] for details). Our sample begins in 2018 (the ZTF survey start), and we further required a flare peak date before July 2020. We excluded only transients for which a TDE origin was ruled out through spectroscopic classification (i.e., our sample contains all unclassified candidates and all classified TDEs). To compute the sky source density at any given time, we conservatively estimated their average lifetime at

1 yr after discovery, yielding an effective source density of 1.7×10^{-4} per deg² of sky in the ZTF footprint (most TDEs evolve on shorter timescales, which—if accounted for—would reduce the effective source density). When including all 24 neutrinos followed up by our program by September 2021 (covering a combined area of 154.33 deg², see Supplemental Material [57]), the probability of finding any photometrically selected TDE candidate by chance is 2.6×10^{-2} , while the probability of finding two by chance is 3.4×10^{-4} (3.4σ). We emphasize that these estimates rely solely on the optical flux and a nuclear location in the host galaxy, and thus do not account for the additional luminous dust echoes or postflare x-ray detections observed for AT2019dsg and AT2019fdr.

Neutrino emission from AT2019fdr.—With a single neutrino observed in association with AT2019fdr, the inference of the neutrino flux will be subject to a large Eddington bias [202] and hence very uncertain. However, we can make a more robust statement on the neutrino flux by considering the underlying population (see, e.g., Ref. [10]). The detection of two high-energy neutrinos implies a mean expectation for the full TDE catalog in the range $0.36 < N_{\nu, \text{tot}} < 6.30$ at 90% confidence, where $N_{\nu, \text{tot}}$ is the cumulative neutrino expectation for the nuclear transients that ZTF has observed. AT2019fdr emits $\sim 2\%$ of the g -band peak energy flux for the population of nuclear transients, consisting of the 17 published ZTF TDEs (see Ref. [203]) and all TDE candidates as bright as AT2019fdr (see Supplemental Material [57] for the latter). If we take this as a proxy for the contribution of AT2019fdr to the neutrino emission, we would expect a total number of neutrinos $0.007 \lesssim N_{\nu} \lesssim 0.13$ for this source.

This estimate can be compared to model expectations. We present three different models invoking $p\gamma$ and/or pp interactions, where protons are efficiently accelerated in a disk corona, a subrelativistic wind or a relativistic jet (see Supplemental Material [57]). The resulting spectra are shown in Fig. 2. All models can explain the observed




FIG. 2. Neutrino fluence for the three models described here. The reported energy of the neutrino event [49], represented by the dotted vertical line, should be viewed as a lower limit to the neutrino energy.

energy of the IC200530A neutrino event; they also make predictions for the underlying “neutrino light curve,” though this can only be resolved once many neutrinos from TDEs have been detected. The obtained neutrino luminosities $L_\nu \lesssim 0.1L_{\text{Edd}} \simeq 5 \times 10^{44} \text{ erg s}^{-1}$ are consistent with theoretical expectations for most models [39].

In accretion flow models [34,36], the virial theorem implies a cosmic ray acceleration efficiency $\eta_{\text{CR}} < (1/40)(R/10R_s)^{-1}$ [36] for a cosmic-ray luminosity $L_{\text{CR}} = \eta_{\text{CR}}\dot{M}c^2$, where R is the emission radius and $R_s = 2GM/c^2$ is the Schwarzschild radius. Even for a mass accretion rate of $\dot{M} \sim 10L_{\text{Edd}}/c^2$, the neutrino luminosity would not exceed $\sim 10^{44} \text{ erg s}^{-1}$. In the case of AT2019fdr, the Eddington ratio $\lambda_{\text{Edd}} \equiv L/L_{\text{Edd}} \lesssim 0.07\text{--}0.3$ in the first 2 epochs, implying accretion near the Eddington limit around the peak and sub-Eddington accretion around the time of the neutrino detection. For such high accretion the disk plasma is collisional, while the coronal region may allow particle acceleration and nonthermal neutrino production [36]. This model yields $N_\nu \sim 0.007$ when evaluating its spectrum under the effective area of the neutrino alert channel [204]. This is within the expected range, albeit at its lower end. The time delay is consistent with quasisteady coronal emission. Alternatively, because the accretion rate gradually decreases, the neutrino time delay can be attributed to the formation of a collisionless corona that allows ion acceleration [36].

We also considered a subrelativistic wind with a velocity of $\sim 0.1c$, consistent with what was observed for AT2019dsg. Such a wind is naturally launched from the TDE disk (e.g., Ref. [205]), and may interact with tidal disruption debris. A strong shock is also expected from interactions between tidal streams. Ions can be accelerated at the shock via diffusive shock acceleration and produce neutrinos through inelastic pp and $p\gamma$ collisions [36]. In this subrelativistic wind model, the maximum proton energy can be as high as $\sim 10\text{--}100 \text{ PeV}$. If the cosmic ray luminosity is three times the optical luminosity, the expected number of muon neutrinos is $N_\nu \sim 0.002$, which falls outside the empirical range for this baryon loading factor. The neutrino light curve would trace the wind luminosity in the calorimetric limit, and the time delay is consistent with quasisteady radio emission.

In the relativistic jet model, external target photons from the disk are backscattered into the jet frame. Here we followed Ref. [35] for AT2019dsg, but adopted a unified model [206] to extrapolate to higher SMBH masses as given for AT2019fdr. We estimated a thermal far UV to x-ray spectrum with $T \simeq 34 \text{ eV}$. This turned out to be consistent with the late-time x-ray detection within the uncertainties. The isotropization timescale of the photons is expected to be given by the system size, suggesting a possible correlation with the dust echo; as a consequence the isotropized x-ray and dust echo light curves look very similar. The jet model allows for efficient particle acceleration and results in a

relatively large number of 0.027 neutrino events with a maximum $L_\nu \simeq 0.05L_{\text{Edd}}$ thanks to the beaming effect; however, direct signatures of the jet have not been observed.

Conclusions.—AT2019fdr was an exceptionally bright nuclear transient that was already identified in the literature as a probable TDE in an active galaxy [45]. In this Letter, we have presented new observational data, including the identification of a strong dust echo and soft late-time x-ray emission, which further support a TDE origin for this flare.

AT2019fdr was a very long-lived transient, one of the most luminous ever detected. For a TDE, the energy release would require a very massive star [207]. However, unlike for TDEs in quiescent galaxies, the AGN disk in AT2019fdr might provide the system with additional energy [208]. Furthermore, the post star-burst nature of the host increases the expected rate for TDEs [209–211].

AT2019fdr was the second candidate neutrino-TDE identified by our ZTF follow-up program. While AT2019fdr was far more luminous than AT2019dsg, the first TDE associated with a high-energy neutrino, it was also more distant. As a result, the two objects have comparable bolometric energy fluxes. The probability for finding two such bright neutrino-coincident TDEs by chance is just 3.4×10^{-4} , a sevenfold decrease relative to the previously reported single association [10]. The gain due to the second association is somewhat offset by the larger neutrino sample and the more inclusive candidate TDE selection. Within the framework of this Letter, the association of a second object results in a reduction of the chance probability by a factor of 75 versus a single association.

AT2019fdr and AT2019dsg share other similarities beyond their potential association with a high-energy neutrino. Intriguingly, AT2019dsg also displayed an unusually strong dust echo signal [201], indicating that the presence of large amounts of matter and an associated high star formation rate in the environment could be a common signature for high-energy neutrino production in such systems. A dedicated search for further associations based on this signature is presented in Ref. [201] and provides more supporting evidence for neutrino production in TDEs.

We studied neutrino emission from AT2019fdr using models previously applied to explain the observations of AT2019dsg. Similar to AT2019dsg, various plausible cosmic ray acceleration sites have been identified, such as the corona, a subrelativistic wind, or a relativistic jet. The number of expected muon neutrinos predicted by the corona and jet models is consistent with empirical constraints derived from the two TDE-neutrino associations. All models require efficient neutrino production at a neutrino luminosity comparable to a sizable fraction of the Eddington luminosity. The neutrino delay may be related to the size of the newly formed system (jet model) or the formation of a collisionless corona (corona model).

With two objects being associated with IceCube neutrino alerts, out of a number of 11.5 expected astrophysical

neutrinos (summed alert signalness, see Supplemental Material [57]), we obtain a fraction of $18_{-15}^{+38}\%$ (90% confidence level) of astrophysical neutrinos that could be explained due to ZTF-detected TDE candidates. Accounting for the incompleteness of our sample with the procedure in Ref. [10], our results imply that at least 7.8% of astrophysical neutrinos would come from the broader TDE population.

The search for neutrinos resulting in public alerts has a high energy threshold to reduce the background. Even when considering the full energy range of IceCube [212], the expected number of neutrino events from AT2019fdr remains below one. Therefore, the detection of additional lower-energy neutrinos from AT2019fdr is not expected (see also the search by the ANTARES neutrino observatory [213]).

Fully understanding the role of TDEs as particle accelerators will only be possible with comprehensive multi-wavelength and -messenger data. While the detailed production processes remain uncertain, the observations presented here provide further evidence that TDEs are highly efficient sources of high-energy neutrinos.

S. R. acknowledges support by the Helmholtz Weizmann Research School on Multimessenger Astronomy, funded through the Initiative and Networking Fund of the Helmholtz Association, DESY, the Weizmann Institute, the Humboldt University of Berlin, and the University of Potsdam. A. F. acknowledges support by the Initiative and Networking Fund of the Helmholtz Association through the Young Investigator Group program (A. F.). C. L. acknowledges support from the National Science Foundation (NSF) with Grant No. PHY-2012195. The work of K. M. is supported by the NSF Grants No. AST-1908689, No. AST-2108466, and No. AST-2108467, and KAKENHI No. 20H01901 and No. 20H05852. M. C. acknowledges support from the National Science Foundation with Grants No. PHY-2010970 and No. OAC-2117997. S. S. acknowledges support from the G.R.E.A.T research environment, funded by *Vetenskapsrådet*, the Swedish Research Council, Project No. 2016-06012. M. G., P. M., and R. S. acknowledge the partial support of this research by Grant 19-12-00369 from the Russian Science Foundation. S. B. acknowledges financial support by the European Research Council for the ERC Starting grant *MessMapp*, under Contract No. 949555. The research of A. G.-Y. is supported by the EU via ERC Grant No. 725161, the ISF GW excellence center, an IMOS space infrastructure grant and BSF/Transformative and GIF grants, as well as The Benozio Endowment Fund for the Advancement of Science, the Deloro Institute for Advanced Research in Space and Optics, The Veronika A. Rabl Physics Discretionary Fund, Minerva, Yeda-Sela, and the Schwartz/Reisman Collaborative Science Program; A. G.-Y. is the incumbent of the The Arlyn Imberman Professorial Chair. E. C. K. acknowledges support from the G. R. E. A. T research environment funded by

Vetenskapsrådet, the Swedish Research Council, under Project No. 2016-06012, and support from The Wenner-Gren Foundations. M. R. has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Program (Grant Agreement No. 759194—USNAC). N. L. S. is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) via the Walter Benjamin program—461903330. Based on observations obtained with the Samuel Oschin Telescope 48-inch and the 60-inch Telescope at the Palomar Observatory as part of the Zwicky Transient Facility project. ZTF is supported by the National Science Foundation under Grant No. AST-1440341 (until 2020 December 1) and No. AST-2034437 and a collaboration including Caltech, IPAC, the Weizmann Institute for Science, the Oskar Klein Center at Stockholm University, the University of Maryland, Deutsches Elektronen-Synchrotron and Humboldt University, the TANGO Consortium of Taiwan, the University of Wisconsin at Milwaukee, Trinity College Dublin, Lawrence Livermore National Laboratories, IN2P3, University of Warwick, Ruhr University Bochum and Northwestern University. Operations are conducted by COO, IPAC, and UW. This work was supported by the GROWTH (Global Relay of Observatories Watching Transients Happen) project funded by the National Science Foundation Partnership in International Research and Education program under Grant No. 1545949. GROWTH is a collaborative project between California Institute of Technology (USA), Pomona College (USA), San Diego State University (USA), Los Alamos National Laboratory (USA), University of Maryland College Park (USA), University of Wisconsin Milwaukee (USA), Tokyo Institute of Technology (Japan), National Central University (Taiwan), Indian Institute of Astrophysics (India), Inter-University Center for Astronomy and Astrophysics (India), Weizmann Institute of Science (Israel), The Oskar Klein Centre at Stockholm University (Sweden), Humboldt University (Germany). This work is based on observations with the eROSITA telescope on board the SRG observatory. The SRG observatory was built by Roskosmos in the interests of the Russian Academy of Sciences represented by its Space Research Institute (IKI) in the framework of the Russian Federal Space Program, with the participation of the Deutsches Zentrum für Luft- und Raumfahrt (DLR). The SRG/eROSITA X-ray telescope was built by a consortium of German Institutes led by MPE, and supported by DLR. The SRG spacecraft was designed, built, launched, and is operated by the Lavochkin Association and its subcontractors. The science data are downlinked via the Deep Space Network Antennae in Bear Lakes, Ussurijsk, and Baykonur, funded by Roskosmos. The eROSITA data used in this work were processed using the eSASS software system developed by the German eROSITA consortium and proprietary data reduction and analysis software developed by the Russian

eROSITA Consortium. This work was supported by the Australian government through the Australian Research Council's Discovery Projects funding scheme (DP200102471). This work includes data products from the Near-Earth Object Wide-field Infrared Survey Explorer (NEOWISE), which is a project of the Jet Propulsion Laboratory/California Institute of Technology. NEOWISE is funded by the National Aeronautics and Space Administration. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. Based on observations made with the Nordic Optical Telescope, owned in collaboration by the University of Turku and Aarhus University, and operated jointly by Aarhus University, the University of Turku and the University of Oslo, representing Denmark, Finland and Norway, the University of Iceland and Stockholm University at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofísica de Canarias.

S. R. first identified AT2019fdr as a candidate neutrino source, performed the SED and the dust echo analysis and was the primary author of the manuscript. A. F., M. K., R. St. and S. v. V. contributed to the manuscript, the data analysis, and the source modeling. A. F. and M. K. have initiated the ZTF neutrino follow-up program. C. L., K. M. and W. W. performed the neutrino production analysis. J. C. A. M.-J. contributed the VLA observations. M. Gi. performed the SRG/eROSITA observations and data analysis. S. B. and S. Ga. analyzed the Fermi data. S. R. analyzed the Swift-XRT data. S. A., K. D., and S. R. performed and analyzed the P200 IR observations. S. Ge. and S. S. requested and reduced the Swift-UVOT data. V. S. P. contributed the black hole mass estimate. E. Z. contributed to the neutrino event rate calculation. M. G. analyzed the archival radio data. S. S. K and B. T. Z. performed parts of the neutrino production analysis. V. K. and N. L. S. contributed PTF forced photometry. C. B., T. S., and J. S. ruled out other candidates in the follow-up. T. A., M. W. C., M. M. K., and L. P. S. enabled ZTF ToO observations. J. Ne., S. R., and R. St. developed the analysis pipeline. V. B., J. No. and J. v. S. developed AMPEL, and contributed to the ToO analysis infrastructure. E. C. B., S. B. C., R. D., M. J. G., R. R. L., B. R., M. R., and R. W. contributed to the implementation of ZTF. P. M. and R. Su. contributed to the implementation of SRG/eROSITA. S. F., S. Ge., A. G.-Y., A. K. H. K., E. K., and D. A. P. contributed comments and to discussions. All authors reviewed the contents of the manuscript.

*Corresponding author.
simeon.reusch@desy.de.

†Corresponding author.
marek.kowalski@desy.de

- [1] M. Aartsen *et al.* (IceCube Collaboration), *J. Instrum.* **12**, P03012 (2017).
- [2] M. Aartsen *et al.* (IceCube Collaboration), *Phys. Rev. Lett.* **115**, 081102 (2015).
- [3] M. Aartsen *et al.* (IceCube Collaboration), *Phys. Rev. Lett.* **113**, 101101 (2014).
- [4] M. Aartsen *et al.* (IceCube Collaboration), *Science* **342**, 1242856 (2013).
- [5] M. Ahlers and F. Halzen, *Prog. Part. Nucl. Phys.* **102**, 73 (2018).
- [6] M. Kowalski, *Nat. Astron.* **5**, 732 (2021).
- [7] M. Aartsen *et al.* (IceCube Collaboration), *Science* **361**, eaat1378 (2018).
- [8] M. G. Aartsen *et al.* (IceCube Collaboration), *Science* **361**, 147 (2018).
- [9] S. Garrappa *et al.* (Fermi LAT Collaboration), *Astrophys. J.* **880**, 103 (2019).
- [10] R. Stein *et al.*, *Nat. Astron.* **5**, 510 (2021).
- [11] M. Aartsen *et al.* (IceCube Collaboration), *Phys. Rev. Lett.* **124**, 051103 (2020).
- [12] V. S. Berezinsky, *Proceedings of the International Conference "Neutrino '77"* (1977).
- [13] D. Eichler, *Astrophys. J.* **232**, 106 (1979).
- [14] F. W. Stecker, C. Done, M. H. Salamon, and P. Sommers, *Phys. Rev. Lett.* **66**, 2697 (1991).
- [15] K. Mannheim, *Astron. Astrophys.* **269**, 67 (1993).
- [16] A. Szabo and R. Protheroe, *Astropart. Phys.* **2**, 375 (1994).
- [17] K. Murase, *Neutrino Astronomy* (World Scientific, 2017), p. 15–31, [10.1142/9789814759410_0002](https://doi.org/10.1142/9789814759410_0002).
- [18] K. Murase, D. Guetta, and M. Ahlers, *Phys. Rev. Lett.* **116**, 071101 (2016).
- [19] Andrea Palladino and Walter Winter, *Astron. Astrophys.* **615**, A168 (2018).
- [20] I. Bartos, D. Veske, M. Kowalski, Z. Márka, and S. Márka, *Astrophys. J.* **921**, 45 (2021).
- [21] K. Murase, S. S. Kimura, and P. Mészáros, *Phys. Rev. Lett.* **125**, 011101 (2020).
- [22] G. R. Farrar and A. Gruzinov, *Astrophys. J.* **693**, 329 (2009).
- [23] K. Murase, *γ -Ray Bursts. Proceedings, Conference, (Nanjing, P. R. China, 2008); [AIP Conf. Proc. **1065**, 201 (2008)]*.
- [24] X.-Y. Wang, R.-Y. Liu, Z.-G. Dai, and K. S. Cheng, *Phys. Rev. D* **84**, 081301(R) (2011).
- [25] G. R. Farrar and T. Piran, [arXiv:1411.0704](https://arxiv.org/abs/1411.0704).
- [26] X.-Y. Wang and R.-Y. Liu, *Phys. Rev. D* **93**, 083005 (2016).
- [27] L. Dai and K. Fang, *Mon. Not. R. Astron. Soc.* **469**, 1354 (2017).
- [28] N. Senno, K. Murase, and P. Mészáros, *Astrophys. J.* **838**, 3 (2017).
- [29] Claire Guépin, Kumiko Kotera, Enrico Barausse, Ke Fang, and Kohta Murase, *Astron. Astrophys.* **616**, A179 (2018); **636**, C3(E) (2020).
- [30] C. Lunardini and W. Winter, *Phys. Rev. D* **95**, 123001 (2017).
- [31] L. Dai and K. Fang, *Mon. Not. R. Astron. Soc.* **469**, 1354 (2017).
- [32] B. T. Zhang, K. Murase, F. Oikonomou, and Z. Li, *Phys. Rev. D* **96**, 063007 (2017).

- [33] D. Biehl, D. Boncioli, C. Lunardini, and W. Winter, *Sci. Rep.* **8**, 10828 (2018).
- [34] K. Hayasaki and R. Yamazaki, *Astrophys. J.* **886**, 114 (2019).
- [35] W. Winter and C. Lunardini, *Nat. Astron.* **5**, 472 (2021).
- [36] K. Murase, S. S. Kimura, B. T. Zhang, F. Oikonomou, and M. Petropoulou, *Astrophys. J.* **902**, 108 (2020).
- [37] R.-Y. Liu, S.-Q. Xi, and X.-Y. Wang, *Phys. Rev. D* **102**, 083028 (2020).
- [38] K. Fang, B. D. Metzger, I. Vurm, E. Aydi, and L. Chomiuk, *Astrophys. J.* **904**, 4 (2020).
- [39] W. Winter and C. Lunardini, *Proc. Sci., ICRC2021 (2021)* 997.
- [40] K. Hayasaki, *Nat. Astron.* **5**, 436 (2021).
- [41] G. Cannizzaro *et al.*, *Mon. Not. R. Astron. Soc.* **504**, 792 (2021).
- [42] Y. Cendes, K. D. Alexander, E. Berger, T. Eftekhari, P. K. G. Williams, and R. Chornock, *Astrophys. J.* **919**, 127 (2021).
- [43] P. Mohan, T. An, Y. Zhang, J. Yang, X. Yang, and A. Wang, *Astrophys. J.* **927**, 74 (2022).
- [44] T. Matsumoto, T. Piran, and J. H. Krolik, *Mon. Not. R. Astron. Soc.* **511**, 5085 (2022).
- [45] S. Frederick *et al.*, *Astrophys. J.* **920**, 56 (2021).
- [46] E. C. Bellm *et al.*, *Publ. Astron. Soc. Pac.* **131**, 018002 (2019).
- [47] M. J. Graham *et al.*, *Publ. Astron. Soc. Pac.* **131**, 078001 (2019).
- [48] R. Dekany *et al.*, *Publ. Astron. Soc. Pac.* **132**, 038001 (2020).
- [49] R. Stein, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27865.gcn3>.
- [50] R. Stein, S. Reusch, and J. Necker, nuztf: v2.2.1, 10.5281/zenodo.5567047 (2021).
- [51] J. Nordin *et al.*, *Astron. Astrophys.* **631**, A147 (2019).
- [52] M. Kowalski and A. Mohr, *Astropart. Phys.* **27**, 533 (2007).
- [53] S. Reusch, R. Stein, A. Franckowiak, and S. Gezari, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27872.gcn3>.
- [54] J. Nordin, V. Brinnel, M. Giomi, J. V. Santen, A. Gal-Yam, O. Yaron, and S. Schulze, *TNS Disc. Rep.* **2019–771**, 1 (2019).
- [55] N. M. Law *et al.*, *Publ. Astron. Soc. Pac.* **121**, 1395 (2009).
- [56] D. Calzetti, L. Armus, R. C. Bohlin, A. L. Kinney, J. Koornneef, and T. Storchi-Bergmann, *Astrophys. J.* **533**, 682 (2000).
- [57] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.128.221101> for details on the data reduction and analysis, which includes Refs. [58–175].
- [58] S. Reusch, simeonreusch/at2019fdr: V1.0 Release, (2021).
- [59] M. Rigault, ztfquery, a python tool to access ztf data, (2018).
- [60] Gaia Collaboration, *Astron. Astrophys.* **616**, A1 (2018).
- [61] H. Ayala, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27873.gcn3>.
- [62] V. Savchenko, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27866.gcn3>.
- [63] S. Sazonov *et al.*, *Mon. Not. R. Astron. Soc.* **508**, 3820 (2021).
- [64] N. Ben Bekhti *et al.* (HI4PI Collaboration), *Astron. Astrophys.* **594**, A116 (2016).
- [65] U. S. S. D. Centre, Swift xrt data products generator, https://www.swift.ac.uk/user_objects (2020).
- [66] P. A. Evans *et al.*, *Mon. Not. R. Astron. Soc.* **397**, 1177 (2009).
- [67] M. Arida and E. Sabol, Heasarc webpimms, <https://heasarc.gsfc.nasa.gov/cgi-bin/Tools/w3pimms/w3pimms.pl> (2020).
- [68] S. Reusch, ztffps v1.0.3, 10.5281/zenodo.5119152 (2020).
- [69] M. Rigault, ztflc, <https://github.com/mickaelrigault/ztflc> (2020).
- [70] A. A. Breeveld, W. Landsman, S. T. Holland, P. Roming, N. P. M. Kuin, and M. J. Page, in *American Institute of Physics Conference Series, American Institute of Physics Conference Series* Vol. 1358, edited by J. E. McEnery, J. L. Racusin, and N. Gehrels (2011), pp. 373–376.
- [71] K. De *et al.*, *Publ. Astron. Soc. Pac.* **132**, 025001 (2020).
- [72] M. F. Skrutskie *et al.*, *Astron. J.* **131**, 1163 (2006).
- [73] C. Y. Peng, L. C. Ho, C. D. Impey, and H.-W. Rix, *Astron. J.* **124**, 266 (2002).
- [74] L. Bradley *et al.*, astropy/photutils: 1.0.0 (2020).
- [75] D. G. York *et al.*, *Astrophys. J.* **120**, 1579 (2000).
- [76] S. van Velzen, T. W. S. Holoiën, F. Onori, T. Hung, and I. Arcavi, *Space Sci. Rev.* **216**, 124 (2020).
- [77] D. C. Martin *et al.*, *Astrophys. J.* **619**, L1 (2005).
- [78] C. Million, S. W. Fleming, B. Shiao, M. Seibert, P. Loyd, M. Tucker, M. Smith, R. Thompson, and R. L. White, *Astrophys. J.* **833**, 292 (2016).
- [79] C. Stoughton *et al.*, *Astron. J.* **123**, 485 (2002).
- [80] A. Lawrence *et al.*, *Mon. Not. R. Astron. Soc.* **379**, 1599 (2007).
- [81] C. Conroy and J. E. Gunn, *Astrophys. J.* **712**, 833 (2010).
- [82] D. Foreman-Mackey, J. Sick, and B. Johnson, python-fsps: Python bindings to FSPP (v0.1.1) (2014).
- [83] D. O. Jones, D. M. Scolnic, and S. A. Rodney, PythonPhot: Simple DAOPHOT-type photometry in Python (2015), ascl:1501.010.
- [84] F. Masci, ICORE: Image Co-addition with Optional Resolution Enhancement (2013), ascl:1302.010.
- [85] M. Lacy *et al.*, *Publ. Astron. Soc. Pac.* **132**, 035001 (2020).
- [86] R. Barniol Duran, E. Nakar, and T. Piran, *Astrophys. J.* **772**, 78 (2013).
- [87] G. Morlino and D. Caprioli, *Astron. Astrophys.* **538**, A81 (2012).
- [88] T. Eftekhari, E. Berger, B. A. Zauderer, R. Margutti, and K. D. Alexander, *Astrophys. J.* **854**, 86 (2018).
- [89] A. Horesh *et al.*, *Mon. Not. R. Astron. Soc.* **436**, 1258 (2013).
- [90] A. Generozov, P. Mimica, B. D. Metzger, N. C. Stone, D. Giannios, and M. A. Aloy, *Mon. Not. R. Astron. Soc.* **464**, 2481 (2017).
- [91] M. Newville, T. Stensitzki, D. B. Allen, and A. Ingargiola, LMFIT: Non-Linear Least-Square Minimization and Curve-Fitting for Python (2021).
- [92] B. Kyle, extinction, 10.5281/zenodo.804967 (2016).
- [93] M. Vestergaard and B. M. Peterson, *Astrophys. J.* **641**, 689 (2006).
- [94] H. Guo, Y. Shen, and S. Wang, PyQSOFit: Python code to fit the spectrum of quasars (2018), ascl:1809.008.
- [95] E. L. Fitzpatrick, *Publ. Astron. Soc. Pac.* **111**, 63 (1999).

- [96] D. J. Schlegel, D. P. Finkbeiner, and M. Davis, *Astrophys. J.* **500**, 525 (1998).
- [97] T. A. Boroson and R. F. Green, *Astrophys. J. Suppl. Ser.* **80**, 109 (1992).
- [98] Y. Shen *et al.*, *Astrophys. J. Suppl. Ser.* **194**, 45 (2011).
- [99] H. Guo, X. Liu, Y. Shen, A. Loeb, T. Monroe, and J. X. Prochaska, *Mon. Not. R. Astron. Soc.* **482**, 3288 (2019).
- [100] Y. Shen *et al.*, *Astrophys. J. Suppl. Ser.* **241**, 34 (2019).
- [101] R. J. McLure and J. S. Dunlop, *Mon. Not. R. Astron. Soc.* **331**, 795 (2002).
- [102] M. T. Patterson, E. C. Bellm, B. Rusholme, F. J. Masci, M. Juric, K. Simon Krughoff, V. Zach Golkhou, M. J. Graham, S. R. Kulkarni, and G. Helou, *Publ. Astron. Soc. Pac.* **131**, 018001 (2019).
- [103] F. J. Masci *et al.*, *Publ. Astron. Soc. Pac.* **131**, 018003 (2019).
- [104] E. Blaufuss, GCN Circular (2019), <https://gcn.gsfc.nasa.gov/gcn3/24378.gcn3>.
- [105] R. Stein, A. Franckowiak, J. van Santen, L. Rauch, M. M. Kasliwal, I. Andreoni, T. Ahumada, M. Coughlin, L. P. Singer, and S. Anand, *Astronomer's Telegram* (2019), <https://www.astronomerstelegam.org/?read=12730>.
- [106] E. Blaufuss, GCN Circular (2019), <https://gcn.gsfc.nasa.gov/gcn3/24910.gcn3>.
- [107] R. Stein *et al.*, *Astronomer's Telegram* (2019), <https://www.astronomerstelegam.org/?read=12879>.
- [108] R. Stein, GCN Circular (2019), <https://gcn.gsfc.nasa.gov/gcn3/25225.gcn3>.
- [109] R. Stein, A. Franckowiak, M. M. Kasliwal, I. Andreoni, M. Coughlin, L. P. Singer, F. Masci, and S. van Velzen, *Astronomer's Telegram* (2019), <https://www.astronomerstelegam.org/?read=12974>.
- [110] E. Blaufuss, GCN Circular (2019), <https://gcn.gsfc.nasa.gov/gcn3/25806.gcn3>.
- [111] R. Stein, A. Franckowiak, M. Kowalski, and M. Kasliwal, *Astronomer's Telegram* (2019), <https://www.astronomerstelegam.org/?read=13125>.
- [112] R. Stein, A. Franckowiak, M. Kowalski, and M. Kasliwal, GCN Circular (2019), <https://gcn.gsfc.nasa.gov/gcn3/25824.gcn3> **25824**.
- [113] R. Stein, GCN Circular (2019), <https://gcn.gsfc.nasa.gov/gcn3/25913.gcn3>.
- [114] R. Stein, A. Franckowiak, J. Necker, S. Gezari, and S. v. Velzen, *Astronomer's Telegram* (2019), <https://www.astronomerstelegam.org/?read=13160>.
- [115] R. Stein, A. Franckowiak, J. Necker, S. Gezari, and S. van Velzen, GCN Circular (2019), <https://gcn.gsfc.nasa.gov/gcn3/25929.gcn3>.
- [116] R. Stein, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/26655.gcn3>.
- [117] R. Stein and S. Reusch, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/26667.gcn3>.
- [118] R. Stein, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/26696.gcn3>.
- [119] S. Reusch and R. Stein, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/26747.gcn3>.
- [120] C. Lagunas Gualda, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/26802.gcn3>.
- [121] S. Reusch and R. Stein, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/26813.gcn3>.
- [122] S. Reusch and R. Stein, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/26816.gcn3>.
- [123] C. Lagunas Gualda, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27719.gcn3>.
- [124] S. Reusch, R. Stein, and A. Franckowiak, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27721.gcn3>.
- [125] R. Stein, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27865.gcn3>.
- [126] S. Reusch, R. Stein, A. Franckowiak, and S. Gezari, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27872.gcn3>.
- [127] S. Reusch, R. Stein, A. Franckowiak, J. Sollerman, T. Schweyer, and C. Barbarino, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27910.gcn3>.
- [128] S. Reusch, R. Stein, A. Franckowiak, J. Necker, J. Sollerman, C. Barbarino, and T. Schweyer, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27980.gcn3>.
- [129] M. Santander, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/27997.gcn3>.
- [130] S. Reusch, R. Stein, and A. Franckowiak, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28005.gcn3>.
- [131] E. Blaufuss, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28433.gcn3>.
- [132] S. Reusch, R. Stein, A. Franckowiak, I. Andreoni, and M. Coughlin, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28441.gcn3>.
- [133] S. Reusch, R. Stein, A. Franckowiak, S. Schulze, and J. Sollerman, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28465.gcn3>.
- [134] C. Lagunas Gualda, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28504.gcn3>.
- [135] S. Reusch, R. Stein, S. Weimann, and A. Franckowiak, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28520.gcn3>.
- [136] C. Lagunas Gualda, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28532.gcn3>.
- [137] S. Weimann, R. Stein, S. Reusch, and A. Franckowiak, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28551.gcn3>.
- [138] M. Santander, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28575.gcn3>.
- [139] S. Reusch, S. Weimann, R. Stein, and A. Franckowiak, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28609.gcn3>.
- [140] C. Lagunas Gualda, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28715.gcn3>.
- [141] R. Stein, S. Reusch, S. Weimann, and M. Coughlin, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28757.gcn3>.
- [142] C. Lagunas Gualda, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28969.gcn3>.
- [143] S. Weimann, R. Stein, S. Reusch, and A. Franckowiak, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/28989.gcn3>.
- [144] C. Lagunas Gualda, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/29012.gcn3>.
- [145] S. Reusch, S. Weimann, R. Stein, and A. Franckowiak, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/29031.gcn3>.
- [146] E. Blaufuss, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/29120.gcn3>.

- [147] R. Stein, S. Weimann, S. Reusch, and A. Franckowiak, GCN Circular (2020), <https://gcn.gsfc.nasa.gov/gcn3/29172.gcn3>.
- [148] C. Lagunas Gualda, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/29454.gcn3>.
- [149] S. Reusch, S. Weimann, R. Stein, M. Coughlin, and A. Franckowiak, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/29461.gcn3>.
- [150] M. Santander, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/29976.gcn3>.
- [151] R. Stein, S. Weimann, J. Necker, S. Reusch, and A. Franckowiak, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/29999.gcn3>.
- [152] M. Santander, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/30342.gcn3>.
- [153] J. Necker, R. Stein, S. Weimann, S. Reusch, and A. Franckowiak, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/30349.gcn3>.
- [154] M. Santander, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/30627.gcn3>.
- [155] R. Stein, S. Weimann, S. Reusch, J. Necker, A. Franckowiak, and M. Coughlin, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/30644.gcn3>.
- [156] M. Lincetto, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/30862.gcn3>.
- [157] S. Weimann, S. Reusch, J. Necker, R. Stein, and A. Franckowiak, GCN Circular (2021), <https://gcn.gsfc.nasa.gov/gcn3/30870.gcn3>.
- [158] S. S. Kimura, K. Murase, and K. Toma, *Astrophys. J.* **806**, 159 (2015).
- [159] J. C. McKinney, L. Dai, and M. J. Avara, *Mon. Not. R. Astron. Soc.* **454**, L6 (2015).
- [160] E. Takeo, K. Inayoshi, K. Ohsuga, H. R. Takahashi, and S. Mineshige, *Mon. Not. R. Astron. Soc.* **488**, 2689 (2019).
- [161] A. Mücke, R. Engel, J. P. Rachen, R. J. Protheroe, and T. Stanev, *Comput. Phys. Commun.* **124**, 290 (2000).
- [162] Y. Io and T. K. Suzuki, *Astrophys. J.* **780**, 46 (2014).
- [163] Y.-F. Jiang, J. M. Stone, and S. W. Davis, *Astrophys. J.* **784**, 169 (2014).
- [164] H. R. Takahashi, K. Ohsuga, T. Kawashima, and Y. Sekiguchi, *Astrophys. J.* **826**, 23 (2016).
- [165] L. E. Strubbe and E. Quataert, *Mon. Not. R. Astron. Soc.* **400**, 2070 (2009).
- [166] M. C. Miller, *Astrophys. J.* **805**, 83 (2015).
- [167] B. D. Metzger and N. C. Stone, *Mon. Not. R. Astron. Soc.* **461**, 948 (2016).
- [168] L. Dai, J. C. McKinney, N. Roth, E. Ramirez-Ruiz, and M. C. Miller, *Astrophys. J.* **859**, L20 (2018).
- [169] A. Mummery and S. A. Balbus, *Mon. Not. R. Astron. Soc.* **504**, 4730 (2021).
- [170] A. Mummery, [arXiv:2104.06212](https://arxiv.org/abs/2104.06212).
- [171] A. Mummery and S. A. Balbus, *Mon. Not. R. Astron. Soc.* **505**, 1629 (2021).
- [172] C. S. Kochanek, *Mon. Not. R. Astron. Soc.* **461**, 371 (2016).
- [173] M. J. Rees, *Nature (London)* **333**, 523 (1988).
- [174] F. De Colle, J. Guillochon, J. Naiman, and E. Ramirez-Ruiz, *Astrophys. J.* **760**, 103 (2012).
- [175] K. Murase, Y. Inoue, and C. D. Dermer, *Phys. Rev. D* **90**, 023007 (2014).
- [176] A. Gal-Yam, *Annu. Rev. Astron. Astrophys.* **57**, 305 (2019).
- [177] R. Chornock, P. K. Blanchard, S. Gomez, G. Hosseinzadeh, and E. Berger, *TNS Class. Rep.* **2019-1016**, 1 (2019).
- [178] T. Pitik, I. Tamborra, C. R. Angus, and K. Auchettl, [arXiv:2110.06944](https://arxiv.org/abs/2110.06944).
- [179] P. W. A. Roming *et al.*, *Space Sci. Rev.* **120**, 95 (2005).
- [180] N. Gehrels *et al.*, *Astrophys. J.* **611**, 1005 (2004).
- [181] D. N. Burrows *et al.*, *Space Sci. Rev.* **120**, 165 (2005).
- [182] P. Predehl *et al.*, *Astron. Astrophys.* **647**, A1 (2021).
- [183] R. Sunyaev *et al.*, *Astron. Astrophys.* **656**, A132 (2021).
- [184] R. Saxton, S. Komossa, K. Auchettl, and P. G. Jonker, *Space Sci. Rev.* **216**, 85 (2020).
- [185] K. M. Leighly, *Astrophys. J. Suppl. Ser.* **125**, 317 (1999).
- [186] S. Vaughan, J. Reeves, R. Warwick, and R. Edelson, *Mon. Not. R. Astron. Soc.* **309**, 113 (1999).
- [187] R. Margutti *et al.*, *Astrophys. J.* **864**, 45 (2018).
- [188] K. Barbary *et al.*, *Astrophys. J.* **690**, 1358 (2009).
- [189] B. T. Gänsicke, A. J. Levan, T. R. Marsh, and P. J. Wheatley, *Astrophys. J.* **697**, L129 (2009).
- [190] A. Mainzer *et al.*, *Astrophys. J.* **731**, 53 (2011).
- [191] E. L. Wright *et al.*, *Astron. J.* **140**, 1868 (2010).
- [192] J. C. Wilson *et al.*, in *Instrument Design and Performance for Optical/Infrared Ground-based Telescopes, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 4841*, edited by M. Iye and A. F. M. Moorwood (SPIE, 2003), pp. 451–458.
- [193] S. van Velzen, A. J. Mendez, J. H. Krolik, and V. Gorjian, *Astrophys. J.* **829**, 19 (2016).
- [194] S. Dong *et al.*, *Science* **351**, 257 (2016).
- [195] G. Leloudas *et al.*, *Nat. Astron.* **1**, 0002 (2017).
- [196] E. Kankare *et al.*, *Nat. Astron.* **1**, 865 (2017).
- [197] E. C. Kool *et al.*, *Mon. Not. R. Astron. Soc.* **498**, 2167 (2020).
- [198] S. Mattila *et al.*, *Science* **361**, 482 (2018).
- [199] R. A. Perley, C. J. Chandler, B. J. Butler, and J. M. Wrobel, *Astrophys. J. Lett.* **739**, L1 (2011).
- [200] W. B. Atwood *et al.*, *Astrophys. J.* **697**, 1071 (2009).
- [201] S. van Velzen *et al.*, [arXiv:2111.09391](https://arxiv.org/abs/2111.09391).
- [202] N. L. Strotjohann, M. Kowalski, and A. Franckowiak, *Astron. Astrophys.* **622**, L9 (2019).
- [203] S. van Velzen *et al.*, *Astrophys. J.* **908**, 4 (2021).
- [204] E. Blaufuss, T. Kintscher, L. Lu, and C. F. Tung, *Proc. Sci., ICRC2019 (2019)* 1021.
- [205] Y.-F. Jiang, J. M. Stone, and S. W. Davis, *Astrophys. J.* **880**, 67 (2019).
- [206] A. Mummery, *Mon. Not. R. Astron. Soc.* **504**, 5144 (2021).
- [207] T. Ryu, J. Krolik, and T. Piran, *Astrophys. J.* **904**, 73 (2020).
- [208] C.-H. Chan, T. Piran, J. H. Krolik, and D. Saban, *Astrophys. J.* **881**, 113 (2019).
- [209] I. Arcavi *et al.*, *Astrophys. J.* **793**, 38 (2014).
- [210] K. D. French, T. Wevers, J. Law-Smith, O. Graur, and A. I. Zabludoff, *Space Sci. Rev.* **216**, 32 (2020).
- [211] N. C. Stone, A. Generozov, E. Vasiliev, and B. D. Metzger, *Mon. Not. R. Astron. Soc.* **480**, 5060 (2018).
- [212] IceCube Collaboration, All-sky point-source IceCube data: Years 2010-2012. Dataset, [10.21234/B4F04V](https://arxiv.org/abs/10.21234/B4F04V) (2018).
- [213] A. Albert *et al.* (ANTARES Collaboration), *Astrophys. J.* **920**, 50 (2021).