

International consensus on the most useful physical examination tests used by physiotherapists for patients with headache: A Delphi study

Luedtke, K.¹; Boissonnault, W.²; Caspersen, N.³; Castien, R.⁴; Chaibi, A.^{5,6}; Falla, D.^{7,8}; Fernández-de-las-Peñas, C.⁹; Hall, T.¹⁰; Hirsvang, J.R.³; Horre, T.¹¹; Hurley, D.¹²; Jull, G.¹³; Krøll, L.S.³; Madsen, B.K.³; Mallwitz, J.¹⁴; Miller, C.¹⁵; Schäfer, B.¹⁶; Schöttker-Königer, T.¹⁷; Starke, W.¹⁷; von Piekartz, H.¹⁸; Watson, D.¹⁹; Westerhuis, P.²⁰; May, A.¹.

Affiliations:

¹Department of Systems Neuroscience, University Medical Centre Hamburg-Eppendorf, Martinistr. 52, 20246 Hamburg, Germany

²University of Wisconsin-Madison, Department of Orthopedics and Rehabilitation, Wisconsin, USA

³Danish Headache Centre, Department of Neurology, Glostrup Hospital, University of Copenhagen, DK-2600 Glostrup, Denmark

⁴Department of General Practice and Elderly Care Medicine, EMGO+Institute for Health Care Research, VU University Medical Center, Amsterdam, The Netherlands

⁵Head and Neck Research Group, Research Centre, Akershus University Hospital, 1478, Oslo, Norway

⁶Institute of Clinical Medicine, Akershus University Hospital and University of Oslo, 1474, Nordbyhagen, Norway

⁷Pain Clinic, Center for Anaesthesiology, Emergency and Intensive Care Medicine, University Hospital Göttingen, Göttingen, Germany

⁸Department of Neurorehabilitation Engineering, Bernstein Focus Neurotechnology, Bernstein Center for Computational Neuroscience, University Medical Center Göttingen, Georg-August University, Göttingen, Germany

⁹Department of Physical Therapy, Occupational Therapy, Rehabilitation and Physical Medicine, Universidad Rey Juan Carlos, Alcorcón, Madrid, Spain

¹⁰School of Physiotherapy and Curtin Health Innovation Research, Curtin University, P.O. Box U1987, Perth, WA 6845, Australia

¹¹Physiotherapy practice Jensen&Team, Nienstedtener Marktplatz 29, 22609 Hamburg, Germany

¹²Physiotherapy, Birmingham Community Healthcare NHS Trust, Birmingham, England

¹³Physiotherapy, School of Health and Rehabilitation Sciences, The University of Queensland, Australia

¹⁴Spine Center "Am Michel", Ludwig-Erhard-Str. 18, 20459 Hamburg, Germany

¹⁵Physiotherapy Department, University Hospital Birmingham Foundation Trust, Birmingham B15 2TH

¹⁶Migraine and Headache Clinic Königstein im Taunus, Ölmühlenweg 31, 61462 Königstein im Taunus, Germany

¹⁷Hochschule für angewandte Wissenschaft und Kunst Hildesheim / Holzminen / Göttingen, Fakultät Soziale Arbeit und Gesundheit, Studiengang Ergotherapie, Logopädie und Physiotherapie, Goschentor 1, 31134 Hildesheim; Germany

¹⁸University of Applied Sciences Osnabrueck, Fakultät Wirtschafts- und Sozialwissenschaften, Caprivistr. 30a, 49076 Osnabrück, Germany

¹⁹Watson Headache Institute, GPO Box 4714, Sydney, NSW 2001, Australia

²⁰Heimlisbergstr. 47, 4513 Langendorf, Switzerland

Funding: Deutsche Migräne- und Kopfschmerzgesellschaft (DMKG)

Corresponding author:

Kerstin Luedtke, PhD
Department of Systems Neuroscience
University Medical Centre Hamburg-Eppendorf
Martinistr. 52,
20246 Hamburg
Germany

Telephone: +49 40 741058943

International consensus on the most useful physical examination tests used by physiotherapists for patients with headache: A Delphi study

Abstract

Background

A wide range of physical tests have been published for use in the assessment of musculoskeletal dysfunctions in patients with headache. Which tests are used depends on a physiotherapist's clinical and scientific background as there is little guidance on the most clinically useful tests.

Objectives

To identify which physical examination tests international experts in physiotherapy consider the most clinically useful for the assessment of patients with headache.

Design / Methods

Delphi survey with pre-specified procedures based on a systematic search of the literature for physical examination tests proposed for the assessment of musculoskeletal dysfunctions in patients with headache.

Results

Seventeen experts completed all three rounds of the survey. Fifteen tests were included in round one with eleven additional tests suggested by the experts. Finally eleven physical examination tests were considered clinically useful: manual joint palpation, the cranio-cervical flexion test, the cervical flexion-rotation test, active range of cervical movement, head forward position, trigger point palpation, muscle tests of the shoulder girdle, passive physiological intervertebral movements, reproduction and resolution of headache symptoms, screening of the thoracic spine, and combined movement tests.

Conclusions

Eleven tests are suggested as a minimum standard for the physical examination of musculoskeletal dysfunctions in patients with headache.

Key words: headache, migraine, Delphi, consensus, examination, physical therapy.

Introduction

The lifetime prevalence of any headache in the European population is more than 90% indicating the relevance and burden of headaches (Steiner et al. 2014). A recent study reported that tension-type headache (TTH) and migraine were also the most common recurrent headaches with prevalences of 20.77% and 14.7%, respectively (Vos et al. 2012). Headache compromises a person's quality of life and reduces their functional capability as well as their work capacity (D'Amico et al. 2013; Raggi et al. 2013). This results in high socioeconomic cost mainly due to days off work (Burton et al. 2002; Pradalier et al. 2004; Berg and Stovner 2005; Bloudek et al. 2012; Lanteri-Minet 2014).

There is ongoing scientific debate about the contribution of cervical neuromusculoskeletal dysfunction to headache. Although the results of a systematic review suggest that there is probably no influence of musculoskeletal dysfunctions in the pathogenesis of migraine (Robertson and Morris 2008), migraine patients have a high prevalence of neck pain (Ashina et al. 2015) and muscle or joint dysfunctions have been hypothesised to act as triggers for migraine attacks (Vincent 2011). In TTH patients, muscle tenderness and trigger points are consistent findings (Fernández-de-las-Peñas et al. 2007; Abboud et al. 2013), while the evidence for muscle tension is conflicting (Bendtsen and Fernández-de-la-Peñas 2011).

Cervicogenic headache (CGH) is the most likely sub-group of headache to present with musculoskeletal dysfunction in the neck (Sjaastad et al. 1998; Zito et al. 2006; Jull et al. 2007). The IHS classification version III beta details the clinical features of CGH under section 11.2.1 (Headache Classification Committee of the International Headache Society (IHS) 2013). However, the diagnosis CGH itself is most controversial: Some strongly argue for the neck as a potential headache generator (Bogduk 1992; Sjaastad et al. 1998; Antonaci et al. 2001; Bogduk and Govind 2009; Becker 2010) but others doubt the existence of CGH and propose that the reported prevalence, estimated e.g. as 4.1% in the general population (Sjaastad and Bakketeig 2008), is due to misdiagnosed migraine or TTH (Pöllmann et al. 1997; Leone et al. 1998).

Patients suffering from headache commonly use self-management strategies such as self-massage, posture correction, stretching and exercises to reduce symptoms (Haque et al. 2012) and seek to reduce medication intake by using additional non-pharmacological therapies, most commonly physiotherapy (Chaibi et al. 2011; Chaibi and Russell 2012, 2014; Kristoffersen et al. 2012). A physiotherapist will conduct a patient interview to record the headache history, its temporal pattern, symptom behaviour, the patient's medication intake and other relevant health history to ensure that the headache is benign and to classify the type of headache. In order to decide whether and which type of physiotherapy interventions (eg. manual therapy, exercise,

soft tissue techniques, or postural correction) are likely to influence the patient's complaints, the subsequent physical examination focuses on the assessment for the presence (or absence) of musculoskeletal dysfunction.

A wide range of physical examination tests have been published for the assessment of cervical musculoskeletal dysfunctions in patients with headache. These include measurement of head/neck posture (Fernández-de-Las-Peñas et al. 2006b), range of cervical motion (Zito et al. 2006; Jull et al. 2007) including the flexion-rotation test (Hall and Robinson 2004) manual examination to detect painful cervical joint dysfunction (Zito et al. 2006; Jull et al. 2007), tests of cervical muscle function (Zito et al. 2006; Jull et al. 2007), tests for temporo-mandibular dysfunction (Caspersen et al. 2013; Harry von Piekartz 2015), palpation for trigger points (Fernández-de-Las-Peñas et al. 2006b) and many others. Currently, there are no published guidelines to support the decision on the most important physical examination techniques. Individual studies have evaluated tests for only one specific headache type (TTH, migraine or CGH) (i.e. Kidd and Nelson 1993; Jensen and Rasmussen 1996; Ashina et al. 1999; Calandre et al. 2006; Fernández-de-las-Peñas et al. 2006; Fernández-de-Las-Peñas et al. 2006b, 2007; Couppé et al. 2007; Bevilaqua-Grossi et al. 2009), or within specific sub-populations (e.g. with associated facial or neck pain (this includes -by definition- all studies on CGH). While this serves as an indicator for clinical usefulness of specific tests in specific populations, it does not inform on the usefulness of tests in other headache patients.

This Delphi survey was conducted to identify physical examination tests that international experts in physiotherapy consider the most clinically useful when examining a patient with headache and to specify for which types of headache or in which clinical situation the tests would be useful.

Methods

Study design

A Delphi-Survey was designed and conducted following the recommendations of Hasson et al. (Hasson et al. 2000), and published examples of Delphi Surveys with similar research questions (Sinha et al. 2011; Chiarotto et al. 2014). The survey was guided by a systematic evaluation of the literature on physical examination tests proposed for the assessment of musculoskeletal dysfunctions in patients with headache. Ethical approval (No: PV5011) was granted by the local ethics authority (Aerztekammer Hamburg). The data collection was conducted between October 2014 and April 2015.

Selection of experts was based on personal contacts and a search for authors of peer-reviewed publications on physical examination tests in headache populations. There is no guideline to define who is an expert for a Delphi Survey or how many experts are required (Hsu and Sandford 2007). Twenty international experts (defined as physiotherapists with a special interest in headaches) from nine countries were identified and invited via email to participate. The letter of invitation included the researcher's details and information on the purpose of the survey. It further explained how the acquired data will be used to design subsequent survey rounds. Seventeen experts with backgrounds ranging from dominantly clinical to dominantly research-based, participated in the survey and completed all three rounds of the survey. Participants remained anonymous during all rounds of the survey with the aim that they can express opinions without influence of dominant characters (Murphy et al. 1998) . After the final survey round, all experts were invited to be a co-author of the publication and subsequently received a first draft of the survey report for revision and a final draft for approval prior to submission.

The survey procedures were pre-designed and included the following steps:

1.) Initially a list of physical examination tests was generated based on a systematic evaluation of the literature to identify any type of publication that focused on physical examination tests in headache populations conducted within a physiotherapy setting. PubMed, Cinahl, EMBASE and PEDro were searched for articles published prior to 01.10.2014 using the following search strategy:

((((headache[Title/Abstract] OR migraine[Title/Abstract]))) AND ((assessment OR measurement OR examination OR test))) AND ((dysfunction OR impairment))) AND ((strength OR length OR endurance OR sensitivity OR control OR mobility OR motion OR tension OR posture))) AND adult Filters: Humans

The literature search returned 3 reviews (Jensen and Rasmussen 1996; Robertson and Morris 2008; Abboud et al. 2013) and 43 additional articles that reported a total of 15 physical

examination tests (Lous and Olesen 1982; Langemark and Olesen 1987; Bovim 1992; Hatch et al. 1992; Garrett et al. 1993; Jensen et al. 1993; Kidd and Nelson 1993; Watson and Trott 1993; Jensen 1996; Lipchik et al. 1996; Sand et al. 1997; Zwart 1997; Ashina et al. 1999; Jull et al. 1999, 2007; Neufeld et al. 2000; Dumas et al. 2001; Mørk et al. 2003; Giacomini et al. 2004; Fernández-de-las-Peñas et al. 2006; Fernández-de-Las-Peñas et al. 2006a, 2006b, 2007, 2008; Sandrini et al. 2006; Zito et al. 2006; Amiri et al. 2007; Glaros et al. 2007, 2007; Ogince et al. 2007; Fernández-de-las-Peñas et al. 2008; Hall et al. 2008, 2010, 2010; Bevilaqua-Grossi et al. 2009; Peddireddy et al. 2009; Uthaikeup et al. 2009; Sohn et al. 2010, 2013; Stuginski-Barbosa et al. 2010; Melo et al. 2012; Watson and Drummond 2012; Marchand et al. 2014; Tali et al. 2014).

2.) In the first Delphi round, experts were asked to rate the clinical usefulness of the 15 tests (manual joint palpation, cranio-cervical flexion test, flexion-rotation test, active range of cervical movement, head forward position, trigger point palpation, muscle tests of the shoulder girdle, pressure pain thresholds, cervical slump test, skin-roll-test, electromyography, temporomandibular screening, goal oriented movements, cervical muscle strength, and joint position error) on a 5-point Likert scale (“extremely useful”, “useful”, “don’t know”, “probably not useful”, “definitely not useful”) for TTH, migraine and CGH patients. References were provided for each physical examination test detailing test procedures. In addition, the experts were invited to provide additional physical examination tests they considered as clinically relevant. The questionnaire used for the rating procedure was sent as a word document via email after it had been tested amongst work group members for comprehensibility. If individual ratings were missing or questionnaires were not returned, reminder emails were sent to clarify responses and to inquire whether the expert was still willing to continue the study.

3.) For the second round the group responses were analysed and fed back to the experts (via email) including the newly suggested tests. These were again rated for clinical usefulness. Since experts were from various backgrounds with a range of special interests and in order to reflect this variety in the results, tests that were rated as clinically useful by >50% of the participants were automatically included in the subsequent rounds, while tests that were rated as clinically useful by \leq 50% of participants during the first and second round were further investigated in subsequent rounds by inviting comments on special clinical situations for which this tests may be useful.

4.) A third round was conducted in the same manner as the second round to include all new information that emerged from round 2.

Data analysis

All tests rated as “definitely not useful” or “probably not useful” were excluded from subsequent Delphi rounds (based on the mode value across the experts). Tests that received an overall rating (mode value) of “don’t know” or that were rated as clinically useful by $\leq 50\%$ of participants during the first and second round required further investigation: experts were asked to indicate specific clinical situations in which they would use these tests. These responses were analysed in a qualitative manner and summarised in a pre-designed table.

RESULTS

Results from round 1:

The most common finding was that the majority of experts (>50%) used similar tests for patients with different headache diagnoses. Individual experts supported the decision not to distinguish between headache diagnoses, when deciding on physical examination tests, with the following statements: “Physical examination tests are conducted to identify musculoskeletal dysfunctions and to develop a treatment plan independent of the diagnosis”; “there is substantial overlap of musculoskeletal findings in different headache types”; “patients can have more than one type of headache”. Figures 1a-c present the ratings of clinical usefulness for each suggested physical examination test for each headache type.

By calculating the mode value across headache types for each suggested physical examination test, manual joint palpation, the cranio-cervical flexion test, the flexion-rotation test, active range of cervical movement, head forward position, trigger point palpation, muscle tests of the shoulder girdle, pressure pain thresholds, and cervical slump test, were considered useful/extremely useful. Rated as “probably not useful” or “definitely not useful” were: skin-roll-test, electromyography, goal oriented movements, cervical muscle strength, and joint position error. Further evaluation during the subsequent round was required for temporomandibular screening because the overall rating was “don’t know” (Figure 1d). Participants were also asked in the subsequent round to indicate specific clinical situations for the use of pressure pain thresholds and the cervical slump test since these tests were not considered clinically useful by the majority of participants.




Fig 1a: Distribution of responses for tension-type headache

HFP=head forward posture; ROM=range of motion; PPT=pressure pain thresholds; Cx=cervical; CCFT=cranio-cervical flexion test; JPE=joint position error; mvts=movements; CMD=craniomandibular dysfunction.




Fig 1b: Distribution of responses for migraine headache

HFP=head forward posture; ROM=range of motion; PPT=pressure pain thresholds; Cx=cervical; CCFT=cranio-cervical flexion test; JPE=joint position error; mvts=movements; CMD=craniomandibular dysfunction.




Fig 1c: Distribution of responses for cervicogenic headache

HFP=head forward posture; ROM=range of motion; PPT=pressure pain thresholds; Cx=cervical; CCFT=cranio-cervical flexion test; JPE=joint position error; mvts=movements; CMD=craniomandibular dysfunction.




Figure 1d: Mode values across headache types for all participants
HFP=head forward posture; ROM=range of motion; PPT=pressure pain thresholds; Cx=cervical; CCFT=cranio-cervical flexion test; JPE=joint position error; mvts=movements; CMD=craniomandibular dysfunction.

The following new tests were suggested in round 1 and evaluated in round 2: passive physiological intervertebral movements, combined movements other than flexion-rotation, cranial nerve palpation, muscle lengths tests, thoracic spine screening, headache reproduction and resolution (to sustained palpation), left-right recognition, facial emotional recognition, and other neurodynamic tests.

Results from rounds 2 and 3:

There were no objections to the decision to rate the clinical usefulness of physical examination tests across all headache types rather than individually for specific headache diagnoses. Results indicated that muscle lengths tests were considered “probably not useful” and were thus excluded. Left-right-recognition, facial emotional recognition and other neurodynamic tests received a combined rating of “don’t know” (Figures 2a and b). Table 1 summarises the qualitative responses to clarify clinical situations in which the use of tests was specified that were rated as either “don’t know” or that were not perceived as clinically useful by the majority of participants. Two further tests were suggested: the total tenderness score and the McKenzie classification. The total tenderness score reached a summary score of “probably useful” but did not meet the >50% target of “extremely/probably useful” ratings. The McKenzie tests received an overall rating of “don’t know” and qualitative responses for both tests were subsequently added to Table 1.




Figure 2a: Distribution of responses from rounds 2 and 3

PPIVMs= passive physiological intervertebral movements; Tx=thoracic; L-R=left-right; TTS= total tenderness score.




Figure 2b: Results from the second and third survey rounds. Values in the graph indicate the mode value across participants for each test.

PPIVMs= passive physiological intervertebral movements; Tx=thoracic; L-R=left-right; TTS= total tenderness score.

Table 1: Qualitative responses for clinical indications for tests that received a summary rating of “don’t know” or were supported by <51% of the 17 participants. The number in brackets indicates the number of agreed responses on a statement.

Physical examination test	Might be useful in these specific clinical situations:
Temporo-mandibular screening	<ul style="list-style-type: none"> - orofacial pain or pain with mandibular movement (10) - cervical component suspected (3) - all other tests negative (2) - tension suspected as main pain mechanism (2) - insufficient improvement with the treatment of the cervical spine (2) - to exclude CMD (1)
Left-right-recognition	<ul style="list-style-type: none"> - CGH (1) - central sensitisation (1) - face or neck pain unilateral > 3 months (1)
Facial emotional recognition	<ul style="list-style-type: none"> - orofacial pain (2) - TTH (1)
Other Neurodynamic Tests e.g. upper cervical flexion with bilateral shoulder abduction	<ul style="list-style-type: none"> - to screen for neural tissue mechanosensitivity (3) - cervical spine flexion or retraction is restricted and painful (2) - CGH (2) - trigeminal neuralgia (1) - clinical pattern indicates neurodynamic component (1) - neuropathic pain pattern (1)
Pressure pain thresholds	<ul style="list-style-type: none"> - exclusively for research purposes (6) - to evaluate central sensitisation (6) - in chronic TTH as a re-assessment tool (2) - fibromyalgia suspected (1)
Cervical slump test	<ul style="list-style-type: none"> - CGH in cases where a neurodynamic component was suspected (9) - advanced test, if less aggressive test did not show conclusive response (2) - as a safety tests prior to cervical manipulation (1)
Cranial nerve palpation	<ul style="list-style-type: none"> - only if neuropathic component /cervical neural sensitisation suspected (5) - migraine (2) - no other findings (2) - to determine the cranial nerve sensitivity (2)
Total tenderness score	<ul style="list-style-type: none"> - suspected TMJ involvement (3) - TTH (3) - re-assessment (3) - all patients (1) - suspected central sensitisation (1) - to indicate musculoskeletal involvement (1) - myofascial pain syndrome (1) - if headaches symptoms were consistent with the referral pattern of the affected muscles (1) - to further investigate the relationship between arthrogenic and myogenic structure (1)
McKenzie tests	<ul style="list-style-type: none"> - no evidence (3) - maybe for CGH (2) - only if repeated movements change headache intensity/quality (2) - other tests more informative (1) - disc as a source of pain (1) - posture as a contributing factor (1) - to further identify segmental level (1)

TTH= tension-type headache; CGH= cervicogenic headache; TMJ= temporomandibular joint; CMD= craniomandibular dysfunction.

Discussion

Seventeen international experts in physiotherapy with expertise in headache treatment rated the following 11 tests as the most clinically useful for the evaluation of musculoskeletal dysfunctions in patients with headache: manual joint palpation, the cranio-cervical flexion test, the flexion-rotation test, active range of cervical movement, head forward position, trigger point palpation, muscle tests of the shoulder girdle, passive physiological intervertebral movements, reproduction and resolution of headache symptoms, screening of the thoracic spine, and combined movement tests.

Based on the patient interview and the physiotherapists' clinical reasoning, additional tests might be required or even prioritised. This is particularly so in specific clinical situations such as orofacial pain, or the presence of temporo-mandibular signs where the additional thorough examination of the temporo-mandibular joint and cranial nerve palpation is essential. Another example is the presence of dizziness or light-headedness, where tests discarded from the cluster of tests suggested in this article, such as proprioception tests, might become the most important assessment parameter.

There was a high participation rate (17/20) for all three rounds of the survey. Experts were from various clinical and scientific backgrounds. This strengthens the transferability of the suggested cluster of tests because experts rated the suggested tests for the assessment of musculoskeletal dysfunctions with a background in different patient populations: acute and chronic, patients referred by medical doctors or seen in a primary care setting, patients consulting an orthopaedic or a neurological clinic. Dominantly research-based physiotherapists pursued strong clinical aspect in their headache research and were therefore also well suited to comment on the clinical usefulness of physical examination tests.

The physical examination tests suggested for the first survey round were based on a systematic search of the literature, hence had been formally evaluated in a headache population in at least one publication. The additional tests suggested by the experts as clinically useful tools for the examination of headache patients, that were subsequently included in the final cluster of test, were not all based on evidence from the literature. Reproduction and resolution had previously been shown to differentiate between migraine or TTH and control participants (Watson and Drummond 2012). However, PPIVMs, combined cervical movements and impairments of the thoracic spine have never been published to differentiate between headache populations and headache free participants. Additionally, test properties such as inter-/intra-rater reliability, have only been previously assessed for manual joint palpation (Watson and Trott 1993; Jull et al. 1997, 2007; Van Suijlekom et al. 2000; Zito et al. 2006; Hall et al. 2010), the cranio-cervical

flexion test, (Zito et al. 2006; Jull et al. 2007), the flexion-rotation test, (Ogince et al. 2007; Hall et al. 2008), active range of cervical movement (Van Suijlekom et al. 2000; Fernández-de-Las-Peñas et al. 2006b; Zito et al. 2006; Jull et al. 2007; Bevilaqua-Grossi et al. 2009), head forward position, (Watson and Trott 1993; Fernández-de-Las-Peñas et al. 2006b; Zito et al. 2006), reproduction and resolution of headache symptoms, (Watson and Trott 1993), and trigger point palpation (Fernández-de-Las-Peñas et al. 2006b). Information or data presented to evaluate the sensitivity and specificity of physical examination tests for headache populations is restricted to the flexion-rotation test (Ogince et al. 2007; Hall et al. 2008), manual joint palpation (Zito et al. 2006; Jull et al. 2007), range of cervical movement (Zito et al. 2006), and the craniocervical flexion test (Zito et al. 2006). Nevertheless, the results of this consensus paper are based on the expertise of participants who valued the suggested 11 tests as clinically useful even in the absence of sufficient evidence on test properties. There is a clear need for future research to close this gap between clinical beliefs and scientific evidence to promote a more evidence based approach in clinical physiotherapy.

Physical examination tests for the assessment of musculoskeletal dysfunctions were more frequently considered useful / extremely useful in CGH and least useful for migraine. However no test emerged as being regarded as especially useful for a specific headache type. Given the pathophysiological concepts one could discuss whether there should be a focus on muscle tests for TTH, a focus on the central nervous system in migraine, and a focus on joint dysfunctions for CGH. However, when reflecting on the clinical purpose of such tests, it is clear that one of the main purposes of the physical examination is to identify the presence or absence of cervical musculoskeletal findings and to specify which structures and systems are most affected and are hence the most promising treatment target.

It was the general consensus after round one to evaluate the clinical usefulness across headache types rather than for each headache type individually, even though the ratings for individual headache diagnoses during the first survey round indicated that the clinical usefulness of tests was perceived differently for different headache diagnoses. The majority of experts considered only two physical examination tests useful in the evaluation of musculoskeletal impairments in patients diagnosed with migraine: cranio-cervical flexion test and manual joint palpation. The same tests were identified by an earlier publication as the two tests (combined with restricted cervical movement) that distinguished best between CGH and TTH / migraine (Jull et al. 2007). This is acknowledged by the IHS classification (Headache Classification Committee of the International Headache Society (IHS) 2013) that includes provocation manoeuvres and restriction of cervical movement as clinical signs of CGH. However, according to the IHS classification, the diagnosis CGH is only justified in the presence of a pathological finding in the

cervical spine. These restrictions are the reason why physiotherapists often refer to one of the two alternative classification systems that provide a more functional description of CGH (www.iasp.org; (Sjaastad et al. 1998). This is rather problematic, as studies which are scientifically sound and statistically well done are still not comparable to each other which is a major restriction in the field. For comparability reasons and to support intra- and interprofessional communication, studies evaluating the usefulness of physiotherapeutic interventions for headache should clearly state which classification was used. Moreover, for clinical routine but especially for research purposes it will be mandatory in the future to harmonise the divergence between these classification systems.

Limitations

The internationality of this study and the different clinical and scientific backgrounds of experts might have introduced variability in responses. During the survey process a discussion emerged on the question whether such tests are used to make a diagnosis or to develop a treatment plan. This was not specified during the survey because not all experts had a first contact status and were therefore responsible for making a diagnosis. The suggested cluster of tests can therefore be applied in both situations and used by physiotherapists independent of the first contact status.

Since some of the tests have not been evaluated for sensitivity and specificity as well as reliability and validity, the suggested tests should be used with caution and within a sound clinical reasoning framework.

Conclusion

This is the first international consensus study providing guidance on the most useful physical examination tests for the evaluation of cervical musculoskeletal impairments in patients with headaches. The cluster of tests was based on a comprehensive evaluation of the literature and on the opinion of international experts in physiotherapy. Eleven physical examination tests for the assessment of musculoskeletal dysfunctions were considered useful by the majority of experts for the evaluation of patients with headache. Independent of the diagnosis of migraine, TTH or CGH, we suggest the use of this cluster of tests as a minimum standard for the evaluation of musculoskeletal dysfunction in patients with headache. However, a consensus study, can only reflect the opinion of experts and results should stimulate future discussion and as well as scientific evaluation.

Of note, eight additional tests were considered useful for specific clinical situations. This consensus paves the way for future research to further evaluate the measurement properties in many of these tests and to compare the prevalence of positive findings in patients with different headache types.

References:

- Abboud J, Marchand A-A, Sorra K, Descarreaux M. Musculoskeletal physical outcome measures in individuals with tension-type headache: a scoping review. *Cephalalgia Int J Headache*. 2013 Dec;33(16):1319–36.
- Amiri M, Jull G, Bullock-Saxton J, Darnell R, Lander C. Cervical musculoskeletal impairment in frequent intermittent headache. Part 2: subjects with concurrent headache types. *Cephalalgia Int J Headache*. 2007 Aug;27(8):891–8.
- Antonaci F, Fredriksen TA, Sjaastad O. Cervicogenic headache: clinical presentation, diagnostic criteria, and differential diagnosis. *Curr Pain Headache Rep*. 2001 Aug;5(4):387–92.
- Ashina M, Bendtsen L, Jensen R, Sakai F, Olesen J. Muscle hardness in patients with chronic tension-type headache: relation to actual headache state. *Pain*. 1999 Feb;79(2-3):201–5.
- Ashina S, Bendtsen L, Lyngberg AC, Lipton RB, Hajiyeva N, Jensen R. Prevalence of neck pain in migraine and tension-type headache: A population study. *Cephalalgia Int J Headache*. 2015 Mar;35(3):211–9.
- Becker WJ. Cervicogenic headache: evidence that the neck is a pain generator. *Headache*. 2010 Apr;50(4):699–705.
- Bendtsen L, Fernández-de-la-Peñas C. The role of muscles in tension-type headache. *Curr Pain Headache Rep*. 2011 Dec;15(6):451–8.
- Berg J, Stovner LJ. Cost of migraine and other headaches in Europe. *Eur J Neurol*. 2005 Jun 1;12:59–62.
- Bevilaqua-Grossi D, Pegoretti KS, Goncalves MC, Speciali JG, Bordini CA, Bigal ME. Cervical mobility in women with migraine. *Headache*. 2009 May;49(5):726–31.
- Bloudek LM, Stokes M, Buse DC, Wilcox TK, Lipton RB, Goadsby PJ, et al. Cost of healthcare for patients with migraine in five European countries: results from the International Burden of Migraine Study (IBMS). *J Headache Pain*. 2012 Jul;13(5):361–78.
- Bogduk N. The anatomical basis for cervicogenic headache. *J Manipulative Physiol Ther*. 1992 Jan;15(1):67–70.
- Bogduk N, Govind J. Cervicogenic headache: an assessment of the evidence on clinical diagnosis, invasive tests, and treatment. *Lancet Neurol*. 2009 Oct;8(10):959–68.
- Bovim G. Cervicogenic headache, migraine, and tension-type headache. Pressure-pain threshold measurements. *Pain*. 1992 Nov;51(2):169–73.
- Burton WN, Conti DJ, Chen C-Y, Schultz AB, Edington DW. The economic burden of lost productivity due to migraine headache: a specific worksite analysis. *J Occup Environ Med Am Coll Occup Environ Med*. 2002 Jun;44(6):523–9.
- Calandre EP, Hidalgo J, García-Leiva JM, Rico-Villademoros F. Trigger point evaluation in migraine patients: an indication of peripheral sensitization linked to migraine predisposition? *Eur J Neurol Off J Eur Fed Neurol Soc*. 2006 Mar;13(3):244–9.
- Caspersen N, Hirsvang JR, Kroell L, Jadidi F, Baad-Hansen L, Svensson P, et al. Is There a Relation between Tension-Type Headache, Temporomandibular Disorders and Sleep? *Pain Res Treat*. 2013;2013:845684.
- Chaibi A, Russell MB. Manual therapies for cervicogenic headache: a systematic review. *J Headache Pain*. 2012 Jul;13(5):351–9.
- Chaibi A, Russell MB. Manual therapies for primary chronic headaches: a systematic review of randomized controlled trials. *J Headache Pain*. 2014;15:67.
- Chaibi A, Tuchin PJ, Russell MB. Manual therapies for migraine: a systematic review. *J Headache Pain*. 2011 Apr;12(2):127–33.
- Chiarotto A, Terwee CB, Deyo RA, Boers M, Lin C-WC, Buchbinder R, et al. A core

outcome set for clinical trials on non-specific low back pain: study protocol for the development of a core domain set. *Trials*. 2014;15:511.

Couppé C, Torelli P, Fuglsang-Frederiksen A, Andersen KV, Jensen R. Myofascial trigger points are very prevalent in patients with chronic tension-type headache: a double-blinded controlled study. *Clin J Pain*. 2007 Jan;23(1):23–7.

D’Amico D, Grazzi L, Usai S, Leonardi M, Raggi A. Disability and quality of life in headache: where we are now and where we are heading. *Neurol Sci Off J Ital Neurol Soc Ital Soc Clin Neurophysiol*. 2013 May;34 Suppl 1:S1–5.

Dumas JP, Arsenault AB, Boudreau G, Magnoux E, Lepage Y, Bellavance A, et al. Physical impairments in cervicogenic headache: traumatic vs. nontraumatic onset. *Cephalalgia Int J Headache*. 2001 Nov;21(9):884–93.

Fernández-de-Las-Peñas C, Alonso-Blanco C, Cuadrado ML, Gerwin RD, Pareja JA. Myofascial trigger points and their relationship to headache clinical parameters in chronic tension-type headache. *Headache*. 2006a Sep;46(8):1264–72.

Fernández-de-las-Peñas C, Alonso-Blanco C, Cuadrado ML, Pareja JA. Forward head posture and neck mobility in chronic tension-type headache: a blinded, controlled study. *Cephalalgia Int J Headache*. 2006 Mar;26(3):314–9.

Fernández-de-Las-Peñas C, Coppieters MW, Cuadrado ML, Pareja JA. Patients with chronic tension-type headache demonstrate increased mechano-sensitivity of the supra-orbital nerve. *Headache*. 2008 Apr;48(4):570–7.

Fernández-de-las-Peñas C, Cuadrado ML, Arendt-Nielsen L, Simons DG, Pareja JA. Myofascial trigger points and sensitization: an updated pain model for tension-type headache. *Cephalalgia Int J Headache*. 2007 May;27(5):383–93.

Fernández-de-Las-Peñas C, Cuadrado ML, Pareja JA. Myofascial trigger points, neck mobility and forward head posture in unilateral migraine. *Cephalalgia Int J Headache*. 2006b Sep;26(9):1061–70.

Fernández-de-Las-Peñas C, Cuadrado ML, Pareja JA. Myofascial trigger points, neck mobility, and forward head posture in episodic tension-type headache. *Headache*. 2007 May;47(5):662–72.

Fernández-de-las-Peñas C, Falla D, Arendt-Nielsen L, Farina D. Cervical muscle co-activation in isometric contractions is enhanced in chronic tension-type headache patients. *Cephalalgia Int J Headache*. 2008 Jul;28(7):744–51.

Garrett TR, Youdas JW, Madson TJ. Reliability of measuring forward head posture in a clinical setting. *J Orthop Sports Phys Ther*. 1993 Mar;17(3):155–60.

Giacomini PG, Alessandrini M, Evangelista M, Napolitano B, Lanciani R, Camaioni D. Impaired postural control in patients affected by tension-type headache. *Eur J Pain Lond Engl*. 2004 Dec;8(6):579–83.

Glaros AG, Urban D, Locke J. Headache and temporomandibular disorders: evidence for diagnostic and behavioural overlap. *Cephalalgia Int J Headache*. 2007 Jun;27(6):542–9.

Hall T, Briffa K, Hopper D, Robinson K. Reliability of manual examination and frequency of symptomatic cervical motion segment dysfunction in cervicogenic headache. *Man Ther*. 2010 Dec;15(6):542–6.

Hall TM, Robinson KW, Fujinawa O, Akasaka K, Pyne EA. Intertester reliability and diagnostic validity of the cervical flexion-rotation test. *J Manipulative Physiol Ther*. 2008 May;31(4):293–300.

Hall T, Robinson K. The flexion-rotation test and active cervical mobility--a comparative measurement study in cervicogenic headache. *Man Ther*. 2004 Nov;9(4):197–202.

Haque B, Rahman KM, Hoque A, Hasan ATMH, Chowdhury RN, Khan SU, et al. Precipitating and relieving factors of migraine versus tension type headache. *BMC Neurol*. 2012;12:82.

Harry von Piekartz. Temporomandibular Disorders: Neuromusculoskeletal Assessment and Management. *Grieves Mod Musculoskelet Physiother* 4th Ed. 2015.

Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs*. 2000 Oct;32(4):1008–15.

Hatch JP, Moore PJ, Cyr-Provost M, Boutros NN, Seleshi E, Borcharding S. The use of electromyography and muscle palpation in the diagnosis of tension-type headache with and without pericranial muscle involvement. *Pain*. 1992 May;49(2):175–8.

Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia Int J Headache*. 2013 Jul;33(9):629–808.

Hsu and Sandford. The Delphi technique: making sense of consensus. *Pract Assess Res Eval*. 2007;12:1–8.

Jensen R. Mechanisms of spontaneous tension-type headaches: an analysis of tenderness, pain thresholds and EMG. *Pain*. 1996 Feb;64(2):251–6.

Jensen R, Rasmussen BK. Muscular disorders in tension-type headache. *Cephalalgia Int J Headache*. 1996 Apr;16(2):97–103.

Jensen R, Rasmussen BK, Pedersen B, Olesen J. Muscle tenderness and pressure pain thresholds in headache. A population study. *Pain*. 1993 Feb;52(2):193–9.

Jull G, Amiri M, Bullock-Saxton J, Darnell R, Lander C. Cervical musculoskeletal impairment in frequent intermittent headache. Part 1: Subjects with single headaches. *Cephalalgia Int J Headache*. 2007 Jul;27(7):793–802.

Jull G, Barrett C, Magee R, Ho P. Further clinical clarification of the muscle dysfunction in cervical headache. *Cephalalgia Int J Headache*. 1999 Apr;19(3):179–85.

Jull G, Zito G, Trott P, Potter H, Shirley D. Inter-examiner reliability to detect painful upper cervical joint dysfunction. *Aust J Physiother*. 1997;43(2):125–9.

Kidd RF, Nelson R. Musculoskeletal dysfunction of the neck in migraine and tension headache. *Headache*. 1993 Dec;33(10):566–9.

Kristoffersen ES, Grande RB, Aaseth K, Lundqvist C, Russell MB. Management of primary chronic headache in the general population: the Akershus study of chronic headache. *J Headache Pain*. 2012 Mar;13(2):113–20.

Langemark M, Olesen J. Pericranial tenderness in tension headache. A blind, controlled study. *Cephalalgia Int J Headache*. 1987 Dec;7(4):249–55.

Lanteri-Minet M. Economic burden and costs of chronic migraine. *Curr Pain Headache Rep*. 2014 Jan;18(1):385.

Leone M, D'Amico D, Grazzi L, Attanasio A, Bussone G. Cervicogenic headache: a critical review of the current diagnostic criteria. *Pain*. 1998 Oct;78(1):1–5.

Lipchik GL, Holroyd KA, France CR, Kvaal SA, Segal D, Cordingley GE, et al. Central and peripheral mechanisms in chronic tension-type headache. *Pain*. 1996 Mar;64(3):467–75.

Lous I, Olesen J. Evaluation of pericranial tenderness and oral function in patients with common migraine, muscle contraction headache and “combination headache.” *Pain*. 1982 Apr;12(4):385–93.

Marchand A-A, Cantin V, Murphy B, Stern P, Descarreaux M. Is performance in goal oriented head movements altered in patients with tension type headache? *BMC Musculoskelet Disord*. 2014;15:179.

Melo C-E-B, Oliveira J-L-G, Jesus A-C-F, Maia M-LM, de Santana J-C-V, Andrade L-S-O, et al. Temporomandibular disorders dysfunction in headache patients. *Med Oral Patol Oral Cir Bucal*. 2012 Nov;17(6):e1042–6.

Mørk H, Ashina M, Bendtsen L, Olesen J, Jensen R. Induction of prolonged tenderness in patients with tension-type headache by means of a new experimental model of myofascial pain. *Eur J Neurol Off J Eur Fed Neurol Soc*. 2003 May;10(3):249–56.

Murphy MK, Black NA, Lamping DL, McKee CM, Sanderson CF, Askham J, et al. Consensus development methods, and their use in clinical guideline development. *Health Technol Assess Winch Engl*. 1998;2(3):i – iv, 1–88.

Neufeld JD, Holroyd KA, Lipchik GL. Dynamic assessment of abnormalities in central pain transmission and modulation in tension-type headache sufferers. *Headache*. 2000 Feb;40(2):142–51.

Ogince M, Hall T, Robinson K, Blackmore AM. The diagnostic validity of the cervical flexion-rotation test in C1/2-related cervicogenic headache. *Man Ther*. 2007 Aug;12(3):256–62.

Peddireddy A, Wang K, Svensson P, Arendt-Nielsen L. Stretch reflex and pressure pain thresholds in chronic tension-type headache patients and healthy controls. *Cephalalgia Int J Headache*. 2009 May;29(5):556–65.

Pöllmann W, Keidel M, Pfaffenrath V. Headache and the cervical spine: a critical review. *Cephalalgia Int J Headache*. 1997 Dec;17(8):801–16.

Pradalier A, Auray J-P, El Hasnaoui A, Alzahouri K, Dartigues J-F, Duru G, et al. Economic impact of migraine and other episodic headaches in France: data from the GRIM2000 study. *PharmacoEconomics*. 2004;22(15):985–99.

Raggi A, Leonardi M, Bussone G, D'Amico D. A 3-month analysis of disability, quality of life, and disease course in patients with migraine. *Headache*. 2013 Feb;53(2):297–309.

Robertson BA, Morris ME. The role of cervical dysfunction in migraine: a systematic review. *Cephalalgia Int J Headache*. 2008 May;28(5):474–83.

Sandrini G, Rossi P, Milanov I, Serrao M, Cecchini AP, Nappi G. Abnormal modulatory influence of diffuse noxious inhibitory controls in migraine and chronic tension-type headache patients. *Cephalalgia Int J Headache*. 2006 Jul;26(7):782–9.

Sand T, Zwart JA, Helde G, Bovim G. The reproducibility of cephalic pain pressure thresholds in control subjects and headache patients. *Cephalalgia Int J Headache*. 1997 Nov;17(7):748–55.

Sinha IP, Smyth RL, Williamson PR. Using the Delphi technique to determine which outcomes to measure in clinical trials: recommendations for the future based on a systematic review of existing studies. *PLoS Med*. 2011;8(1):e1000393.

Sjaastad O, Bakketeig LS. Prevalence of cervicogenic headache: Vågå study of headache epidemiology. *Acta Neurol Scand*. 2008 Mar 1;117(3):173–80.

Sjaastad O, Fredriksen TA, Pfaffenrath V. Cervicogenic headache: diagnostic criteria. The Cervicogenic Headache International Study Group. *Headache*. 1998 Jun;38(6):442–5.

Sohn J-H, Choi H-C, Jun A-Y. Differential patterns of muscle modification in women with episodic and chronic tension-type headache revealed using surface electromyographic analysis. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol*. 2013 Feb;23(1):110–7.

Sohn J-H, Choi H-C, Lee S-M, Jun A-Y. Differences in cervical musculoskeletal impairment between episodic and chronic tension-type headache. *Cephalalgia Int J Headache*. 2010 Dec;30(12):1514–23.

Steiner TJ, Stovner LJ, Katsarava Z, Lainez JM, Lampl C, Lantéri-Minet M, et al. The impact of headache in Europe: principal results of the Eurolight project. *J Headache Pain*. 2014;15:31.

Stuginski-Barbosa J, Macedo HR, Bigal ME, Speciali JG. Signs of temporomandibular disorders in migraine patients: a prospective, controlled study. *Clin J Pain*. 2010 Jun;26(5):418–21.

Van Suijlekom HA, De Vet HC, Van Den Berg SG, Weber WE. Interobserver reliability in physical examination of the cervical spine in patients with headache. *Headache*. 2000 Aug;40(7):581–6.

Tali D, Menahem I, Vered E, Kalichman L. Upper cervical mobility, posture and myofascial trigger points in subjects with episodic migraine: Case-control study. *J Bodyw Mov Ther.* 2014 Oct;18(4):569–75.

Uthaikhup S, Sterling M, Jull G. Cervical musculoskeletal impairment is common in elders with headache. *Man Ther.* 2009 Dec;14(6):636–41.

Vincent MB. Headache and Neck. *Curr Pain Headache Rep.* 2011 Apr 5;15(4):324–31.

Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet.* 2012 Dec 15;380(9859):2163–96.

Watson DH, Drummond PD. Head pain referral during examination of the neck in migraine and tension-type headache. *Headache.* 2012 Sep;52(8):1226–35.

Watson DH, Trott PH. Cervical headache: an investigation of natural head posture and upper cervical flexor muscle performance. *Cephalalgia Int J Headache.* 1993 Aug;13(4):272–84; discussion 232.

Zito G, Jull G, Story I. Clinical tests of musculoskeletal dysfunction in the diagnosis of cervicogenic headache. *Man Ther.* 2006 May;11(2):118–29.

Zwart JA. Neck mobility in different headache disorders. *Headache.* 1997 Jan;37(1):6–11.

Figure Captions

Fig 1a: Distribution of responses for tension-type headache

HFP=head forward posture; ROM=range of motion; PPT=pressure pain thresholds; Cx=cervical; CCFT=cranio-cervical flexion test; JPE=joint position error; mvts=movements; CMD=craniomandibular dysfunction.




Fig 1b: Distribution of responses for migraine headache

HFP=head forward posture; ROM=range of motion; PPT=pressure pain thresholds; Cx=cervical; CCFT=cranio-cervical flexion test; JPE=joint position error; mvts=movements; CMD=craniomandibular dysfunction.




Fig 1c: Distribution of responses for cervicogenic headache

HFP=head forward posture; ROM=range of motion; PPT=pressure pain thresholds; Cx=cervical; CCFT=cranio-cervical flexion test; JPE=joint position error; mvts=movements; CMD=craniomandibular dysfunction.




Figure 1d: Mode values across headache types for all participants

HFP=head forward posture; ROM=range of motion; PPT=pressure pain thresholds; Cx=cervical; CCFT=cranio-cervical flexion test; JPE=joint position error; mvts=movements; CMD=craniomandibular dysfunction.




Figure 2a: Distribution of responses from rounds 2 and 3

PPIVMs= passive physiological intervertebral movements; Tx=thoracic; L-R=left-right; TTS= total tenderness score.




Figure 2b: Results from the second and third survey rounds. Values in the graph indicate the mode value across participants for each test.

PPIVMs= passive physiological intervertebral movements; Tx=thoracic; L-R=left-right; TTS= total tenderness score.

